

Case Number:	CM13-0016289		
Date Assigned:	03/03/2014	Date of Injury:	02/21/2002
Decision Date:	04/14/2014	UR Denial Date:	08/11/2013
Priority:	Standard	Application Received:	08/26/2013

HOW THE IMR FINAL DETERMINATION WAS MADE

MAXIMUS Federal Services sent the complete case file to an expert reviewer. He/she has no affiliation with the employer, employee, providers or the claims administrator. The expert reviewer is Board Certified in Family Medicine, has a subspecialty in Preventive Medicine and is licensed to practice in California. He/she has been in active clinical practice for more than five years and is currently working at least 24 hours a week in active practice. The expert reviewer was selected based on his/her clinical experience, education, background, and expertise in the same or similar specialties that evaluate and/or treat the medical condition and disputed items/services. He/she is familiar with governing laws and regulations, including the strength of evidence hierarchy that applies to Independent Medical Review determinations.

CLINICAL CASE SUMMARY

The expert reviewer developed the following clinical case summary based on a review of the case file, including all medical records:

This is a 47-year-old male claimant who sustained an injury on February 21, 2002 resulting in chronic neck, back, shoulder and lower extremity pain. He had developed radiculopathy for which he had lumbar spine surgery. For several years he has been taking Soma for muscle spasms. Examination note in July 2013 indicated the claimant had eight out of 10 pain without medications and four out of 10 pain with medications. He had been on Norco and Soma. At the time of his examination was notable for muscle tenderness in the paraspinal region as well as limited range of motion any positive leg raise. He was recommended to continue this soma.

IMR ISSUES, DECISIONS AND RATIONALES

The Final Determination was based on decisions for the disputed items/services set forth below:

SOMA 350MG, #60: Upheld

Claims Administrator guideline: Decision based on MTUS Chronic Pain Treatment Guidelines Page(s): 63-66.

MAXIMUS guideline: Decision based on MTUS Chronic Pain Treatment Guidelines Carsiprodolol, Page(s): 29.

Decision rationale: Not recommended. This medication is not indicated for long-term use. Carisoprodol is a commonly prescribed, centrally acting skeletal muscle relaxant whose primary active metabolite is meprobamate (a schedule-IV controlled substance). Final Determination

Letter for IMR Case Number [REDACTED] Carisoprodol is now scheduled in several states but not on a federal level. It has been suggested that the main effect is due to generalized sedation and treatment of anxiety. Abuse has been noted for sedative and relaxant effects. In regular abusers the main concern is the accumulation of meprobamate. Carisoprodol abuse has also been noted in order to augment or alter effects of other drugs. This includes the following: (1) increasing sedation of benzodiazepines or alcohol; (2) use to prevent side effects of cocaine; (3) use with tramadol to produce relaxation and euphoria; (4) as a combination with hydrocodone, an effect that some abusers claim is similar to heroin (referred to as a "Las Vegas Cocktail"); & (5) as a combination with codeine (referred to as "Soma Coma"). (Reeves, 1999) (Reeves, 2001) (Reeves, 2008) (Schears, 2004) There was a 300% increase in numbers of emergency room episodes related to carisoprodol from 1994 to 2005. (DHSS, 2005) Intoxication appears to include subdued consciousness, decreased cognitive function, and abnormalities of the eyes, vestibular function, appearance, gait and motor function. Intoxication includes the effects of both carisoprodol and meprobamate, both of which act on different neurotransmitters. (Bramness, 2007) (Bramness, 2004) A withdrawal syndrome has been documented that consists of insomnia, vomiting, tremors, muscle twitching, anxiety, and ataxia when abrupt discontinuation of large doses occurs. This is similar to withdrawal from meprobamate. (Reeves, 2007) (Reeves, 2004) There is little research in terms of weaning of high dose carisoprodol and there is no standard treatment regimen for patients with known dependence. Most treatment includes treatment for symptomatic complaints of withdrawal. Another option is to switch to phenobarbital to prevent withdrawal with subsequent tapering. A maximum dose of phenobarbital is 500 mg/day and the taper is 30 mg/day with a slower taper in an outpatient setting. Tapering should be individualized for each patient. (Boothby, 2003) For more information and references, see Muscle relaxants. See also Weaning of medications.