

Case Number:	CM13-0025788		
Date Assigned:	11/20/2013	Date of Injury:	01/01/2000
Decision Date:	01/14/2014	UR Denial Date:	09/04/2013
Priority:	Standard	Application Received:	09/18/2013

HOW THE IMR FINAL DETERMINATION WAS MADE

MAXIMUS Federal Services sent the complete case file to a physician reviewer. He/she has no affiliation with the employer, employee, providers or the claims administrator. The physician reviewer is Board Certified in Family Medicine, has a subspecialty in Preventive Medicine and is licensed to practice in California. He/she has been in active clinical practice for more than five years and is currently working at least 24 hours a week in active practice. The physician reviewer was selected based on his/her clinical experience, education, background, and expertise in the same or similar specialties that evaluate and/or treat the medical condition and disputed items/services. He/she is familiar with governing laws and regulations, including the strength of evidence hierarchy that applies to Independent Medical Review determinations.

CLINICAL CASE SUMMARY

The expert reviewer developed the following clinical case summary based on a review of the case file, including all medical records:

This is a 59 yr old male claimant that sustained a work injury in April 2006 resulting in neck pain that radiates to his lower extremities. He has 10/10 pain, constantly falls, has difficulty sleeping and has minimal improvement on anti-inflammatories and therapy. He has been using Vicodin for several months and Dendracin topically for pain. A urine toxicology screen in February 2013 was consistent with the medications he was taking and he had no signs of abuse.

IMR ISSUES, DECISIONS AND RATIONALES

The Final Determination was based on decisions for the disputed items/services set forth below:

Retrospective request for 1 urine drug screen: Upheld

Claims Administrator guideline: Decision based on MTUS Chronic Pain Treatment Guidelines.

MAXIMUS guideline: Decision based on MTUS Chronic Pain Treatment Guidelines Section 9792.20 - 9792.26. Decision based on Non-MTUS Citation Official Disability Guidelines (ODG) Urine Toxicology..

Decision rationale: According to the California MTUS Chronic Pain Treatment Guidelines, urine toxicology screen is used to assess presence of illicit drugs or to monitor adherence to prescription medication program. There's no documentation from the provider to suggest that there was illicit drug use or noncompliance. There were no prior urine drug screen results that

indicated noncompliance, substance-abuse or other inappropriate activity. Furthermore screening for addiction risk should be performed with questionnaires such as the Cage, Skinner trauma, Opioid Risk Tools, etc. Such screening tests were also not indicated in the documentation. The ODG guidelines on Urine Toxicology screening state the following: Indications for UDT: At the onset of treatment: (1) UDT is recommended at the onset of treatment of a new patient who is already receiving a controlled substance or when chronic opioid management is considered. Urine drug testing is not generally recommended in acute treatment settings (i.e. when opioids are required for nociceptive pain). (2) In cases in which the patient asks for a specific drug. This is particularly the case if this drug has high abuse potential, the patient refuses other drug treatment and/or changes in scheduled drugs, or refuses generic drug substitution. (3) If the patient has a positive or "at risk" addiction screen on evaluation. This may also include evidence of a history of comorbid psychiatric disorder such as depression, anxiety, bipolar disorder, and/or personality disorder (4) If aberrant behavior or misuse is suspected and/or detected. Ongoing monitoring: (1) If a patient has evidence of a "high risk" of addiction (including evidence of a comorbid psychiatric disorder (such as depression, anxiety, attention-deficit disorder, obsessive-compulsive disorder, bipolar disorder, and/or schizophrenia), has a history of aberrant behavior, personal or family history of substance dependence (addiction), or a personal history of sexual or physical trauma, ongoing urine drug testing is indicated as an adjunct to monitoring along with clinical exams and pill counts (2) If dose increases are not decreasing pain and increasing function, consideration of UDT should be made to aid in evaluating medication compliance and adherence. Based on the above references and clinical history a urine toxicology screen is not medically necessary.