

<b>Case Number:</b>	CM13-0043930		
<b>Date Assigned:</b>	03/24/2014	<b>Date of Injury:</b>	07/09/2008
<b>Decision Date:</b>	06/30/2014	<b>UR Denial Date:</b>	10/21/2013
<b>Priority:</b>	Standard	<b>Application Received:</b>	10/31/2013

### HOW THE IMR FINAL DETERMINATION WAS MADE

MAXIMUS Federal Services sent the complete case file to an expert reviewer. He/she has no affiliation with the employer, employee, providers or the claims administrator. The expert reviewer is Board Certified in Neurology, has a subspecialty in Neuromuscular Medicine and is licensed to practice in California. He/she has been in active clinical practice for more than five years and is currently working at least 24 hours a week in active practice. The expert reviewer was selected based on his/her clinical experience, education, background, and expertise in the same or similar specialties that evaluate and/or treat the medical condition and disputed items/services. He/she is familiar with governing laws and regulations, including the strength of evidence hierarchy that applies to Independent Medical Review determinations.

### CLINICAL CASE SUMMARY

The expert reviewer developed the following clinical case summary based on a review of the case file, including all medical records:

██████████ is 58 year old man who sustained a work related injury on July 9 2008. Subsequently, he developed a chronic back pain and bilateral knee pain. According to a note dated on October 7 2013, the patient still have pain 3 to 7/10 intensity. The patient was treated with physical therapy, pain medications including Norco, Gabapentin and Zolpidem. The provider requested authorization to use Norco and Oxycodone.

### IMR ISSUES, DECISIONS AND RATIONALES

The Final Determination was based on decisions for the disputed items/services set forth below:

**NORCO 7.5/325 MG #120 WITH THREE REFILLS:** Upheld

**Claims Administrator guideline:** The Claims Administrator did not base their decision on the MTUS. Decision based on Non-MTUS Citation MTUS: CHRONIC PAIN MEDICAL TREATMENT GUIDELINES, ,

**MAXIMUS guideline:** Decision based on MTUS Chronic Pain Treatment Guidelines Chronic Pain Medical Treatment Guidelines < Criteria for use of opioids Page(s): 179.

**Decision rationale:** According to MTUS guidelines, Norco (Hydrocodone/Acetaminophen) is a synthetic opioid indicated for the pain management but not recommended as a first line oral analgesic. In addition and according to MTUS guidelines, ongoing use of opioids should follow specific rules: <(a) Prescriptions from a single practitioner taken as directed, and all

prescriptions from a single pharmacy. (b) The lowest possible dose should be prescribed to improve pain and function. (c) Office: Ongoing review and documentation of pain relief, functional status, appropriate medication use, and side effects. Pain assessment should include: current pain; the least reported pain over the period since last assessment; average pain; intensity of pain after taking the opioid; how long it takes for pain relief; and how long pain relief lasts. Satisfactory response to treatment may be indicated by the patient's decreased pain, increased level of function, or improved quality of life. Information from family members or other caregivers should be considered in determining the patient's response to treatment. The 4 A's for Ongoing Monitoring: Four domains have been proposed as most relevant for ongoing monitoring of chronic pain patients on opioids: pain relief, side effects, physical and psychosocial functioning, and the occurrence of any potentially aberrant (or non adherent) drug-related behaviors. These domains have been summarized as the "4 A's" (analgesia, activities of daily living, adverse side effects, and aberrant drug taking behaviors). The monitoring of these outcomes over time should affect therapeutic decisions and provide a framework> There is no clear evidence of objective and recent functional and pain improvement with previous use of opioids (Norco). There no clear documentation of the efficacy/safety of previous use of Norco. There is no recent evidence of objective monitoring of compliance of the patient with his medications. There is no clear justification for the need to continue the use of Norco. Therefore, the prescription of NORCO 7.5/325 MG #120 WITH THREE REFILLS is not medically necessary at this time

**OXYCODONE 10 MG #60 WITH THREE REFILLS:** Upheld

**Claims Administrator guideline:** The Claims Administrator did not base their decision on the MTUS. Decision based on Non-MTUS Citation MTUS: CHRONIC PAIN MEDICAL TREATMENT GUIDELINES, ,

**MAXIMUS guideline:** Decision based on MTUS Chronic Pain Treatment Guidelines Chronic Pain Medical Treatment Guidelines < Opioids Page(s): 75-81.

**Decision rationale:** According to MTUS guidelines, Oxycodone as well as other short acting opioids are indicated for intermittent or breakthrough pain (page 75). It can be used in acute post operative pain. It is not recommended for chronic pain of longterm use as prescribed in this case. In addition and according to MTUS guidelines, ongoing use of opioids should follow specific rules: <(a) Prescriptions from a single practitioner taken as directed, and all prescriptions from a single pharmacy. (b) The lowest possible dose should be prescribed to improve pain and function. (c) Office: Ongoing review and documentation of pain relief, functional status, appropriate medication use, and side effects. Pain assessment should include: current pain; the least reported pain over the period since last assessment; average pain; intensity of pain after taking the opioid; how long it takes for pain relief; and how long pain relief lasts. Satisfactory response to treatment may be indicated by the patient's decreased pain, increased level of function, or improved quality of life. Information from family members or other caregivers should be considered in determining the patient's response to treatment. The 4 A's for Ongoing Monitoring: Four domains have been proposed as most relevant for ongoing monitoring of chronic pain patients on opioids: pain relief, side effects, physical and psychosocial functioning, and the occurrence of any potentially aberrant (or non adherent) drug-related behaviors. These

domains have been summarized as the "4 A's" (analgesia, activities of daily living, adverse side effects, and aberrant drug taking behaviors). The monitoring of these outcomes over time should affect therapeutic decisions and provide a framework> There is no documentation that the patient have pain breakthrough. In addition, there is no rational for combining Norco with Oxycodone. There is no rational for a continuous and chronic use of Oxycodone. Therefore, the prescription of OXYCODONE 10 MG #60 WITH THREE REFILLS is not medically necessary at this time.