

<b>Case Number:</b>	CM13-0047998		
<b>Date Assigned:</b>	04/30/2014	<b>Date of Injury:</b>	06/17/2013
<b>Decision Date:</b>	07/08/2014	<b>UR Denial Date:</b>	10/24/2013
<b>Priority:</b>	Standard	<b>Application Received:</b>	11/04/2013

### HOW THE IMR FINAL DETERMINATION WAS MADE

MAXIMUS Federal Services sent the complete case file to an expert reviewer. He/she has no affiliation with the employer, employee, providers or the claims administrator. The expert reviewer is Board Certified in Neurology, has a subspecialty in Neuromuscular Medicine, and is licensed to practice in California. He/she has been in active clinical practice for more than five years and is currently working at least 24 hours a week in active practice. The expert reviewer was selected based on his/her clinical experience, education, background, and expertise in the same or similar specialties that evaluate and/or treat the medical condition and disputed items/services. He/she is familiar with governing laws and regulations, including the strength of evidence hierarchy that applies to Independent Medical Review determinations.

### CLINICAL CASE SUMMARY

The expert reviewer developed the following clinical case summary based on a review of the case file, including all medical records:

██████████ is a 53-year-old who sustained a work related injury on June 17 2013. Subsequently, she developed a chronic lower extremities pain. According to a note dated on October 17 2013, the patient reported left ankle pain. Her physical examination showed left ankle tenderness with reduced range of motion and weakness of the left ankle. The provider requested authorization for additional 6 sessions of physical therapy.

### IMR ISSUES, DECISIONS AND RATIONALES

The Final Determination was based on decisions for the disputed items/services set forth below:

#### **6 SESSIONS OF PHYSICAL THERAPY FOR THE LEFT FOOT/ANKLE: Upheld**

**Claims Administrator guideline:** Decision based on MTUS ACOEM Chapter 14 Ankle and Foot Complaints.

**MAXIMUS guideline:** Decision based on MTUS Chronic Pain Treatment Guidelines PHYSICAL MEDICINE Page(s): 98.

**Decision rationale:** According to the Chronic Pain Medical Treatment Guidelines, Physical Medicine is recommended as indicated below. Passive therapy (those treatment modalities that do not require energy expenditure on the part of the patient) can provide short term relief during the early phases of pain treatment and are directed at controlling symptoms such as pain, inflammation and swelling and to improve the rate of healing soft tissue injuries. They can be

used sparingly with active therapies to help control swelling, pain and inflammation during the rehabilitation process. Active therapy is based on the philosophy that therapeutic exercise and/or activity are beneficial for restoring flexibility, strength, endurance, function, range of motion, and can alleviate discomfort. Active therapy requires an internal effort by the individual to complete a specific exercise or task. This form of therapy may require supervision from a therapist or medical provider such as verbal, visual and/or tactile instruction(s). Patients are instructed and expected to continue active therapies at home as an extension of the treatment process in order to maintain improvement levels. Home exercise can include exercise with or without mechanical assistance or resistance and functional activities with assistive devices. (Colorado, 2002) (Airaksinen, 2006) Patient-specific hand therapy is very important in reducing swelling, decreasing pain, and improving range of motion in CRPS. (Li, 2005) The use of active treatment modalities (e.g., exercise, education, activity modification) instead of passive treatments is associated with substantially better clinical outcomes. In a large case series of patients with low back pain treated by physical therapists, those adhering to guidelines for active rather than passive treatments incurred fewer treatment visits, cost less, and had less pain and less disability. The overall success rates were 64.7% among those adhering to the active treatment recommendations versus 36.5% for passive treatment. (Fritz, 2007). Although the patient still has residual left ankle dysfunction, there is no clear justification for the addition of six sessions of physical therapy. Therefore the request for six sessions of physical therapy for the left foot/ankle is not medically necessary or appropriate.