

Case Number:	CM13-0059464		
Date Assigned:	12/30/2013	Date of Injury:	05/16/2012
Decision Date:	11/05/2014	UR Denial Date:	11/26/2013
Priority:	Standard	Application Received:	12/02/2013

HOW THE IMR FINAL DETERMINATION WAS MADE

MAXIMUS Federal Services sent the complete case file to an expert reviewer. He/she has no affiliation with the employer, employee, providers or the claims administrator. The expert reviewer is Board Certified in Anesthesiology, has a subspecialty in Pain Medicine and is licensed to practice in Massachusetts. He/she has been in active clinical practice for more than five years and is currently working at least 24 hours a week in active practice. The expert reviewer was selected based on his/her clinical experience, education, background, and expertise in the same or similar specialties that evaluate and/or treat the medical condition and disputed items/services. He/she is familiar with governing laws and regulations, including the strength of evidence hierarchy that applies to Independent Medical Review determinations.

CLINICAL CASE SUMMARY

The expert reviewer developed the following clinical case summary based on a review of the case file, including all medical records:

According to the documents available for review, the patient is a 35-year-old male. The date of injury is May 16, 2012. The patient sustained an injury to the left hip. The specific mechanism of injury was not fully elaborated on in the notes available for review. The patient underwent left hip arthroscopy on April 12, 2013. The patient currently complains of pain in the left hip and left groin worse with movement. The patient has undergone 28 prior physical therapy visits. A request for continued postoperative physical therapy two times a week for six weeks to the left was denied.

IMR ISSUES, DECISIONS AND RATIONALES

The Final Determination was based on decisions for the disputed items/services set forth below:

CONTINUED POSTOPERATIVE PHYSICAL THERAPY 2 TIMES A WEEK FOR 6 WEEKS TO THE LEFT HIP QTY: 12.00: Upheld

Claims Administrator guideline: Decision based on MTUS Postsurgical Treatment Guidelines Page(s): 23.

MAXIMUS guideline: Decision based on MTUS Chronic Pain Treatment Guidelines Physical Medicine Page(s): 98.

Decision rationale: Per MTUS Physical medicine is recommended as indicated below. Passive therapy (those treatment modalities that do not require energy expenditure on the part of the

patient) can provide short term relief during the early phases of pain treatment and are directed at controlling symptoms such as pain, inflammation and swelling and to improve the rate of healing soft tissue injuries. They can be used sparingly with active therapies to help control swelling, pain and inflammation during the rehabilitation process. Active therapy is based on the philosophy that therapeutic exercise and/or activity are beneficial for restoring flexibility, strength, endurance, function, range of motion, and can alleviate discomfort. Active therapy requires an internal effort by the individual to complete a specific exercise or task. This form of therapy may require supervision from a therapist or medical provider such as verbal, visual and/or tactile instruction(s). Patients are instructed and expected to continue active therapies at home as an extension of the treatment process in order to maintain improvement levels. Home exercise can include exercise with or without mechanical assistance or resistance and functional activities with assistive devices. Patient-specific hand therapy is very important in reducing swelling, decreasing pain, and improving range of motion in CRPS. (Li, 2005) The use of active treatment modalities (e.g., exercise, education, activity modification) instead of passive treatments is associated with substantially better clinical outcomes. In a large case series of patients with low back pain treated by physical therapists, those adhering to guidelines for active rather than passive treatments incurred fewer treatment visits, cost less, and had less pain and less disability. The overall success rates were 64.7% among those adhering to the active treatment recommendations versus 36.5% for passive treatment. Physical Medicine Guidelines Allow for fading of treatment frequency (from up to 3 visits per week to 1 or less), plus active self-directed home Physical Medicine. Myalgia and myositis, unspecified: 9-10 visits over 8 weeks; Neuralgia, neuritis, and radiculitis, unspecified 8-10 visits over 4 weeks; Reflex sympathetic dystrophy (CRPS): 24 visits over 16 weeks. According to the documents available for review, the patient has previously undergone 28 prior sessions of physical therapy. Further therapy sessions as requested would be in contrast to the recommendations as outlined in the MTUS. Therefore at this time the requirements for treatment have not been met and medical necessity has not been established.