

Case Number:	CM14-0105323		
Date Assigned:	09/24/2014	Date of Injury:	05/01/2000
Decision Date:	11/28/2014	UR Denial Date:	06/03/2014
Priority:	Standard	Application Received:	07/07/2014

HOW THE IMR FINAL DETERMINATION WAS MADE

MAXIMUS Federal Services sent the complete case file to an expert reviewer. He/she has no affiliation with the employer, employee, providers or the claims administrator. The expert reviewer is Board Certified in Physical Medicine and Rehabilitation, has a subspecialty in Pain Medicine and is licensed to practice in California. He/she has been in active clinical practice for more than five years and is currently working at least 24 hours a week in active practice. The expert reviewer was selected based on his/her clinical experience, education, background, and expertise in the same or similar specialties that evaluate and/or treat the medical condition and disputed items/services. He/she is familiar with governing laws and regulations, including the strength of evidence hierarchy that applies to Independent Medical Review determinations.

CLINICAL CASE SUMMARY

The expert reviewer developed the following clinical case summary based on a review of the case file, including all medical records:

This is a patient with a date of injury of 5/1/00. A utilization review determination dated 6/3/14 recommends non-certification of Norco, Soma, lorazepam, and topical flurbiprofen. 4/1/14 medical report identifies lumbosacral pain radiating down both legs with numbness and tingling. On exam, there is tenderness, decreased ROM, and positive SLR bilaterally at 20 degrees. Recommendations include continued medications and right L3-S1 microdiscectomy.

IMR ISSUES, DECISIONS AND RATIONALES

The Final Determination was based on decisions for the disputed items/services set forth below:

Norco 10/325 mg # 120 refills 4: Upheld

Claims Administrator guideline: Decision based on MTUS Chronic Pain Treatment Guidelines Page(s): 91.

MAXIMUS guideline: Decision based on MTUS Chronic Pain Treatment Guidelines Page(s): 44, 47, 75-79, 120 of 127.

Decision rationale: Regarding the request for Norco, California Pain Medical Treatment Guidelines note that it is an opiate pain medication. Due to high abuse potential, close follow-up is recommended with documentation of analgesic effect, objective functional improvement, side effects, and discussion regarding any aberrant use. Guidelines go on to recommend discontinuing opioids if there is no documentation of improved function and pain. Within the documentation

available for review, there is no indication that the medication is improving the patient's function or pain (in terms of specific examples of functional improvement and percent reduction in pain or reduced NRS), no documentation regarding side effects, and no discussion regarding aberrant use. As such, there is no clear indication for ongoing use of the medication. Opioids should not be abruptly discontinued, but unfortunately, there is no provision to modify the current request to allow tapering. In light of the above issues, the currently requested Norco is not medically necessary.

Soma 350mg # 90 refills 4: Upheld

Claims Administrator guideline: Decision based on MTUS Chronic Pain Treatment Guidelines Page(s): 29.

MAXIMUS guideline: Decision based on MTUS Chronic Pain Treatment Guidelines Page(s): 63-66 of 127.

Decision rationale: Regarding the request for Soma, Chronic Pain Medical Treatment Guidelines support the use of non-sedating muscle relaxants to be used with caution as a 2nd line option for the short-term treatment of acute exacerbations of pain. Within the documentation available for review, there is no identification of a specific analgesic benefit or objective functional improvement as a result of the medication. Additionally, it does not appear that this medication is being prescribed for the short-term treatment of an acute exacerbation, as recommended by guidelines. In the absence of such documentation, the currently requested Soma is not medically necessary.

Lorazepam 2mg # 90 x4: Upheld

Claims Administrator guideline: Decision based on MTUS Chronic Pain Treatment Guidelines Page(s): 24.

MAXIMUS guideline: Decision based on MTUS Chronic Pain Treatment Guidelines Page(s): 24 of 127.

Decision rationale: Regarding the request for lorazepam, Chronic Pain Medical Treatment Guidelines state the benzodiazepines are "Not recommended for long-term use because long-term efficacy is unproven and there is a risk of dependence. Most guidelines limit use to 4 weeks... Tolerance to anxiolytic effects occurs within months and long-term use may actually increase anxiety. A more appropriate treatment for anxiety disorder is an antidepressant." Within the documentation available for review, there is no documentation identifying any objective functional improvement as a result of the use of the medication and no rationale provided for long-term use of the medication despite the CA MTUS recommendation against long-term use. Benzodiazepines should not be abruptly discontinued, but unfortunately, there is no provision to modify the current request to allow tapering. In the absence of such documentation, the currently requested lorazepam is not medically necessary.

Flurbiprofen 120gm tube: Upheld

Claims Administrator guideline: Decision based on MTUS Chronic Pain Treatment Guidelines Topical Analgesics Page(s): 111-113.

MAXIMUS guideline: Decision based on MTUS Chronic Pain Treatment Guidelines Page(s): 111-113 of 127.

Decision rationale: Regarding the request for topical flurbiprofen, CA MTUS states that topical NSAIDs are indicated for "Osteoarthritis and tendinitis, in particular, that of the knee and elbow or other joints that are amenable to topical treatment: Recommended for short-term use (4-12 weeks). There is little evidence to utilize topical NSAIDs for treatment of osteoarthritis of the spine, hip or shoulder. Neuropathic pain: Not recommended as there is no evidence to support use." Within the documentation available for review, none of the abovementioned criteria have been documented. Furthermore, there is no clear rationale for the use of topical medications rather than the FDA-approved oral forms for this patient. In light of the above issues, the requested topical flurbiprofen is not medically necessary.