

Case Number:	CM14-0111509		
Date Assigned:	07/28/2014	Date of Injury:	04/30/2001
Decision Date:	09/17/2014	UR Denial Date:	06/24/2014
Priority:	Standard	Application Received:	06/24/2014

HOW THE IMR FINAL DETERMINATION WAS MADE

MAXIMUS Federal Services sent the complete case file to an expert reviewer. He/she has no affiliation with the employer, employee, providers or the claims administrator. The expert reviewer is Board Certified in Occupational Medicine and is licensed to practice in California. He/she has been in active clinical practice for more than five years and is currently working at least 24 hours a week in active practice. The expert reviewer was selected based on his/her clinical experience, education, background, and expertise in the same or similar specialties that evaluate and/or treat the medical condition and disputed items/services. He/she is familiar with governing laws and regulations, including the strength of evidence hierarchy that applies to Independent Medical Review determinations.

CLINICAL CASE SUMMARY

The expert reviewer developed the following clinical case summary based on a review of the case file, including all medical records:

Injured worker is a female with date of injury 4/30/2001. Per primary treating physician's progress report dated 5/28/2014, the injured worker complains of constant pain in her neck with 3/10 in severity. She describes her pain as throbbing. On examination blood pressure is 108/70, weight 168 pounds. There are trigger areas noted in the aracervical muscles. She has restricted range of motion on lateral bending to the right. Diagnoses include 1) cervical strain, neck 2) unspecified site of shoulder and upper arm.

IMR ISSUES, DECISIONS AND RATIONALES

The Final Determination was based on decisions for the disputed items/services set forth below:

Flexeril 10 Mg #60: Upheld

Claims Administrator guideline: Decision based on MTUS Chronic Pain Treatment Guidelines Muscle Relaxants Page(s): 63.

MAXIMUS guideline: Decision based on MTUS Chronic Pain Treatment Guidelines Cyclobenzaprine section, Muscle Relaxants Page(s): 41, 42, 63, 64.

Decision rationale: The injured worker has been using this medication for at least one year. Cyclobenzaprine is recommended by the MTUS Guidelines for short periods with acute exacerbations, but not for chronic or extended use. These guidelines report that the effect of cyclobenzaprine is greatest in the first four days of treatment. Cyclobenzaprine is associated with

a number needed to treat of three at two weeks for symptoms improvement in low back pain and is associated with drowsiness and dizziness. Chronic use of cyclobenzaprine may cause dependence, and sudden discontinuation may result in withdrawal symptoms. Discontinuation should include a tapering dose to decrease withdrawal symptoms. This request however is not for a tapering dose. The request for Flexril 10 mg #60 is determined to not be medically necessary.

Tramadol 37.5/325mg: Upheld

Claims Administrator guideline: Decision based on MTUS Chronic Pain Treatment Guidelines Tramadol Page(s): 113.

MAXIMUS guideline: Decision based on MTUS Chronic Pain Treatment Guidelines Opioids section, Weaning of Medications section Page(s): 74-95, 124.

Decision rationale: The injured worker has been using this medication for at least one year. Tramadol is a central acting synthetic opioid that exhibits opioid activity with a mechanism of action that inhibits the reuptake of serotonin and norepinephrine with side effects similar to traditional opioids. The MTUS Guidelines do not recommend the use of opioid pain medications, in general, for the management of chronic pain. There is guidance for the rare instance where opioids are needed in maintenance therapy, but the emphasis should remain on non-opioid pain medications and active therapy. Long-term use may be appropriate if the patient is showing measurable functional improvement and reduction in pain in the absence of non-compliance. Functional improvement is defined by either significant improvement in activities of daily living or a reduction in work restriction as measured during the history and physical exam. It is not recommended to discontinue opioid treatment abruptly, as weaning of medications is necessary to avoid withdrawal symptoms when opioids have been used chronically. This request however is not for a weaning treatment, but to continue treatment. The request for Tramadol 37.5/325MG is determined to not be medically necessary.