

<b>Case Number:</b>	CM14-0153889		
<b>Date Assigned:</b>	09/23/2014	<b>Date of Injury:</b>	11/24/1999
<b>Decision Date:</b>	10/24/2014	<b>UR Denial Date:</b>	08/25/2014
<b>Priority:</b>	Standard	<b>Application Received:</b>	09/22/2014

### HOW THE IMR FINAL DETERMINATION WAS MADE

MAXIMUS Federal Services sent the complete case file to an expert reviewer. He/she has no affiliation with the employer, employee, providers or the claims administrator. The expert reviewer is Board Certified in Family Medicine and is licensed to practice in California. He/she has been in active clinical practice for more than five years and is currently working at least 24 hours a week in active practice. The expert reviewer was selected based on his/her clinical experience, education, background, and expertise in the same or similar specialties that evaluate and/or treat the medical condition and disputed items/services. He/she is familiar with governing laws and regulations, including the strength of evidence hierarchy that applies to Independent Medical Review determinations.

### CLINICAL CASE SUMMARY

The expert reviewer developed the following clinical case summary based on a review of the case file, including all medical records:

This is a 60-year-old male who reported an industrial injury to the back on 11/24/1999, almost 15 years ago, attributed to the performance of his usual and customary job tasks. The patient subsequently underwent surgical intervention to the lumbar spine with a L2-S1 fusion. The patient has been treated with medications, PT, activity modification, acupuncture, massage, and ongoing care. The patient also underwent 46 days of a Functional Restoration Program. The patient continues to complain of low back pain. The objective findings on examination included tenderness and mild spasm palpable over bilateral paralumbar muscles; tenderness in the sciatic notch is; positive bilateral SLR. The patient was assessed as TDD. Patient was prescribed hydrocodone 10/Sen. 25 mg; soma; Protonix; gabapentin 600 mg; and topical compounded analgesics. Patient was also prescribed Suboxone 2-0.5 mg #30.

### IMR ISSUES, DECISIONS AND RATIONALES

The Final Determination was based on decisions for the disputed items/services set forth below:

**Hydrocodone 10/325mg #120:** Upheld

**Claims Administrator guideline:** Decision based on MTUS ACOEM Chapter 3 Initial Approaches to Treatment Page(s): 47-47, 115, Chronic Pain Treatment Guidelines Opioids Page(s): 80-82.

**MAXIMUS guideline:** Decision based on MTUS ACOEM, Chronic Pain Treatment Guidelines opioids Page(s): 74-97. Decision based on Non-MTUS Citation Official Disability Guidelines (ODG) pain chapter-opioids

**Decision rationale:** Evidence-based guidelines recommend short-term use of opioids for the management of chronic nonmalignant moderate to severe pain. Long-term use is not recommended for nonmalignant pain due to addiction, dependency, intolerance, abuse, misuse and/or side effects. Ongoing opioid management criteria are required for long-term use with evidence of reduce pain and improve function as compared to baseline measurements or a return to work. The prescription for Hydrocodone-APAP (Norco) 10/325 mg #120 for short acting pain is being prescribed as an opioid analgesic for the treatment of chronic pain to the back for the date of injury 15 years ago. The objective findings on examination do not support the medical necessity for continued opioid analgesics. The patient is being prescribed opioids for chronic mechanical low back pain, which is inconsistent with the recommendations of the CA MTUS. There is no objective evidence provided to support the continued prescription of opioid analgesics for the cited diagnoses and effects of the industrial claim. The patient should be titrated down and off the prescribed Hydrocodone. The patient is 15 years s/p DOI with reported continued issues postoperatively; however, there is no rationale supported with objective evidence to continue the use of opioids. There is no demonstrated medical necessity for the continuation of opioids for the effects of the industrial injury. The chronic use of Hydrocodone-APAP/Norco is not recommended by the CA MTUS, the ACOEM Guidelines, or the Official Disability Guidelines for the long-term treatment of chronic back/knee pain. There is no demonstrated sustained functional improvement from the prescribed opioids. The prescription of opiates on a continued long-term basis is inconsistent with the CA MTUS and the Official Disability Guidelines recommendations for the use of opiate medications for the treatment of chronic pain. There is objective evidence that supports the use of opioid analgesics in the treatment of this patient over the use of NSAIDs for the treatment of chronic pain. The current prescription of opioid analgesics is inconsistent with evidence-based guidelines. The prescription of opiates on a continued long-term basis is inconsistent with the Official Disability Guidelines recommendations for the use of opiate medications for the treatment of chronic pain. There is objective evidence that supports the use of opioid analgesics in the treatment of this patient over the use of NSAIDs for the treatment of chronic pain issues. Evidence-based guidelines necessitate documentation that the patient has signed an appropriate pain contract, functional expectations have been agreed to by the clinician, and the patient, pain medications will be provided by one physician only, and the patient agrees to use only those medications recommended or agreed to by the clinician to support the medical necessity of treatment with opioids. The ACOEM Guidelines updated chapter on chronic pain states, "Opiates for the treatment of mechanical and compressive etiologies: rarely beneficial. Chronic pain can have a mixed physiologic etiology of both neuropathic and nociceptive components. In most cases, analgesic treatment should begin with acetaminophen, aspirin, and NSAIDs (as suggested by the WHO step-wise algorithm). When these drugs do not satisfactorily reduce pain, opioids for moderate to moderately severe pain may be added to (not substituted for) the less efficacious drugs. A major concern about the use of opioids for chronic pain is that most randomized controlled trials have been limited to a short-term period (70 days). This leads to a concern about confounding issues; such as, tolerance, opioid-induced hyperalgesia, long-range adverse effects, such as, hypogonadism and/or opioid abuse, and the influence of placebo as a variable for treatment effect. ACOEM guidelines state that opioids appear to be no more effective than safer analgesics for managing most musculoskeletal symptoms; they should be used only if needed for severe pain and only for a short time. The long-term use of opioid medications may be considered in the treatment of chronic musculoskeletal pain, if: The patient has signed an appropriate pain contract; Functional expectations have been agreed to by the clinician and the patient; Pain

medications will be provided by one physician only; The patient agrees to use only those medications recommended or agreed to by the clinician. ACOEM also notes, "Pain medications are typically not useful in the subacute and chronic phases and have been shown to be the most important factor impeding recovery of function There is no clinical documentation by with objective findings on examination to support the medical necessity of Hydrocodone- APAP for this long period of time or to support ongoing functional improvement. There is no provided evidence that the patient has received benefit or demonstrated functional improvement with the prescribed Hydrocodone-APAP. There is no demonstrated medical necessity for the prescribed Opioids. There is no demonstrated medical necessity for the current prescription of tramadol with Norco. The continued prescription for Norco 10/325 mg #120 is not demonstrated to be medically necessary. The patient should be weaned down and off the prescribed hydrocodone-APAP.