

Case Number:	CM14-0163795		
Date Assigned:	10/08/2014	Date of Injury:	08/28/1996
Decision Date:	10/31/2014	UR Denial Date:	09/25/2014
Priority:	Standard	Application Received:	10/06/2014

HOW THE IMR FINAL DETERMINATION WAS MADE

MAXIMUS Federal Services sent the complete case file to an expert reviewer. He/she has no affiliation with the employer, employee, providers or the claims administrator. The expert reviewer is Board Certified in Family Medicine and is licensed to practice in North Carolina. He/she has been in active clinical practice for more than five years and is currently working at least 24 hours a week in active practice. The expert reviewer was selected based on his/her clinical experience, education, background, and expertise in the same or similar specialties that evaluate and/or treat the medical condition and disputed items/services. He/she is familiar with governing laws and regulations, including the strength of evidence hierarchy that applies to Independent Medical Review determinations.

CLINICAL CASE SUMMARY

The expert reviewer developed the following clinical case summary based on a review of the case file, including all medical records:

The patient is a 58-year-old with a reported date of injury of 08/28/1996 which occurred while moving heavy bags of concrete at work. The patient has the diagnoses of lumbago, post laminectomy syndrome of the lumbar region and lumbar degenerative disc disease. Per the most recent progress notes provided for review by the primary treating physician dated 09/11/2014, the patient had complaints of increased pain in the low back and right buttocks. The physical exam noted a positive lumbar spine Faber test, right hip sacroiliac joint tenderness and a positive Ober's and Patrick's test on the right. Treatment plan recommendations included IM injection, appeal for SI joint injection, continue TENS unit, physical therapy and pain medications.

IMR ISSUES, DECISIONS AND RATIONALES

The Final Determination was based on decisions for the disputed items/services set forth below:

Percocet 10/325mg quantity requested: 270.00: Upheld

Claims Administrator guideline: Decision based on MTUS Chronic Pain Treatment Guidelines Opioids Page(s): 92.

MAXIMUS guideline: Decision based on MTUS Chronic Pain Treatment Guidelines opioids Page(s): 76-84.

Decision rationale: The California chronic pain medical treatment guidelines section on opioids states:On-Going Management. Actions Should Include:(a) Prescriptions from a single

practitioner taken as directed, and all prescriptions from a single pharmacy.(b) The lowest possible dose should be prescribed to improve pain and function.(c) Office: Ongoing review and documentation of pain relief, functional status,appropriate medication use, and side effects. Pain assessment should include: current pain; the least reported pain over the period since last assessment; average pain; intensity of pain after taking the opioid; how long it takes for pain relief; and how long pain relief lasts. Satisfactory response to treatment may be indicated by the patient's decreased pain, increased level of function, or improved quality of life. Information from family members or other caregivers should be considered in determining the patient's response to treatment. The 4 A's for Ongoing Monitoring: Four domains have been proposed as most relevant for ongoing monitoring of chronic pain patients on opioids: pain relief, side effects, physical and psychosocial functioning, and the occurrence of any potentially aberrant (or non-adherent) drug-related behaviors. These domains have been summarized as the "4 A's" (analgesia, activities of daily living, adverse side effects, and aberrant drug taking behaviors). The monitoring of these outcomes over time should affect therapeutic decisions and provide a framework for documentation of the clinical use of these controlled drugs. (Passik, 2000)(d) Home: To aid in pain and functioning assessment, the patient should be requested to keep a pain diary that includes entries such as pain triggers, and incidence of end-of-dose pain. It should be emphasized that using this diary will help in tailoring the opioid dose. This should not be a requirement for pain management.(e) Use of drug screening or inpatient treatment with issues of abuse, addiction, or poor pain control.(f) Documentation of misuse of medications (doctor-shopping, uncontrolled drug escalation, drug diversion).(g) Continuing review of overall situation with regard to non-opioid means of pain control.(h) Consideration of a consultation with a multidisciplinary pain clinic if doses of opioids are required beyond what is usually required for the condition or pain does not improve on opioids in 3 months. Consider a psych consult if there is evidence of depression, anxiety or irritability. Consider an addiction medicine consult if there is evidence of substance misuse. When to Continue Opioids(a) If the patient has returned to work(b) If the patient has improved functioning and pain(Washington, 2002) (Colorado, 2002) (Ontario, 2000) (VA/DoD, 2003) (Maddox-AAPM/APS, 1997) (Wisconsin, 2004) (Warfield, 2004)- Chronic back pain: Appears to be efficacious but limited for short-term pain relief, and long term efficacy is unclear (>16 weeks), but also appears limited. Failure to respond to a time limited course of opioids has led to the suggestion of reassessment and consideration of alternative therapy. There is no evidence to recommend one opioid over another. In patients taking opioids for back pain, the prevalence of lifetime substance use disorders has ranged from 36% to 56% (a statistic limited by poor study design). Limited information indicated that up to one-fourth of patients who receive opioids exhibit aberrant medication-taking behavior. (Martell- Annals, 2007) (Chou, 2007) There are three studies comparing Tramadol to placebo that have reported pain relief, but this increase did not necessarily improve function. (Deshpande, 2007) The long-term use of this medication is not recommended unless certain objective outcome measures have been met as defined above. There is no provided objective outcome measure that shows significant improvement in function while on the medication or a return to work. There is no evidence of failure of other conservative treatment modalities besides physical therapy and other first line choices for chronic pain. In addition this patient's cumulative morphine equivalent daily dose is greater than 120 which is not recommended per the California MTUS. For these reasons criteria for ongoing and continued use of the medication have not been met. Therefore the request is not medically necessary.

Baclofen 10mg quantity requested: 180.00: Upheld

Claims Administrator guideline: Decision based on MTUS Chronic Pain Treatment Guidelines Muscle relaxants Page(s): 23, 64, 113.

MAXIMUS guideline: Decision based on MTUS Chronic Pain Treatment Guidelines muscle relaxants Page(s): 63-65.

Decision rationale: The California chronic pain medical treatment guidelines section on muscle relaxants states: Recommend non-sedating muscle relaxants with caution as a second-line option for short-term treatment of acute exacerbations in patients with chronic LBP. (Chou, 2007) (Mens, 2005) (Van Tulder, 1998) (van Tulder, 2003) (van Tulder, 2006) (Schnitzer, 2004) (See, 2008) Muscle relaxants may be effective in reducing pain and muscle tension, and increasing mobility. However, in most LBP cases, they show no benefit beyond NSAIDs in pain and overall

improvement. Also there is no additional benefit shown in combination with NSAIDs. Efficacy appears to diminish over time, and prolonged use of some medications in this class may lead to dependence. (Homik, 2004) Baclofen (Lioresal, generic available): The mechanism of action is blockade of the pre- and post-synaptic GABAB receptors. It is recommended orally for the treatment of spasticity and muscle spasm related to multiple sclerosis and spinal cord injuries. Baclofen has been noted to have benefits for treating lancinating, paroxysmal neuropathic pain (trigeminal neuralgia, non-FDA approved). (ICSI, 2007). The long term chronic use of this medication is not recommended per the California MTUS. The medication has not been prescribed for the acute flare up of chronic low back pain. The patient does not have multiple sclerosis or spinal cord injury. The specific use of this medication for greater than 3 weeks is not recommended per the California MTUS. The criteria set forth above for its use has not been met. Therefore the request is not medically necessary.

Oxycodone HCL 10mg quantity requested: 270.00: Upheld

Claims Administrator guideline: Decision based on MTUS Chronic Pain Treatment Guidelines Opioids Page(s): 92.

MAXIMUS guideline: Decision based on MTUS Chronic Pain Treatment Guidelines opioids Page(s): 76-84.

Decision rationale: The California chronic pain medical treatment guidelines section on opioids states: On-Going Management. Actions Should Include: (a) Prescriptions from a single practitioner taken as directed, and all prescriptions from a single pharmacy. (b) The lowest possible dose should be prescribed to improve pain and function. (c) Office: Ongoing review and documentation of pain relief, functional status, appropriate medication use, and side effects. Pain assessment should include: current pain; the least reported pain over the period since last assessment; average pain; intensity of pain after taking the opioid; how long it takes for pain relief; and how long pain relief lasts. Satisfactory response to treatment may be indicated by the patient's decreased pain, increased level of function, or improved quality of life. Information from family members or other caregivers should be considered in determining the patient's response to treatment. The 4 A's for Ongoing Monitoring: Four domains have been proposed as most relevant for ongoing monitoring of chronic pain patients on opioids: pain relief, side effects, physical and psychosocial functioning, and the occurrence of any potentially aberrant (or non-adherent) drug-related behaviors. These domains have been summarized as the "4 A's" (analgesia, activities of daily living, adverse side effects, and aberrant drug taking behaviors). The monitoring of these outcomes over time should affect therapeutic decisions and provide a framework for documentation of the clinical use of these controlled drugs. (Passik, 2000) (d) Home: To aid in pain and functioning assessment, the patient should be requested to keep a pain diary that includes entries such as pain triggers, and incidence of end-of-dose pain. It should be emphasized that using this diary will help in tailoring the opioid dose. This should not be a requirement for pain management. (e) Use of drug screening or inpatient treatment with issues of abuse, addiction, or poor pain control. (f) Documentation of misuse of medications (doctor-shopping, uncontrolled drug escalation, drug diversion). (g) Continuing review of overall situation with regard to non-opioid means of pain control. (h) Consideration of a consultation with a multidisciplinary pain clinic if doses of opioids are required beyond what is usually required for

the condition or pain does not improve on opioids in 3 months. Consider a psych consult if there is evidence of depression, anxiety or irritability. Consider an addiction medicine consult if there is evidence of substance misuse. When to Continue Opioids(a) If the patient has returned to work(b) If the patient has improved functioning and pain(Washington, 2002) (Colorado, 2002) (Ontario, 2000) (VA/DoD, 2003) (Maddox-AAPM/APS, 1997) (Wisconsin, 2004) (Warfield, 2004)- Chronic back pain: Appears to be efficacious but limited for short-term pain relief, and long term efficacy is unclear (>16 weeks), but also appears limited. Failure to respond to a time limited course of opioids has led to the suggestion of reassessment and consideration of alternative therapy. There is no evidence to recommend one opioid over another. In patients taking opioids for back pain, the prevalence of lifetime substance use disorders has ranged from 36% to 56% (a statistic limited by poor study design). Limited information indicated that up to one-fourth of patients who receive opioids exhibit aberrant medication-taking behavior. (Martell- Annals, 2007) (Chou, 2007) There are three studies comparing Tramadol to placebo that have reported pain relief, but this increase did not necessarily improve function. (Deshpande, 2007) The long-term use of this medication is not recommended unless certain objective outcome measures have been met as defined above. There is no provided objective outcome measure that shows significant improvement in function while on the medication or a return to work. There is no evidence of failure of other conservative treatment modalities besides physical therapy and other first line choices for chronic pain. In addition this patient's cumulative morphine equivalent daily dose is greater than 120 which is not recommended per the California MTUS. For these reasons criteria for ongoing and continued use of the medication have not been met. Therefore the request is not medically necessary.