

Case Number:	CM14-0169910		
Date Assigned:	10/20/2014	Date of Injury:	06/25/2007
Decision Date:	11/20/2014	UR Denial Date:	09/22/2014
Priority:	Standard	Application Received:	10/14/2014

HOW THE IMR FINAL DETERMINATION WAS MADE

MAXIMUS Federal Services sent the complete case file to an expert reviewer. He/she has no affiliation with the employer, employee, providers or the claims administrator. The expert reviewer is Board Certified in anesthesia, has a subspecialty in pain medicine and is licensed to practice in Massachusetts. He/she has been in active clinical practice for more than five years and is currently working at least 24 hours a week in active practice. The expert reviewer was selected based on his/her clinical experience, education, background, and expertise in the same or similar specialties that evaluate and/or treat the medical condition and disputed items/services. He/she is familiar with governing laws and regulations, including the strength of evidence hierarchy that applies to Independent Medical Review determinations.

CLINICAL CASE SUMMARY

The expert reviewer developed the following clinical case summary based on a review of the case file, including all medical records:

According to the documents available for review, the patient is an injured female worker. The date of injury is June 25, 2007. The patient sustained an injury to the right knee. The specific mechanism of injury was not fully elaborated on in the notes available for review. The patient currently complains of pain in the knee worsened with ambulation and rated 9/10 in severity. The patient is maintained on the multimodal pain medication regimen including Motrin, Norco, Voltaren gel and Flexeril. A request for Motrin, Norco, Voltaren Gel and Flexeril was denied.

IMR ISSUES, DECISIONS AND RATIONALES

The Final Determination was based on decisions for the disputed items/services set forth below:

Motrin 800 mg TID PRN #90 with 3 refills: Upheld

Claims Administrator guideline: Decision based on MTUS Chronic Pain Treatment Guidelines.

MAXIMUS guideline: Decision based on MTUS Chronic Pain Treatment Guidelines NSAIDs Page(s): 67, 70-73.

Decision rationale: According to the MTUS Anti-inflammatories is the traditional first line of treatment, to reduce pain so activity and functional restoration can resume, but long-term use may not be warranted. (Van Tulder-Cochrane, 2000) Recommended with cautions below. Disease-State Warnings for all NSAIDs: All NSAIDs have [U.S. Boxed Warning]: for

associated risk of adverse cardiovascular events, including, MI, stroke, and new onset or worsening of pre-existing hypertension. NSAIDs should never be used right before or after a heart surgery (CABG - coronary artery bypass graft). NSAIDs can cause ulcers and bleeding in the stomach and intestines at any time during treatment (FDA Medication Guide). See NSAIDs, GI Symptoms and Cardiovascular Risks. Other disease-related concerns (non-boxed warnings): Hepatic: Use with caution in patients with moderate hepatic impairment and not recommended for patients with severe hepatic impairment. Borderline elevations of one or more liver enzymes may occur in up to 15% of patients taking NSAIDs. Renal: Use of NSAIDs may compromise renal function. FDA Medication Guide is provided by FDA mandate on all prescriptions dispensed for NSAIDs. Routine Suggested Monitoring: Package inserts for NSAIDs recommend periodic lab monitoring of a CBC and chemistry profile (including liver and renal function tests). There has been a recommendation to measure liver transaminases within 4 to 8 weeks after starting therapy, but the interval of repeating lab tests after this treatment duration has not been established. Routine blood pressure monitoring is recommended. Overall Dosing Recommendation: It is generally recommended that the lowest effective dose be used for all NSAIDs for the shortest duration of time consistent with the individual patient treatment goals. According to the documents available for review, it appears that the patient is taking this medication for long-term therapy of a chronic condition. Given the increased risks associated with long-term use of this medication and no documented evidence that the lowest possible dose is being used for the shortest period of time, the requirements for treatment have not been met and medical necessity has not been established.

Norco 10/325 mg Q6H PRN #120 with 2 refills: Upheld

Claims Administrator guideline: Decision based on MTUS Chronic Pain Treatment Guidelines.

MAXIMUS guideline: Decision based on MTUS Chronic Pain Treatment Guidelines Opioids, On-Going Management Page(s): 74-97.

Decision rationale: There is no current documentation of baseline pain, pain score with use of opioids, functional improvement on current regimen, side effects or review of potentially aberrant drug taking behaviors as outlined in the MTUS and as required for ongoing treatment. Therefore, at this time, the requirements for treatment have not been met and medical necessity has not been established.

Voltaren cream 100 mg apply three times daily PRN with 2 refills: Upheld

Claims Administrator guideline: Decision based on MTUS Chronic Pain Treatment Guidelines.

MAXIMUS guideline: Decision based on MTUS Chronic Pain Treatment Guidelines Voltaren Page(s): 112.

Decision rationale: According to the MTUS, Voltaren Gel 1% (diclofenac): Indicated for relief of osteoarthritis pain in joints that lend themselves to topical treatment (ankle, elbow, foot, hand,

knee, and wrist). It has not been evaluated for treatment of the spine, hip or shoulder. Maximum dose should not exceed 32 g per day (8 g per joint per day in the upper extremity and 16 g per joint per day in the lower extremity). The most common adverse reactions were dermatitis and pruritus. (Voltaren package insert) For additional adverse effects: See NSAIDs, GI symptoms and cardiovascular risk; and NSAIDs, hypertension and renal function. Additionally, accordingly to the ODG, Voltaren gel is not recommended as a first-line treatment. Voltaren Gel is recommended for osteoarthritis after failure of an oral NSAID, or contraindications to oral NSAIDs, or for patients who cannot swallow solid oral dosage forms, and after considering the increased risk profile with diclofenac, including topical formulations. According to the documents available for review, there is no indication that the patient has had a failure of oral NSAIDs, a contraindication to oral NSAIDs or cannot swallow solid oral dosage forms. Therefore, at this time, the requirements for treatment have not been met and medical necessity has not been established.

Flexeril 10 mg BID PRN #90 with 3 refills: Upheld

Claims Administrator guideline: Decision based on MTUS Chronic Pain Treatment Guidelines.

MAXIMUS guideline: Decision based on MTUS Chronic Pain Treatment Guidelines Flexeril Page(s): 41-42.

Decision rationale: Accordingly to the MTUS, current treatment guidelines recommend this medication is an option for chronic pain using a short course of therapy. The effect of Flexeril is great is the first four days of treatment, suggesting a shorter course as many better. This medication is not recommended as an addition to other medications. Longer course of Flexeril also are not recommended to be for longer than 2 to 3 weeks as prolonged use me lead to dependence. According to the records, the patient has been taking his medication chronically. Therefore, at this time, the requirements for treatment have not been met and medical necessity has not been established.