

Case Number:	CM14-0184199		
Date Assigned:	11/12/2014	Date of Injury:	03/06/2003
Decision Date:	12/16/2014	UR Denial Date:	10/18/2014
Priority:	Standard	Application Received:	11/05/2014

HOW THE IMR FINAL DETERMINATION WAS MADE

MAXIMUS Federal Services sent the complete case file to an expert reviewer. He/she has no affiliation with the employer, employee, providers or the claims administrator. The expert reviewer is Board Certified in General Preventive Medicine and is licensed to practice in Indiana. He/she has been in active clinical practice for more than five years and is currently working at least 24 hours a week in active practice. The expert reviewer was selected based on his/her clinical experience, education, background, and expertise in the same or similar specialties that evaluate and/or treat the medical condition and disputed items/services. He/she is familiar with governing laws and regulations, including the strength of evidence hierarchy that applies to Independent Medical Review determinations.

CLINICAL CASE SUMMARY

The expert reviewer developed the following clinical case summary based on a review of the case file, including all medical records:

This employee is a 59 year old male with date of injury of 3/6/2003. A review of the medical records indicate that the patient is undergoing treatment for cervical and lumbar disc degeneration. Subjective complaints include continued pain in her neck and shoulders with radiation down both upper extremities; continued pain in the lower back with radiation down both lower extremities. Objective findings include limited range of motion of the cervical and lumbar spine with tenderness to palpation of the paravertebrals; positive straight leg raise bilaterally. Treatment has included physical therapy, epidural steroid injections, and Norco. The utilization review dated 10/18/2014 partially-certified Norco and Omeprazole.

IMR ISSUES, DECISIONS AND RATIONALES

The Final Determination was based on decisions for the disputed items/services set forth below:

Hydrocodone/APAP 5/325 MG #120: Upheld

Claims Administrator guideline: Decision based on MTUS Chronic Pain Treatment Guidelines Opioids.

MAXIMUS guideline: Decision based on MTUS Chronic Pain Treatment Guidelines Opioids Page(s): 74-96. Decision based on Non-MTUS Citation Official Disability Guidelines (ODG) Neck and Upper Back (Acute and Chronic), Low Back - Lumbar & Thoracic (Acute & Chronic), Shoulder, Pain, Opioids

Decision rationale: ODG does not recommend the use of opioids for neck, low back, and shoulder pain "except for short use for severe cases, not to exceed 2 weeks." The patient has exceeded the 2 week recommended treatment length for opioid usage. MTUS does not discourage use of opioids past 2 weeks, but does state that "ongoing review and documentation of pain relief, functional status, appropriate medication use, and side effects. Pain assessment should include: current pain; the least reported pain over the period since last assessment; average pain; intensity of pain after taking the opioid; how long it takes for pain relief; and how long pain relief lasts. Satisfactory response to treatment may be indicated by the patient's decreased pain, increased level of function, or improved quality of life." The treating physician does not fully document the least reported pain over the period since last assessment, intensity of pain after taking opioid, pain relief, increased level of function, or improved quality of life. Additionally, medical documents indicate that the patient has been on Norco in excess of the recommended 2-week limit. As such, the question for Hydrocodone/APAP 5/325 MG #120 is not medically necessary.

Omeprazole 20 MG: Upheld

Claims Administrator guideline: Decision based on MTUS Chronic Pain Treatment Guidelines Non-Steroidal Anti-Inflammatory Drugs (NSAIDs).

MAXIMUS guideline: Decision based on MTUS Chronic Pain Treatment Guidelines NSAIDs, GI symptoms & cardiovascular risk Page(s): 68-69. Decision based on Non-MTUS Citation Official Disability Guidelines (ODG) Pain (Chronic), NSAIDs, GI symptoms & cardiovascular risk

Decision rationale: MTUS and ODG states, "Determine if the patient is at risk for gastrointestinal events: (1) age > 65 years; (2) history of peptic ulcer, GI bleeding or perforation; (3) concurrent use of ASA, corticosteroids, and/or an anticoagulant; or (4) high dose/multiple NSAID (e.g., NSAID + low-dose ASA)." And "Patients at intermediate risk for gastrointestinal events and no cardiovascular disease:(1) A non-selective NSAID with either a PPI (Proton Pump Inhibitor, for example, 20 mg omeprazole daily) or misoprostol (200 g four times daily) or(2) a Cox-2 selective agent. Long-term PPI use (> 1 year) has been shown to increase the risk of hip fracture (adjusted odds ratio 1.44)." The medical documents provided do not establish the patient as having documented GI bleeding/perforation/peptic ulcer or other GI risk factors as outlined in MTUS. As such, the request for Omeprazole 20mg is not medically necessary.