

Case Number:	CM14-0021241		
Date Assigned:	06/11/2014	Date of Injury:	06/11/1997
Decision Date:	12/12/2014	UR Denial Date:	02/13/2014
Priority:	Standard	Application Received:	02/20/2014

HOW THE IMR FINAL DETERMINATION WAS MADE

MAXIMUS Federal Services sent the complete case file to an expert reviewer. He/she has no affiliation with the employer, employee, providers or the claims administrator. The expert reviewer is Board Certified in Family Medicine and is licensed to practice in California. He/she has been in active clinical practice for more than five years and is currently working at least 24 hours a week in active practice. The expert reviewer was selected based on his/her clinical experience, education, background, and expertise in the same or similar specialties that evaluate and/or treat the medical condition and disputed items/services. He/she is familiar with governing laws and regulations, including the strength of evidence hierarchy that applies to Independent Medical Review determinations.

CLINICAL CASE SUMMARY

The expert reviewer developed the following clinical case summary based on a review of the case file, including all medical records:

The patient is a 57 year-old woman who was injured at work on 6/11/1997. The injury was primarily to her neck and back. She is requesting review of denial for the following: Norco 10/325mg, Q12 hours, #90 and Soma 350mg, #90. Medical records corroborate ongoing care for her injuries. Her orthopedic surgeon's notes describe the following chronic diagnoses: Status Post Anterior Cervical Discectomy and Fusion, C3-4, with removal of hardware at C4-5 and C5-6 on 8/19/2010; Lumbar Sprain/Strain with Radiculopathy; Rotator Cuff Tear, Status Post Right Shoulder Arthroscopy on 5/7/2008; and Status Post Right Knee Arthroscopy with Degenerative Osteoarthritis, Chondromalacia, and Meniscal Injury. She has been on Norco and Soma at least since 12/31/2013.

IMR ISSUES, DECISIONS AND RATIONALES

The Final Determination was based on decisions for the disputed items/services set forth below:

Norco 10/325mg, Q12hrs PRN #90: Upheld

Claims Administrator guideline: Decision based on MTUS Chronic Pain Treatment Guidelines Opioids and Chronic Pain.

MAXIMUS guideline: Decision based on MTUS Chronic Pain Treatment Guidelines Opioids Page(s): 76-78, 80.

Decision rationale: The MTUS/Chronic Pain Medical Treatment Guidelines comment on the long-term use of opioids. These guidelines have established criteria on the use of opioids for the ongoing management of pain. Actions should include: prescriptions from a single practitioner and from a single pharmacy. The lowest possible dose should be prescribed to improve pain and function. There should be an ongoing review and documentation of pain relief, functional status, appropriate medication use and side effects. Pain assessment should include: current pain, the least reported pain over the period since last assessment; average pain; intensity of pain after taking the opioid; how long it takes for pain relief; and how long pain relief lasts. Satisfactory response to treatment may be indicated by the patient's decreased pain, increased level of function, or improved quality of life. There should be evidence of documentation of the "4 A's for Ongoing Monitoring." These four domains include: pain relief, side effects, physical and psychological functioning, and the occurrence of any potentially aberrant drug-related behaviors. Further, there should be consideration of a consultation with a multidisciplinary pain clinic if doses of opioids are required beyond what is usually required for the condition or pain that does not improve on opioids in 3 months. There should be consideration of an addiction medicine consult if there is evidence of substance misuse (Pages 76-78). Finally, the guidelines indicate that for chronic back pain, the long-term efficacy of opioids is unclear. Failure to respond to a time-limited course of opioids has led to the suggestion of reassessment and consideration of alternative therapy (Page 80). Based on the review of the medical records, there is insufficient documentation in support of these stated MTUS/Chronic Pain Medical Treatment Guidelines for the ongoing use of opioids. There is insufficient documentation of the "4 A's for Ongoing Monitoring." The treatment course of opioids in this patient has extended well beyond the timeframe required for a reassessment of therapy. In summary, there is insufficient documentation to support the chronic use of an opioid in this patient. Treatment with Norco is not considered as medically necessary.

Soma 350mg BID #90: Upheld

Claims Administrator guideline: Decision based on MTUS Chronic Pain Treatment Guidelines.

MAXIMUS guideline: Decision based on MTUS Chronic Pain Treatment Guidelines Carisoprodol (Soma) Page(s): 29.

Decision rationale: The MTUS/Chronic Pain Medical Treatment Guidelines comment on the use of carisoprodol (Soma) for the treatment of pain. Soma is not recommended for this purpose. This medication is not indicated for long-term use. Carisoprodol is a commonly prescribed, centrally acting skeletal muscle relaxant whose primary active metabolite is meprobamate (a schedule-IV controlled substance). Carisoprodol is now scheduled in several states but not on a federal level. It has been suggested that the main effect is due to generalized sedation and treatment of anxiety. Abuse has been noted for sedative and relaxant effects. In regular abusers the main concern is the accumulation of meprobamate. Carisoprodol abuse has also been noted in order to augment or alter effects of other drugs. This includes the following: (1) increasing sedation of benzodiazepines or alcohol; (2) use to prevent side effects of cocaine; (3) use with tramadol to produce relaxation and euphoria; (4) as a combination with hydrocodone, an effect that some abusers claim is similar to heroin (referred to as a "Las Vegas Cocktail"); & (5) as a

combination with codeine (referred to as ""Soma Coma""). (Reeves, 1999) (Reeves, 2001) (Reeves, 2008) (Schears, 2004) There was a 300% increase in numbers of emergency room episodes related to carisoprodol from 1994 to 2005. (DHSS, 2005) Intoxication appears to include subdued consciousness, decreased cognitive function, and abnormalities of the eyes, vestibular function, appearance, gait and motor function. Intoxication includes the effects of both carisoprodol and meprobamate, both of which act on different neurotransmitters. (Bramness, 2007) (Bramness, 2004) A withdrawal syndrome has been documented that consists of insomnia, vomiting, tremors, muscle twitching, anxiety, and ataxia when abrupt discontinuation of large doses occurs. This is similar to withdrawal from meprobamate. (Reeves, 2007) (Reeves, 2004) There is little research in terms of weaning of high dose carisoprodol and there is no standard treatment regimen for patients with known dependence. Tapering should be individualized for each patient. (Boothby, 2003) In summary, Soma is not recommended per the above stated guidelines. The use of Soma in this patient is not considered as a medically necessary treatment.