

Case Number:	CM14-0043031		
Date Assigned:	07/25/2014	Date of Injury:	04/09/2003
Decision Date:	09/25/2014	UR Denial Date:	03/28/2014
Priority:	Standard	Application Received:	04/10/2014

HOW THE IMR FINAL DETERMINATION WAS MADE

MAXIMUS Federal Services sent the complete case file to an expert reviewer. He/she has no affiliation with the employer, employee, providers or the claims administrator. The expert reviewer is Board Certified in Family Medicine, and is licensed to practice in California. He/she has been in active clinical practice for more than five years and is currently working at least 24 hours a week in active practice. The expert reviewer was selected based on his/her clinical experience, education, background, and expertise in the same or similar specialties that evaluate and/or treat the medical condition and disputed items/services. He/she is familiar with governing laws and regulations, including the strength of evidence hierarchy that applies to Independent Medical Review determinations.

CLINICAL CASE SUMMARY

The expert reviewer developed the following clinical case summary based on a review of the case file, including all medical records:

This is a 46-year-old female who reported an industrial injury to the back on 4/9/2003, over 11 years ago, attributed to the performance of her customary job tasks reported as lifting a five-gallon can. The patient is receiving opioid therapy for chronic low back pain with a radiculopathy. The patient complained of low back pain radiating to the upper buttock and possibly to the posterior thigh. The patient reported intermittent tingling to the right lower extremity. The objective findings on examination included no evidence of scarring; no spasms; no atrophy; tenderness to the bilateral lumbar paravertebral regions at L4 and L5; tenderness to palpation of the right SI joint and medial proximal buttock; lumbar spine range of motion was diminished; SLR was negative; neurologic exam revealed no focal motor deficits or muscle atrophy of the lower extremities; intact sensation; reflexes 2+; weight 240 pounds. The patient was noted to of had two prior SCS implantations which were subsequently removed subsequent the failure. The patient was prescribed Ambien CR; Theramine; ducal locks; tizanidine 4 mg; Lyrica 75 mg; OxyContin 80 mg; Norco 10/325 mg; Fioricet 50/325/40 mg and venlafaxine 75 mg. The patient was noted to be receiving the morphine equivalent dose of 480 mg per day. The patient was prescribed OxyContin 80 mg #84.

IMR ISSUES, DECISIONS AND RATIONALES

The Final Determination was based on decisions for the disputed items/services set forth below:

Oxycontin 80 mg # 84: Upheld

Claims Administrator guideline: Decision based on MTUS Chronic Pain Treatment Guidelines Chronic Pain Medical Treatment Guidelines Page(s): 13, 23, 78-92 and 105.

MAXIMUS guideline: Decision based on MTUS ACOEM Chapter 12 Low Back Complaints Page(s): 300-306, Chronic Pain Treatment Guidelines opioids Page(s): 74-97. Decision based on Non-MTUS Citation Official Disability Guidelines (ODG) pain chapter opioids American College of Occupational and Environmental Medicine (ACOEM), 2nd Edition, (2004) chapter 6 pages 114-116.

Decision rationale: The prescription for OxyContin 80 mg #84 for short acting pain is being prescribed as an opioid analgesic for the treatment of chronic pain to the back for the date of injury 11 years ago. The objective findings on examination do not support the medical necessity for continued opioid analgesics for chronic mechanical back pain. The patient is noted to be taking 480 MEDs per day without a demonstrated functional improvement. The patient is being prescribed opioids for mechanical back pain, which is inconsistent with the recommendations of the CA MTUS. There is no objective evidence provided to support the continued prescription of opioid analgesics for the cited diagnoses and effects of the industrial claim. The patient should be titrated down and off the prescribed OxyContin 80 mg #84. The patient is 11 years s/p DOI with reported continued issues. There is no demonstrated medical necessity for the continuation of opioids for the effects of the industrial injury. The chronic use of OxyContin 80 mg #84 is not recommended by the CA MTUS, the ACOEM Guidelines, or the Official Disability Guidelines for the long-term treatment of chronic back pain. The prescription of opiates on a continued long-term basis is inconsistent with the CA MTUS and the Official Disability Guidelines recommendations for the use of opiate medications for the treatment of chronic pain. There is objective evidence that supports the use of opioid analgesics in the treatment of this patient over the use of NSAIDs for the treatment of chronic pain. The current prescription of opioid analgesics is inconsistent with evidence-based guidelines. The prescription of opiates on a continued long-term basis is inconsistent with the Official Disability Guidelines recommendations for the use of opiate medications for the treatment of chronic pain. There is objective evidence that supports the use of opioid analgesics in the treatment of this patient over the use of NSAIDs for the treatment of chronic pain issues. Evidence-based guidelines necessitate documentation that the patient has signed an appropriate pain contract, functional expectations have been agreed to by the clinician, and the patient, pain medications will be provided by one physician only, and the patient agrees to use only those medications recommended or agreed to by the clinician to support the medical necessity of treatment with opioids. The ACOEM Guidelines updated chapter on chronic pain states, "Opiates for the treatment of mechanical and compressive etiologies: rarely beneficial. Chronic pain can have a mixed physiologic etiology of both neuropathic and nociceptive components. In most cases, analgesic treatment should begin with acetaminophen, aspirin, and NSAIDs (as suggested by the WHO step-wise algorithm). When these drugs do not satisfactorily reduce pain, opioids for moderate to moderately severe pain may be added to (not substituted for) the less efficacious drugs. A major concern about the use of opioids for chronic pain is that most randomized controlled trials have been limited to a short-term period (70 days). This leads to a concern about confounding issues; such as, tolerance, opioid-induced hyperalgesia, long-range adverse effects such as hypogonadism and/or opioid abuse, and the influence of placebo as a variable for treatment effect." ACOEM guidelines state that opioids appear to be no more effective than safer analgesics for managing most

musculoskeletal symptoms; they should be used only if needed for severe pain and only for a short time. The long-term use of opioid medications may be considered in the treatment of chronic musculoskeletal pain, if: The patient has signed an appropriate pain contract; Functional expectations have been agreed to by the clinician and the patient; Pain medications will be provided by one physician only; The patient agrees to use only those medications recommended or agreed to by the clinician. ACOEM also notes, "Pain medications are typically not useful in the subacute and chronic phases and have been shown to be the most important factor impeding recovery of function." There is no clinical documentation by with objective findings on examination to support the medical necessity of OxyContin 80 mg for this long period of time or to support ongoing functional improvement. There is no provided evidence that the patient has received benefit or demonstrated functional improvement with the prescribed OxyContin 80 mg. There is no demonstrated medical necessity for the prescribed Opioids as there is no demonstrated functional improvement for the prescribed high dose opioids. The continued prescription for OxyContin 80 mg #84 is not demonstrated to be medically necessary.