

Case Number:	CM14-0064796		
Date Assigned:	07/11/2014	Date of Injury:	07/31/2002
Decision Date:	12/19/2014	UR Denial Date:	04/28/2014
Priority:	Standard	Application Received:	05/08/2014

HOW THE IMR FINAL DETERMINATION WAS MADE

MAXIMUS Federal Services sent the complete case file to an expert reviewer. He/she has no affiliation with the employer, employee, providers or the claims administrator. The expert reviewer is Board Certified in Anesthesiology, has a subspecialty in Acupuncture & Pain Medicine, and is licensed to practice in California. He/she has been in active clinical practice for more than five years and is currently working at least 24 hours a week in active practice. The expert reviewer was selected based on his/her clinical experience, education, background, and expertise in the same or similar specialties that evaluate and/or treat the medical condition and disputed items/services. He/she is familiar with governing laws and regulations, including the strength of evidence hierarchy that applies to Independent Medical Review determinations.

CLINICAL CASE SUMMARY

The expert reviewer developed the following clinical case summary based on a review of the case file, including all medical records:

65y/o female injured worker with date of injury 7/31/02 with related left shoulder pain. Per progress report dated 4/10/14, the injured worker complained of left shoulder pain secondary to osteoarthritis of the left shoulder. She reported insomnia due to chronic pain. Physical exam findings were not documented. The documentation submitted for review did not state whether physical therapy was utilized. Treatment to date has included medication management. The date of UR decision was 4/18/14.

IMR ISSUES, DECISIONS AND RATIONALES

The Final Determination was based on decisions for the disputed items/services set forth below:

Soma 350mg #60: Upheld

Claims Administrator guideline: Decision based on MTUS Chronic Pain Treatment Guidelines. Decision based on Non-MTUS Citation Official disability guidelines

MAXIMUS guideline: Decision based on MTUS Chronic Pain Treatment Guidelines Carisoprodol Page(s): 29.

Decision rationale: Per MTUS CPMTG p29, "Not recommended. This medication is not indicated for long-term use. Carisoprodol is a commonly prescribed, centrally acting skeletal muscle relaxant whose primary active metabolite is meprobamate (a schedule-IV controlled substance). Carisoprodol is now scheduled in several states but not on a federal level. It has been

suggested that the main effect is due to generalized sedation and treatment of anxiety. Abuse has been noted for sedative and relaxant effects. In regular abusers the main concern is the accumulation of meprobamate. Carisoprodol abuse has also been noted in order to augment or alter effects of other drugs."As this medication is not recommended by MTUS, it is not medically necessary.

Aciphex 20mg #60: Upheld

Claims Administrator guideline: Decision based on MTUS Chronic Pain Treatment Guidelines.

MAXIMUS guideline: Decision based on MTUS Chronic Pain Treatment Guidelines NSAIDs,GI Symptoms & Cardiovascular Risk Page(s): 68. Decision based on Non-MTUS Citation Official Disability Guidelines (ODG) Pain, Proton Pump Inhibitors

Decision rationale: In the treatment of dyspepsia secondary to NSAID therapy, the MTUS recommends stopping the NSAID, switching to a different NSAID, or considering the use of an H2-receptor antagonist or a PPI. The MTUS Chronic Pain Medical Treatment Guidelines recommend the use of proton pump inhibitors in conjunction with NSAIDs in situations in which the patient is at risk for gastrointestinal events including: (1) age > 65 years; (2) history of peptic ulcer, GI bleeding or perforation; (3) concurrent use of ASA, corticosteroids, and/or an anticoagulant; or (4) high dose/multiple NSAID (e.g., NSAID + low-dose ASA). CPMTG guidelines further specify: "Recommendations: Patients with no risk factor and no cardiovascular disease: Non-selective NSAIDs OK (e.g, ibuprofen, naproxen, etc.). Patients at intermediate risk for gastrointestinal events and no cardiovascular disease : (1) a non-selective NSAID with either a PPI (Proton Pump Inhibitor, for example, 20 mg omeprazole daily) or misoprostol (200 four times daily) or (2) a Cox-2 selective agent. Long-term PPI use (> 1 year) has been shown to increase the risk of hip fracture (adjusted odds ratio 1.44). Patients at high risk for gastrointestinal events with no cardiovascular disease: A Cox-2 selective agent plus a PPI if absolutely necessary. Patients at high risk of gastrointestinal events with cardiovascular disease: If GI risk is high the suggestion is for a low-dose Cox-2 plus low dose Aspirin (for cardioprotection) and a PPI. If cardiovascular risk is greater than GI risk the suggestion is naproxyn plus low-dose aspirin plus a PPI. (Laine, 2006) (Scholmerich, 2006) (Nielsen, 2006) (Chan, 2004) (Gold, 2007) (Laine, 2007)" Per ODG TWC, "many prescribers believe that this class of drugs is innocuous, but much information is available to demonstrate otherwise. A trial of omeprazole or lansoprazole is recommended before Nexium therapy. The other PPIs, Protonix, Dexilant, and Aciphex, should also be second-line."As there is no documentation of peptic ulcer, GI bleeding or perforation, or cardiovascular disease in the records available for my review, the injured worker's risk for gastrointestinal events is low, as such, medical necessity cannot be affirmed. Furthermore, as noted per the guidelines, Aciphex is a second-line medication. The medical records do not establish whether the patient has failed attempts at first line PPIs, such as omeprazole or lansoprazole, which should be considered prior to prescribing a second line PPI such as Protonix. The request is not medically necessary.

Ibuprofen 400mg #100: Overturned

Claims Administrator guideline: Decision based on MTUS Chronic Pain Treatment Guidelines NSAIDs.

MAXIMUS guideline: Decision based on MTUS Chronic Pain Treatment Guidelines NSAIDs
Page(s): 67-68.

Decision rationale: Current guidelines note that evidence is limited to make an initial recommendation with acetaminophen, and that NSAIDs may be more efficacious for treatment. The selection of acetaminophen as a first-line treatment appears to be made primarily based on side effect profile in osteoarthritis guidelines. The most recent Cochrane review on this subject suggests that non-steroidal anti-inflammatory drugs (NSAIDs) are more efficacious for osteoarthritis in terms of pain reduction, global assessments and improvement of functional status. The request is indicated for the injured worker's shoulder osteoarthritis. I respectfully disagree with the UR physician's denial based on proximity to a previous approval. The documentation submitted for review contains no evidence of the aforementioned approval. The request is medically necessary.

Methadone 10mg #630: Upheld

Claims Administrator guideline: Decision based on MTUS Chronic Pain Treatment Guidelines Opioids.

MAXIMUS guideline: Decision based on MTUS Chronic Pain Treatment Guidelines Opioids
Page(s): 61, 78.

Decision rationale: With regard to methadone, the MTUS CPMTG states: "Recommended as a second-line drug for moderate to severe pain if the potential benefit outweighs the risk. The FDA reports that they have received reports of severe morbidity and mortality with this medication. This appears, in part, secondary to the long half-life of the drug (8-59 hours). Pain relief on the other hand only lasts from 4-8 hours. Methadone should only be prescribed by providers experienced in using it." Per MTUS Chronic Pain Medical Treatment Guidelines p78 regarding on-going management of opioids "Four domains have been proposed as most relevant for ongoing monitoring of chronic pain patients on opioids: Pain relief, side effects, physical and psychosocial functioning, and the occurrence of any potentially aberrant (or nonadherent) drug related behaviors. These domains have been summarized as the '4As' (Analgesia, activities of daily living, adverse side effects, and any aberrant drug-taking behaviors). The monitoring of these outcomes over time should affect therapeutic decisions and provide a framework for documentation of the clinical use of these controlled drugs." Review of the available medical records reveals neither documentation to support the medical necessity of methadone nor any documentation addressing the '4 A's' domains, which is a recommended practice for the on-going management of opioids. Specifically, the notes do not appropriately review and document pain relief, functional status improvement, appropriate medication use, or side effects. The MTUS considers this list of criteria for initiation and continuation of opioids in the context of efficacy required to substantiate medical necessity, and they do not appear to have been addressed by the treating physician in the documentation available for review. Furthermore, efforts to rule out aberrant behavior (e.g. CURES report, UDS, opiate agreement) are necessary to assure safe

usage and establish medical necessity. There is no documentation comprehensively addressing this concern in the records available for my review. As MTUS recommends discontinuing opioids if there is no overall improvement in function, medical necessity cannot be affirmed. The request is not medically necessary

Norco 10/325 #60: Upheld

Claims Administrator guideline: Decision based on MTUS Chronic Pain Treatment Guidelines Opioids.

MAXIMUS guideline: Decision based on MTUS Chronic Pain Treatment Guidelines Opioids Page(s): 78, 91.

Decision rationale: Per MTUS Chronic Pain Medical Treatment Guidelines p78 regarding on-going management of opioids "Four domains have been proposed as most relevant for ongoing monitoring of chronic pain patients on opioids: Pain relief, side effects, physical and psychosocial functioning, and the occurrence of any potentially aberrant (or nonadherent) drug related behaviors. These domains have been summarized as the '4As' (Analgesia, activities of daily living, adverse side effects, and any aberrant drug-taking behaviors).The monitoring of these outcomes over time should affect therapeutic decisions and provide a framework for documentation of the clinical use of these controlled drugs." Review of the available medical records reveals neither documentation to support the medical necessity of Norco nor any documentation addressing the '4 A's' domains, which is a recommended practice for the on-going management of opioids. Specifically, the notes do not appropriately review and document pain relief, functional status improvement, appropriate medication use, or side effects. The MTUS considers this list of criteria for initiation and continuation of opioids in the context of efficacy required to substantiate medical necessity, and they do not appear to have been addressed by the treating physician in the documentation available for review. Furthermore, efforts to rule out aberrant behavior (e.g. CURES report, UDS, opiate agreement) are necessary to assure safe usage and establish medical necessity. There is no documentation comprehensively addressing this concern in the records available for my review. As MTUS recommends discontinuing opioids if there is no overall improvement in function, medical necessity cannot be affirmed. The request is not medically necessary.