

<b>Case Number:</b>	CM14-0085399		
<b>Date Assigned:</b>	07/23/2014	<b>Date of Injury:</b>	10/02/2009
<b>Decision Date:</b>	10/14/2014	<b>UR Denial Date:</b>	06/02/2014
<b>Priority:</b>	Standard	<b>Application Received:</b>	06/09/2014

### HOW THE IMR FINAL DETERMINATION WAS MADE

MAXIMUS Federal Services sent the complete case file to an expert reviewer. He/she has no affiliation with the employer, employee, providers or the claims administrator. The expert reviewer is Board Certified in Anesthesiology, has a subspecialty in Pain Management and is licensed to practice in Georgia. He/she has been in active clinical practice for more than five years and is currently working at least 24 hours a week in active practice. The expert reviewer was selected based on his/her clinical experience, education, background, and expertise in the same or similar specialties that evaluate and/or treat the medical condition and disputed items/services. He/she is familiar with governing laws and regulations, including the strength of evidence hierarchy that applies to Independent Medical Review determinations.

### CLINICAL CASE SUMMARY

The expert reviewer developed the following clinical case summary based on a review of the case file, including all medical records:

The claimant is a 47 year old male presenting with chronic pain following a work related injury on 10/02/2009. The claimant was diagnosed with ankle enthesopathy, limb pain, knee pain, brachial plexus disorder and rotator cuff syndrome. On 02/12/2014, the claimant complained of pain in the right shoulder and the right knee, and left foot. MRI of the left foot was significant for neuroma as well as tarsal tunnel syndrome by EMG. The physical exam showed increased pain in the left foot, and large hematoma in the right thumb.

### IMR ISSUES, DECISIONS AND RATIONALES

The Final Determination was based on decisions for the disputed items/services set forth below:

**Norco 10mg-325mg # 50:** Upheld

**Claims Administrator guideline:** Decision based on MTUS Chronic Pain Treatment Guidelines Chronic Pain Medical Treatment Guidelines; Opioids Ongoing Management.

**MAXIMUS guideline:** Decision based on MTUS Chronic Pain Treatment Guidelines Opioids Page(s): 79.

**Decision rationale:** Norco 10mg-325mg # 50 is not medically necessary. Per MTUS Page 79 of MTUS guidelines states that weaning of opioids are recommended if (a) there are no overall improvement in function, unless there are extenuating circumstances (b) continuing pain with evidence of intolerable adverse effects (c) decrease in functioning (d) resolution of pain (e) if

serious non-adherence is occurring (f) the patient requests discontinuing. The claimant's medical records did not document that there was an overall improvement in function or a return to work with previous opioid therapy. The claimant has long-term use with this medication and there was a lack of improved function with this opioid; therefore the requested medication is not medically necessary.

**Soma 350mg #120:** Upheld

**Claims Administrator guideline:** Decision based on MTUS Chronic Pain Treatment Guidelines Chronic Pain Medical Treatment Guidelines; Soma/Carisoprodol.

**MAXIMUS guideline:** Decision based on MTUS Chronic Pain Treatment Guidelines Carisoprodol.

**Decision rationale:** Soma 350mg #120 is not medically necessary. Ca MTUS states that Soma is not recommended. This medication is not indicated for long-term use. Carisoprodol is commonly prescribed, centrally acting skeletal muscle relaxant and its primary active metabolite is meprobamate (schedule for controlled substances). Carisoprodol is now scheduled in several states but not on the federal level. Since been suggested that the main affect is due to generalized sedation and treatment of anxiety. Abuse has been noted for sedatives and relaxants effects. In regular basis to maintain concern is the cannulation of medical date. Carisoprodol abuse has also been noted in order to augment or alter effects of other drugs. This includes the following: Increasing sedation of benzodiazepines or alcohol; used to prevent side effects of cocaine; use with tramadol to produce relaxation and euphoria; as a combination with hydrocodone, and affected some abusers claim is similar to heroin; the combination with codeine. There was a 300% increase in numbers of emergency room episodes related to Terrace Woodall from 1994 2005. Intoxication appears to include subjective consciousness, decreased cognitive function, and abnormalities of the eyes, vestibular function, appearance, gait and motor function. Intoxication includes the effects of both cars up at all and meprobamate, both of which act on different neurotransmitters. A withdrawal syndrome has been documented that consists of insomnia, vomiting, tremors, muscle twitching, anxiety, and ataxia when abrupt discontinuation of large doses occur. This is similar to withdrawal from meprobamate. There is little research in terms of weaning of high dose carries up at all and there is no standard treatment regimen for patients with known dependence. Most treatment includes treatment for symptomatic complaints of a stroke. Another option is to switch to phenobarbital to prevent withdrawal with subsequent tapering. A maximum dose of phenobarbital is 500 mg per day and the taper is 3 mg per day with a slower taper in an outpatient setting. Tapering should be individualized to reach patient. There was no specific time limit for the prescription of this medication or a weaning protocol; therefore Soma is not medically necessary.

**Zolpidem 10mg #30:** Upheld

**Claims Administrator guideline:** The Claims Administrator did not base their decision on the MTUS. Decision based on Non-MTUS Citation Official Disability Guidelines (ODG) Pain; Zolpidem

**MAXIMUS guideline:** The Expert Reviewer did not base their decision on the MTUS. Decision based on Non-MTUS Citation Official Disability Guidelines (ODG) <Insert Section (for example Knee)>, <Insert Topic (for example Total Knee Arthroplasty)>

**Decision rationale:** Zolpidem 10 mg at bedtime # 30 with 2 refills (prescribed 7/3/14) is not medically necessary. The ODG states that Ambien "is not recommended for long term use, but recommended for short-term use. While sleeping pills, so called minor tranquilizers, and anti-anxiety agents are commonly prescribed in chronic pain, pain specialist rarely, if ever, recommend them for long-term use. They can be habit-forming and they may impair function and memory more than opioid pain relievers. There is also concern that they may increase pain and depression over long-term. Ambien is indicated for treatment of insomnia with difficulty of sleep onset and/or sleep maintenance. Longer-term studies have found Ambien ER to be effective for up to 24 weeks in adults. According to the medical records it is unclear how long the claimant was on the sleeping aid medication of this class. Additionally, there is no documentation of sleep disorder requiring this medication. It is more appropriate to set a weaning protocol at this point. Ambien 10mg is not medically necessary.