

Case Number:	CM14-0086417		
Date Assigned:	07/23/2014	Date of Injury:	05/16/2001
Decision Date:	09/19/2014	UR Denial Date:	05/23/2014
Priority:	Standard	Application Received:	06/09/2014

HOW THE IMR FINAL DETERMINATION WAS MADE

MAXIMUS Federal Services sent the complete case file to an expert reviewer. He/she has no affiliation with the employer, employee, providers or the claims administrator. The expert reviewer is Board Certified in Physical Medicine and Rehabilitation, has a subspecialty in Pain Management and is licensed to practice in California. He/she has been in active clinical practice for more than five years and is currently working at least 24 hours a week in active practice. The expert reviewer was selected based on his/her clinical experience, education, background, and expertise in the same or similar specialties that evaluate and/or treat the medical condition and disputed items/services. He/she is familiar with governing laws and regulations, including the strength of evidence hierarchy that applies to Independent Medical Review determinations.

CLINICAL CASE SUMMARY

The expert reviewer developed the following clinical case summary based on a review of the case file, including all medical records:

This is a patient with a date of injury of 5/15/01. A utilization review determination dated 5/23/14 recommends non-certification of stool softener and nabumetone. OxyContin and Percocet were modified from #60 each to #30 each. 5/6/14 medical report identifies patient is stable on the medication with no adverse effects. He has been asked to cut Percocet to two a day and is willing to try this for a month. On exam, "gait/station-walking slow, no antalgic gait." Medication was refilled.

IMR ISSUES, DECISIONS AND RATIONALES

The Final Determination was based on decisions for the disputed items/services set forth below:

Percocet 10/325 #60: Upheld

Claims Administrator guideline: Decision based on MTUS Chronic Pain Treatment Guidelines Percocet Page(s): 78.

MAXIMUS guideline: Decision based on MTUS Chronic Pain Treatment Guidelines 8 C.C.R. 9792.20 - 9792.26 MTUS (Effective July 18, 2009) Page(s): 76-79, 120 of 127.

Decision rationale: Regarding the request for Percocet, California Pain Medical Treatment Guidelines state that, due to high abuse potential, close follow-up is recommended with documentation of analgesic effect, objective functional improvement, side effects, and discussion regarding any aberrant use. Guidelines go on to recommend discontinuing opioids if there is no

documentation of improved function and pain. Within the documentation available for review, there is no indication that the medication is improving the patient's function or pain (in terms of specific examples of functional improvement and percent reduction in pain or reduced NRS) and no discussion regarding aberrant use. As such, there is no clear indication for ongoing use of the medication. Opioids should not be abruptly discontinued, but unfortunately, there is no provision to modify the current request to allow tapering. In light of the above issues, the currently requested Percocet is not medically necessary.

Oxycontin 60mg #60: Upheld

Claims Administrator guideline: Decision based on MTUS Chronic Pain Treatment Guidelines Opioids Page(s): 78.

MAXIMUS guideline: Decision based on MTUS Chronic Pain Treatment Guidelines 8 C.C.R. 9792.20 - 9792.26 MTUS (Effective July 18, 2009) Page(s): 76-79, 120 of 127.

Decision rationale: Regarding the request for OxyContin, California Pain Medical Treatment Guidelines state that, due to high abuse potential, close follow-up is recommended with documentation of analgesic effect, objective functional improvement, side effects, and discussion regarding any aberrant use. Guidelines go on to recommend discontinuing opioids if there is no documentation of improved function and pain. Within the documentation available for review, there is no indication that the medication is improving the patient's function or pain (in terms of specific examples of functional improvement and percent reduction in pain or reduced NRS) and no discussion regarding aberrant use. As such, there is no clear indication for ongoing use of the medication. Opioids should not be abruptly discontinued, but unfortunately, there is no provision to modify the current request to allow tapering. In light of the above issues, the currently requested OxyContin is not medically necessary.

Stool softener 100mg #840: Upheld

Claims Administrator guideline: The Claims Administrator did not cite any medical evidence for its decision.

MAXIMUS guideline: Decision based on MTUS Chronic Pain Treatment Guidelines 8 C.C.R. 9792.20 - 9792.26 MTUS (Effective July 18, 2009) Page(s): 76-79, 120 of 127.

Decision rationale: Regarding the request for stool softener, California MTUS does support the prophylactic treatment of constipation for patients undergoing opioid therapy. However, as the opioids are not medically necessary, the stool softener is not medically necessary.

Nabumetome 750mg #360: Upheld

Claims Administrator guideline: Decision based on MTUS Chronic Pain Treatment Guidelines NSAIDS Page(s): 70,72.

MAXIMUS guideline: Decision based on MTUS Chronic Pain Treatment Guidelines 8 C.C.R. 9792.20 - 9792.26 MTUS (Effective July 18, 2009) Page(s): 67-72 of 127.

Decision rationale: Regarding the request for nabumetone, Chronic Pain Medical Treatment Guidelines state that NSAIDs are recommended at the lowest dose for the shortest period in patients with moderate to severe pain. Within the documentation available for review, there is no indication that Naproxen is providing any specific analgesic benefits (in terms of percent pain reduction, or reduction in numeric rating scale) or any objective functional improvement. In the absence of such documentation, the currently requested nabumetone is not medically necessary.