

Case Number:	CM14-0101218		
Date Assigned:	07/30/2014	Date of Injury:	12/01/2010
Decision Date:	01/06/2015	UR Denial Date:	06/25/2014
Priority:	Standard	Application Received:	07/01/2014

HOW THE IMR FINAL DETERMINATION WAS MADE

MAXIMUS Federal Services sent the complete case file to an expert reviewer. He/she has no affiliation with the employer, employee, providers or the claims administrator. The expert reviewer is Board Certified in Neurology, has a subspecialty in Neuromuscular Medicine and is licensed to practice in New Jersey. He/she has been in active clinical practice for more than five years and is currently working at least 24 hours a week in active practice. The expert reviewer was selected based on his/her clinical experience, education, background, and expertise in the same or similar specialties that evaluate and/or treat the medical condition and disputed items/services. He/she is familiar with governing laws and regulations, including the strength of evidence hierarchy that applies to Independent Medical Review determinations.

CLINICAL CASE SUMMARY

The expert reviewer developed the following clinical case summary based on a review of the case file, including all medical records:

The patient is a 47 year old woman who sustained a work-related injury on December 1 2010. Subsequently, the patient developed a chronic sacroiliac and lumbar pain. According to a progress report dated on May 13 2014, the patient was complaining of right lower back pain. The patient physical examination demonstrated lumbar tenderness with reduced range of motion and positive provocative tests. The patient was diagnosed with lumbar arthropathy and chronic sacroiliac pain. The provider requested authorization for following medications.

IMR ISSUES, DECISIONS AND RATIONALES

The Final Determination was based on decisions for the disputed items/services set forth below:

Tramadol 50mg #60 with 2 refills: Upheld

Claims Administrator guideline: Decision based on MTUS Chronic Pain Treatment Guidelines Opioids, Criteria for Use.

MAXIMUS guideline: Decision based on MTUS Chronic Pain Treatment Guidelines Tramadol Page(s): 113.

Decision rationale: According to MTUS guidelines, Ultram (Tramadol) is a synthetic opioid indicated for the pain management but not recommended as a first line oral analgesic. In addition and according to MTUS guidelines, ongoing use of opioids should follow specific rules:(a) Prescriptions from a single practitioner taken as directed, and all prescriptions from a single

pharmacy.(b) The lowest possible dose should be prescribed to improve pain and function.(c) Office: Ongoing review and documentation of pain relief, functional status, appropriate medication use, and side effects. Four domains have been proposed as most relevant for ongoing monitoring of chronic pain patients on opioids: pain relief, side effects, physical and psychosocial functioning, and the occurrence of any potentially aberrant (or non-adherent) drug-related behaviors. These domains have been summarized as the "4 A's" (analgesia, activities of daily living, adverse side effects, and aberrant drug taking behaviors). The monitoring of these outcomes over time should affect therapeutic decisions and provide a framework. Although, Tramadol may be needed to help with the patient pain, there is no clear evidence of objective and recent functional and pain improvement from its previous use. There is no clear documentation of the efficacy/safety of previous use of Tramadol. There is no recent evidence of objective monitoring of compliance of the patient with his medications. Therefore, the prescription of Tramadol is not medically necessary.

Tizanidine 4mg #30 with 2 refills: Upheld

Claims Administrator guideline: Decision based on MTUS Chronic Pain Treatment Guidelines Muscle Relaxants (For Pain).

MAXIMUS guideline: Decision based on MTUS Chronic Pain Treatment Guidelines Muscle Relaxants Page(s): 63.

Decision rationale: According to MTUS guidelines, non-sedating muscle relaxants are recommended with caution as a second line option for short term treatment of acute exacerbations in patients with chronic lumbosacral pain. Efficacy appears to diminish over time and prolonged use may cause dependence. The patient was previously treated with Tizanidine, which is considered a prolonged use of the drug. There is no continuous and objective documentation of the effect of the drug on patient pain, spasm and function. There is no recent documentation for recent pain exacerbation or failure of first line treatment of medication. Furthermore, there is no clear exacerbation of back pain and spasm and the prolonged use of Zanaflex is not justified. Therefore, the request for Tizanidine 4mg is not medically necessary.