

Case Number:	CM14-0107242		
Date Assigned:	08/01/2014	Date of Injury:	05/10/2014
Decision Date:	01/12/2015	UR Denial Date:	06/27/2014
Priority:	Standard	Application Received:	07/10/2014

HOW THE IMR FINAL DETERMINATION WAS MADE

MAXIMUS Federal Services sent the complete case file to an expert reviewer. He/she has no affiliation with the employer, employee, providers or the claims administrator. The expert reviewer is Board Certified in Anesthesia, has a subspecialty in Acupuncture & Pain Medicine and is licensed to practice in California. He/she has been in active clinical practice for more than five years and is currently working at least 24 hours a week in active practice. The expert reviewer was selected based on his/her clinical experience, education, background, and expertise in the same or similar specialties that evaluate and/or treat the medical condition and disputed items/services. He/she is familiar with governing laws and regulations, including the strength of evidence hierarchy that applies to Independent Medical Review determinations.

CLINICAL CASE SUMMARY

The expert reviewer developed the following clinical case summary based on a review of the case file, including all medical records:

This 54 year old woman sustained an industrial injury on 5/10/2014 after a slip and fall which resulted in cervical, lumbar, and left shoulder sprain/strain injuries. Treatment has included x-rays of the cervical and lumbar spine indicating degenerative disc disease, and a left shoulder x-ray that was negative for acute displaced fracture performed on 5/14/2014. The worker has received physical therapy with instruction on a home exercise regimen and oral medications. Physician notes from 6/12/2014 show the worker was ordered for modified work duty and to continue physical therapy. On 6/26/2014 Utilization Review evaluated the prescription for MRIs to the cervical and lumbar spines and the left shoulder. The physician noted that there were no neurological deficits, 5+/5 strength to all four extremities, and sensation intact with complaints of moderate pain. The requests were denied and subsequently appealed to Independent Medical Review.

IMR ISSUES, DECISIONS AND RATIONALES

The Final Determination was based on decisions for the disputed items/services set forth below:

MRI of the cervical spine: Upheld

Claims Administrator guideline: Decision based on MTUS ACOEM Chapter 8 Neck and Upper Back Complaints.

MAXIMUS guideline: Decision based on MTUS ACOEM Chapter 8 Neck and Upper Back Complaints Page(s): 177.

Decision rationale: ACOEM guidelines support ordering of imaging studies for emergence of red flags, physiologic evidence of tissue insult or neurologic dysfunction, failure to progress in a strengthening program intended to avoid surgery, and clarification of the anatomy prior to an invasive procedure. Physiologic evidence may be in the form of definitive neurologic findings on physical examination, electrodiagnostic studies, laboratory tests, or bone scans. Unequivocal findings that identify specific nerve compromise on the neurologic examination are sufficient evidence to warrant imaging studies if symptoms persist. When the neurologic examination is less clear, however, further physiologic evidence of nerve dysfunction can be obtained before ordering an imaging study. Electromyography (EMG), and nerve conduction velocities (NCV), including H-reflex tests, may help identify subtle focal neurologic dysfunction in patients with neck or arm symptoms, or both, lasting more than three or four weeks. The documentation submitted for review contained no evidence of neurologic dysfunction to warrant MRI study. The request is not medically necessary.

MRI of lumbar spine: Upheld

Claims Administrator guideline: Decision based on MTUS ACOEM Chapter 12 Low Back Complaints.

MAXIMUS guideline: Decision based on MTUS ACOEM Chapter 8 Neck and Upper Back Complaints Page(s): 177.

Decision rationale: ACOEM guidelines support ordering of imaging studies for emergence of red flags, physiologic evidence of tissue insult or neurologic dysfunction, failure to progress in a strengthening program intended to avoid surgery, and clarification of the anatomy prior to an invasive procedure. Physiologic evidence may be in the form of definitive neurologic findings on physical examination, electrodiagnostic studies, laboratory tests, or bone scans. Unequivocal findings that identify specific nerve compromise on the neurologic examination are sufficient evidence to warrant imaging studies if symptoms persist. When the neurologic examination is less clear, however, further physiologic evidence of nerve dysfunction can be obtained before ordering an imaging study. Electromyography (EMG), and nerve conduction velocities (NCV), including H-reflex tests, may help identify subtle focal neurologic dysfunction in patients with neck or arm symptoms, or both, lasting more than three or four weeks. The documentation submitted for review contained no evidence of neurologic dysfunction to warrant MRI study. The request is not medically necessary.

MRI of the left shoulder: Upheld

Claims Administrator guideline: Decision based on MTUS ACOEM Chapter 9 Shoulder Complaints.

MAXIMUS guideline: Decision based on MTUS ACOEM Chapter 9 Shoulder Complaints Page(s): 177.

Decision rationale: ACOEM guidelines support ordering of imaging studies for emergence of red flags, physiologic evidence of tissue insult or neurologic dysfunction, failure to progress in a strengthening program intended to avoid surgery, and clarification of the anatomy prior to an invasive procedure. Physiologic evidence may be in the form of definitive neurologic findings on physical examination, electrodiagnostic studies, laboratory tests, or bone scans. Unequivocal findings that identify specific nerve compromise on the neurologic examination are sufficient evidence to warrant imaging studies if symptoms persist. When the neurologic examination is less clear, however, further physiologic evidence of nerve dysfunction can be obtained before ordering an imaging study. Electromyography (EMG), and nerve conduction velocities (NCV), including H-reflex tests, may help identify subtle focal neurologic dysfunction in patients with neck or arm symptoms, or both, lasting more than three or four weeks. The documentation submitted for review contained no evidence of neurologic dysfunction to warrant MRI study. The request is not medically necessary.