

<b>Case Number:</b>	CM14-0123464		
<b>Date Assigned:</b>	09/16/2014	<b>Date of Injury:</b>	11/08/2002
<b>Decision Date:</b>	07/22/2015	<b>UR Denial Date:</b>	07/14/2014
<b>Priority:</b>	Standard	<b>Application Received:</b>	08/05/2014

### HOW THE IMR FINAL DETERMINATION WAS MADE

MAXIMUS Federal Services sent the complete case file to an expert reviewer. He/she has no affiliation with the employer, employee, providers or the claims administrator. He/she has been in active clinical practice for more than five years and is currently working at least 24 hours a week in active practice. The expert reviewer was selected based on his/her clinical experience, education, background, and expertise in the same or similar specialties that evaluate and/or treat the medical condition and disputed items/Service. He/she is familiar with governing laws and regulations, including the strength of evidence hierarchy that applies to Independent Medical Review determinations.

The Expert Reviewer has the following credentials:  
 State(s) of Licensure: New York  
 Certification(s)/Specialty: Anesthesiology

### CLINICAL CASE SUMMARY

The expert reviewer developed the following clinical case summary based on a review of the case file, including all medical records:

The injured worker is a 56 year old female who sustained an industrial injury on 11/8/02. The injured worker was diagnosed as having lumbar region unspecified disc disorder, shoulder impingement syndrome and lumbar radiculopathy. Currently, the injured worker was with complaints of sacroiliac joint pain. Previous treatments included medication management. The plan of care was for medication prescriptions.

### IMR ISSUES, DECISIONS AND RATIONALES

The Final Determination was based on decisions for the disputed items/services set forth below:

**Norco 10/325mg #240:** Upheld

**Claims Administrator guideline:** Decision based on MTUS Chronic Pain Treatment Guidelines Opioids.

**MAXIMUS guideline:** Decision based on MTUS Chronic Pain Treatment Guidelines Opioids Page(s): 91-97. Decision based on Non-MTUS Citation Official Disability Guidelines (ODG) Opioids.

**Decision rationale:** According to the CA MTUS and ODG, Norco 10/325mg (Hydrocodone/Acetaminophen) is a short-acting opioid analgesic indicated for moderate to moderately severe pain, and is used to manage both acute and chronic pain. The treatment of chronic pain with any opioid analgesic requires review and documentation of pain relief, functional status, appropriate medication use, and side effects. A pain assessment should include current pain, intensity of pain after taking the opiate, and the duration of pain relief. In this case, there is no documentation of the medication's functional benefit. Medical necessity of the requested item has not been established. Of note, discontinuation of an opioid analgesic should include a taper, to avoid withdrawal symptoms. The requested medication is not medically necessary.

**Ultram 150mg #90:** Upheld

**Claims Administrator guideline:** Decision based on MTUS Chronic Pain Treatment Guidelines Opioids.

**MAXIMUS guideline:** Decision based on MTUS Chronic Pain Treatment Guidelines Opioids Page(s): 91-97. Decision based on Non-MTUS Citation Official Disability Guidelines (ODG) Opioids.

**Decision rationale:** According to the California MTUS, Tramadol (Ultram) is a synthetic opioid which affects the central nervous system and is indicated for the treatment of moderate to severe pain. Per CA MTUS Guidelines, certain criteria need to be followed, including an ongoing review and documentation of pain relief and functional status, appropriate medication use, and side effects. Pain assessment should include current pain: last reported pain over the period since last assessment; average pain; intensity of pain after taking the opioid, and the duration of pain relief. According to the medical records, there has been no documentation of the medication's analgesic effectiveness and no clear documentation that the patient has responded to ongoing opioid therapy. Medical necessity of the requested medication has not been established. Of note, discontinuation of an opioid analgesic requires a taper to avoid withdrawal symptoms. The requested medication is not medically necessary.

**Topical Compound: Cyclobenzaprine 60gm tube:** Upheld

**Claims Administrator guideline:** Decision based on MTUS Chronic Pain Treatment Guidelines Topical Analgesics, Muscle Relaxants.

**MAXIMUS guideline:** Decision based on MTUS Chronic Pain Treatment Guidelines Topical Analgesics Page(s): 111-113. Decision based on Non-MTUS Citation Official Disability Guidelines (ODG) Topical analgesics.

**Decision rationale:** According to the California MTUS Guidelines (2009), topical analgesics are primarily recommended for neuropathic pain when trials of antidepressants and anticonvulsants have failed. These agents are applied topically to painful areas with advantages that include lack of systemic side effects, absence of drug interactions, and no need to titrate. Many agents are compounded as monotherapy or in combination for pain control including, for example, NSAIDs, opioids, capsaicin, muscle relaxants, local anesthetics or antidepressants. In this case, the requested topical agent is the muscle relaxant, Cyclobenzaprine. Cyclobenzaprine is not FDA approved for use as a topical application. Medical necessity for the requested topical analgesic has not been established. The requested topical gel is not medically necessary.

**Percocet 10/325 #180:** Upheld

**Claims Administrator guideline:** Decision based on MTUS Chronic Pain Treatment Guidelines Oxycodone/acetaminophen (Percocet).

**MAXIMUS guideline:** Decision based on MTUS Chronic Pain Treatment Guidelines Opioids Page(s): 91-97. Decision based on Non-MTUS Citation Official Disability Guidelines (ODG) Opioids.

**Decision rationale:** According to the CA MTUS and the ODG, Percocet (Oxycodone/Acetaminophen) is a short-acting opioid analgesic indicated for moderate to severe pain, and is used to manage both acute and chronic pain. The treatment of chronic pain with any opioid analgesic requires review and documentation of pain relief, functional status, appropriate medication use, and side effects. A pain assessment should include current pain, intensity of pain after taking the opiate, and the duration of pain relief. In this case, there is no documentation of the medication's pain relief effectiveness, functional status, or response to ongoing opioid analgesic therapy. Also, there is no indication as to why this patient requires Norco and Ultram in addition to Percocet. Medical necessity of the requested medication has not been established. Of note, discontinuation of an opioid analgesic should include a taper, to avoid withdrawal symptoms. The requested treatment with Percocet 10/325 mg is not medically necessary.