

Case Number:	CM14-0131802		
Date Assigned:	08/20/2014	Date of Injury:	12/04/2003
Decision Date:	01/29/2015	UR Denial Date:	08/09/2014
Priority:	Standard	Application Received:	08/18/2014

HOW THE IMR FINAL DETERMINATION WAS MADE

MAXIMUS Federal Services sent the complete case file to an expert reviewer. He/she has no affiliation with the employer, employee, providers or the claims administrator. The expert reviewer is Board Certified in Emergency Medicine and is licensed to practice in New York. He/she has been in active clinical practice for more than five years and is currently working at least 24 hours a week in active practice. The expert reviewer was selected based on his/her clinical experience, education, background, and expertise in the same or similar specialties that evaluate and/or treat the medical condition and disputed items/services. He/she is familiar with governing laws and regulations, including the strength of evidence hierarchy that applies to Independent Medical Review determinations.

CLINICAL CASE SUMMARY

The expert reviewer developed the following clinical case summary based on a review of the case file, including all medical records:

The patient is a 45-year-old female who was injured on December 4, 2003. The patient continued to experience low back pain. Physical examination was notable for tightness in the the trapezius muscles bilaterally, tenderness to low back, intact motor strength, and positive straight leg raise. Diagnoses included lumbar radiculopathy. Treatment included medications and epidural steroid injections. Requests for authorization for Norco 10/325 mg #120 and quazepam 15 mg #30 were submitted for consideration.

IMR ISSUES, DECISIONS AND RATIONALES

The Final Determination was based on decisions for the disputed items/services set forth below:

Norco 10/325mg #120: Upheld

Claims Administrator guideline: Decision based on MTUS Chronic Pain Treatment Guidelines Opioids.

MAXIMUS guideline: Decision based on MTUS Chronic Pain Treatment Guidelines Pain Interventions and Guidelines Page(s): 11, 74-96.

Decision rationale: Norco is the compounded medication containing hydrocodone and acetaminophen. Chronic Pain Medical Treatment Guidelines state that opioids are not recommended as a first line therapy. Opioid should be part of a treatment plan specific for the patient and should follow criteria for use. Criteria for use include establishment of a treatment

plan, determination if pain is nociceptive or neuropathic, failure of pain relief with non-opioid analgesics, setting of specific functional goals, and opioid contract with agreement for random drug testing. If analgesia is not obtained, opioids should be discontinued. The patient should be screened for likelihood that he or she could be weaned from the opioids if there is no improvement in pain of function. It is recommended for short term use if first-line options, such as acetaminophen or NSAIDS have failed. Opioids may be a safer choice for patients with cardiac and renal disease than antidepressants or anticonvulsants. Acetaminophen is recommended for treatment of chronic pain & acute exacerbations of chronic pain. Acetaminophen overdose is a well-known cause of acute liver failure. Hepatotoxicity from therapeutic doses is unusual. Renal insufficiency occurs in 1 to 2% of patients with overdose. The recommended dose for mild to moderate pain is 650 to 1000 mg orally every 4 hours with a maximum of 4 g/day. In this case, the patient has been using Norco since at least November 2013 and has not obtained analgesia. In addition, there is no documentation that the patient has signed an opioid contract. Criteria for long-term opioid use have not been met. The request is not medically necessary.

Quazepam 15mg #30: Upheld

Claims Administrator guideline: Decision based on MTUS Chronic Pain Treatment Guidelines Benzodiazepines.

MAXIMUS guideline: The Expert Reviewer did not base their decision on the MTUS. Decision based on Non-MTUS Citation Official Disability Guidelines (ODG) Pain <Insomnia treatment.

Decision rationale: Insomnia treatment should be based on etiology. Pharmacological agents should only be used after careful evaluation of potential causes of sleep disturbance. Failure of sleep disturbance to resolve in a 7 to 10 day period may indicate a psychiatric and/or medical illness. Quazepam is a benzodiazepine FDA approved for sleep onset insomnia. These medications are only recommended for short-term use due to risk of tolerance, dependence, and adverse events (daytime drowsiness, anterograde amnesia, next-day sedation, impaired cognition, impaired psychomotor function, and rebound insomnia). These drugs have been associated with sleep-related activities such as sleep driving, cooking and eating food, and making phone calls (all while asleep). Particular concern is noted for patients at risk for abuse or addiction. Withdrawal occurs with abrupt discontinuation or large decreases in dose. In this case, the patient had been taking insomnia medication since at least February 2014. The duration of pharmacologic treatment for insomnia surpasses the recommended short-term duration of use. The request is not medically necessary.