

<b>Case Number:</b>	CM14-0137990		
<b>Date Assigned:</b>	09/05/2014	<b>Date of Injury:</b>	08/11/2004
<b>Decision Date:</b>	07/28/2015	<b>UR Denial Date:</b>	08/14/2014
<b>Priority:</b>	Standard	<b>Application Received:</b>	08/26/2014

### HOW THE IMR FINAL DETERMINATION WAS MADE

MAXIMUS Federal Services sent the complete case file to an expert reviewer. He/she has no affiliation with the employer, employee, providers or the claims administrator. He/she has been in active clinical practice for more than five years and is currently working at least 24 hours a week in active practice. The expert reviewer was selected based on his/her clinical experience, education, background, and expertise in the same or similar specialties that evaluate and/or treat the medical condition and disputed items/Service. He/she is familiar with governing laws and regulations, including the strength of evidence hierarchy that applies to Independent Medical Review determinations.

The Expert Reviewer has the following credentials:

State(s) of Licensure: North Carolina

Certification(s)/Specialty: Family Practice

### CLINICAL CASE SUMMARY

The expert reviewer developed the following clinical case summary based on a review of the case file, including all medical records:

The injured worker is a 54 year-old female who sustained an industrial injury on 08/11/04. She reported neck and low back pain after heavy lifting. Current diagnoses include chronic pain syndrome, postlaminectomy syndrome cervical and lumbar, radiculitis cervical and lumbar, cervicgia, spondylosis-cervical, displacement intervertebral disc-cervical, degeneration intervertebral disc-lumbar, lumbago, carpal tunnel syndrome, pain in soft tissues-limb, and lesion of ulnar nerve. In a progress note dated 08/07/14, the treating physician reports she continues to complain of sharp, burning, tingling, low back pain to lower extremity with continued neck, shoulder, and upper extremity pain. The pain is exacerbated by walking, standing, or sitting; it is alleviated by medications. She is losing sensation in her hands and feet with increased upper extremity weakness. She is status post 2 falls; she uses a cane. Examination reveals cervical, thoracic, lumbar paraspinal, and bilateral sciatic notch tenderness. Cervical and Lumbar spine range of motion are decreased; she has give away weakness to upper and lower extremities bilaterally. There is chin-forward thoracic kyphosis and loss of lumbar lordosis. She has positive Spurling's left and right, causing radiating pain in the shoulders. Treatment recommendation includes continuation of medications, physical therapy, epidural and trigger point injections, pain psychology evaluation, and repeat lumbar spine MRI to assess for new herniation. There is no indication she has been able to return to work. Date of Utilization Review: 08/14/14

### IMR ISSUES, DECISIONS AND RATIONALES

The Final Determination was based on decisions for the disputed items/services set forth below:

**Repeat MRI of the lumbar spine:** Upheld

**Claims Administrator guideline:** Decision based on MTUS ACOEM Chapter 12 Low Back Complaints Page(s): 287.

**MAXIMUS guideline:** Decision based on MTUS ACOEM Chapter 12 Low Back Complaints Page(s): 303.

**Decision rationale:** The ACOEM chapter on low back complaints and special diagnostic studies states: Unequivocal objective findings that identify specific nerve compromise on the neurologic examination are sufficient evidence to warrant imaging in patients who do not respond to treatment and who would consider surgery an option. When the neurologic examination is less clear, however, further physiologic evidence of nerve dysfunction should be obtained before ordering an imaging study. Indiscriminant imaging will result in false-positive findings, such as disk bulges, that are not the source of painful symptoms and do not warrant surgery. If physiologic evidence indicates tissue insult or nerve impairment, the practitioner can discuss with a consultant the selection of an imaging test to define a potential cause (magnetic resonance imaging [MRI] for neural or other soft tissue, computed tomography [CT] for bony structures). Relying solely on imaging studies to evaluate the source of low back and related symptoms carries a significant risk of diagnostic confusion (false positive test results) because of the possibility of identifying a finding that was present before symptoms began and therefore has no temporal association with the symptoms. Techniques vary in their abilities to define abnormalities (Table 12-7). Imaging studies should be reserved for cases in which surgery is considered or red-flag diagnoses are being evaluated. Because the overall false-positive rate is 30% for imaging studies in patients over age 30 who do not have symptoms, the risk of diagnostic confusion is great. There is no recorded presence of emerging red flags on the physical exam. There is evidence of nerve compromise on physical exam but there is not mention of consideration for surgery or complete failure of conservative therapy. For these reasons, criteria for imaging as defined above per the ACOEM have not been met. Therefore the request is not medically necessary.