

Case Number:	CM14-0186371		
Date Assigned:	11/14/2014	Date of Injury:	12/18/2008
Decision Date:	02/17/2015	UR Denial Date:	11/05/2014
Priority:	Standard	Application Received:	11/10/2014

HOW THE IMR FINAL DETERMINATION WAS MADE

MAXIMUS Federal Services sent the complete case file to an expert reviewer. He/she has no affiliation with the employer, employee, providers or the claims administrator. The expert reviewer is Board Certified in Preventive Medicine, has a subspecialty in Occupational Medicine and is licensed to practice in Iowa. He/she has been in active clinical practice for more than five years and is currently working at least 24 hours a week in active practice. The expert reviewer was selected based on his/her clinical experience, education, background, and expertise in the same or similar specialties that evaluate and/or treat the medical condition and disputed items/services. He/she is familiar with governing laws and regulations, including the strength of evidence hierarchy that applies to Independent Medical Review determinations.

CLINICAL CASE SUMMARY

The expert reviewer developed the following clinical case summary based on a review of the case file, including all medical records:

This is a 39 year old patient with date of injury of 12/18/2008. Medical records indicate the patient is undergoing treatment for shoulder joint pain, pain in joint of upper arm, pain in joint of wrist, hypermobility syndrome, cervical sprain/strain, myofascial pain. Subjective complaints include pain in left shoulder, right elbow and neck. Objective findings include reduced range of motion in left shoulder, tenderness to palpation of left trap and right lateral epicondyle, cervical PSM, hypertonicity of left trapezoid. Treatment has consisted of physical therapy, yoga, TENS, Terocin, Omeprazole and Diclofenac. The utilization review determination was rendered on 11/05/2014 recommending non-certification of 3 Terocin 120ml, Omeprazole 20mg #60, Diclofenac ER 100mg #60.

IMR ISSUES, DECISIONS AND RATIONALES

The Final Determination was based on decisions for the disputed items/services set forth below:

3 Terocin 120ml: Upheld

Claims Administrator guideline: The Claims Administrator did not cite any medical evidence for its decision.

MAXIMUS guideline: Decision based on MTUS Chronic Pain Treatment Guidelines Topical Analgesics Page(s): 111-113. Decision based on Non-MTUS Citation Official Disability Guidelines (ODG) Pain, Compound creams

Decision rationale: MTUS and ODG recommend usage of topical analgesics as an option, but also further details "primarily recommended for neuropathic pain when trials of antidepressants and anticonvulsants have failed." The medical documents do not indicate failure of antidepressants or anticonvulsants. MTUS states, "There is little to no research to support the use of many of these agents. Any compounded product that contains at least one drug (or drug class) that is not recommended is not recommended." Terocin lotion is topical pain lotion that contains lidocaine and menthol. ODG states regarding lidocaine topical patch, "This is not a first-line treatment and is only FDA approved for post-herpetic neuralgia." Medical documents do not document the patient as having post-herpetic neuralgia. Additionally, Topical analgesics are primarily recommended for neuropathic pain when trials of antidepressants and anticonvulsants have failed. The treating physician did not document a trial of first line agents and the objective outcomes of these treatments. MTUS states regarding topical analgesic creams, "There is little to no research to support the use of many of these agents. Any compounded product that contains at least one drug (or drug class) that is not recommended is not recommended." In this case, topical lidocaine is not indicated. As such, 3 Terocin 120ml is not medically necessary.

Omeprazole 20mg #60: Upheld

Claims Administrator guideline: The Claims Administrator did not cite any medical evidence for its decision.

MAXIMUS guideline: Decision based on MTUS Chronic Pain Treatment Guidelines NSAIDs, GI symptoms & cardiovascular risk Page(s): 68-69. Decision based on Non-MTUS Citation Official Disability Guidelines (ODG) Pain (Chronic), NSAIDs, GI symptoms & cardiovascular risk

Decision rationale: MTUS and ODG states, "Determine if the patient is at risk for gastrointestinal events: (1) age > 65 years; (2) history of peptic ulcer, GI bleeding or perforation; (3) concurrent use of ASA, corticosteroids, and/or an anticoagulant; or (4) high dose/multiple NSAID (e.g., NSAID + low-dose ASA)." And "Patients at intermediate risk for gastrointestinal events and no cardiovascular disease: (1) A non-selective NSAID with either a PPI (Proton Pump Inhibitor, for example, 20 mg omeprazole daily) or misoprostol (200 g four times daily) or (2) a Cox-2 selective agent. Long-term PPI use (> 1 year) has been shown to increase the risk of hip fracture (adjusted odds ratio 1.44)." The medical documents provided do not establish the patient has having documented GI bleeding, perforation, peptic ulcer, high dose NSAID, or other GI risk factors as outlined in MTUS. As such, the request for Omeprazole 20mg #60 is not medically necessary.

Diclofenac ER 100mg #60: Upheld

Claims Administrator guideline: The Claims Administrator did not cite any medical evidence for its decision.

MAXIMUS guideline: Decision based on MTUS Chronic Pain Treatment Guidelines NSAIDs (non-steroidal anti-inflammatory drugs) Page(s): 67-73. Decision based on Non-MTUS Citation Official Disability Guidelines (ODG) Pain (Chronic), Diclofenac

Decision rationale: Voltaran/Zipsor is the name brand version of Diclofenac, which is a NSAID. MTUS specifies four recommendations regarding NSAID use: 1) Osteoarthritis (including knee and hip): Recommended at the lowest dose for the shortest period in patients with moderate to severe pain. 2) Back Pain - Acute exacerbations of chronic pain: Recommended as a second-line treatment after acetaminophen. In general, there is conflicting evidence that NSAIDs are more effective than acetaminophen for acute LBP. 3) Back Pain - Chronic low back pain: Recommended as an option for short-term symptomatic relief. A Cochrane review of the literature on drug relief for low back pain (LBP) suggested that NSAIDs were no more effective than other drugs such as acetaminophen, narcotic analgesics, and muscle relaxants. The review also found that NSAIDs had more adverse effects than placebo and acetaminophen but fewer effects than muscle relaxants and narcotic analgesics. 4) Neuropathic pain: There is inconsistent evidence for the use of these medications to treat long term neuropathic pain, but they may be useful to treat breakthrough and mixed pain conditions such as osteoarthritis (and other nociceptive pain) in with neuropathic pain. The medical documents do not indicate that the patient is being treated for osteoarthritis. The treating physician does not document failure of primary (Tylenol) treatment. Importantly, ODG also states that diclofenac is "Not recommended as first line due to increased risk profile . . . If using diclofenac then consider discontinuing as it should only be used for the shortest duration possible in the lowest effective dose due to reported serious adverse events." Medical documents indicate that the patient has been on diclofenac since at least 06/20/2014, which given the treatment history does not appear to be the shortest duration possible. As such, the request Diclofenac ER 100mg #60 is not medically necessary.