

| | | | |
|-----------------------|--------------|------------------------------|------------|
| Case Number: | CM14-0186963 | | |
| Date Assigned: | 11/17/2014 | Date of Injury: | 03/18/2001 |
| Decision Date: | 01/05/2015 | UR Denial Date: | 10/18/2014 |
| Priority: | Standard | Application Received: | 11/10/2014 |

HOW THE IMR FINAL DETERMINATION WAS MADE

MAXIMUS Federal Services sent the complete case file to an expert reviewer. He/she has no affiliation with the employer, employee, providers or the claims administrator. The expert reviewer is Board Certified in Anesthesia, has a subspecialty in Pain Medicine and is licensed to practice in Massachusetts. He/she has been in active clinical practice for more than five years and is currently working at least 24 hours a week in active practice. The expert reviewer was selected based on his/her clinical experience, education, background, and expertise in the same or similar specialties that evaluate and/or treat the medical condition and disputed items/services. He/she is familiar with governing laws and regulations, including the strength of evidence hierarchy that applies to Independent Medical Review determinations.

CLINICAL CASE SUMMARY

The expert reviewer developed the following clinical case summary based on a review of the case file, including all medical records:

According to the documents available for review, the patient is a 47 year old male. The date of injury is 3/18/2001. The patient sustained an injury to the lumbar spine. The specific mechanism of injury was not fully elaborated on in the notes available for review. The patient currently complains of pain in the low back. The patient has undergone a lumbar spine fusion. The patient is maintained on the multimodal pain medication regimen including Ambien and Percocet . A request for Ambien and Percocet was denied.

IMR ISSUES, DECISIONS AND RATIONALES

The Final Determination was based on decisions for the disputed items/services set forth below:

1 Prescription for Ambien 10mg #30: Upheld

Claims Administrator guideline: The Claims Administrator did not base their decision on the MTUS. Decision based on Non-MTUS Citation Official Disability Guidelines

MAXIMUS guideline: The Expert Reviewer did not base their decision on the MTUS. Decision based on Non-MTUS Citation Official Disability Guidelines (ODG) Pain (Chronic) Ambien (Zolpidem)

Decision rationale: Zolpidem is a prescription short-acting nonbenzodiazepine hypnotic, which is approved for the short-term (usually two to six weeks) treatment of insomnia. Proper sleep hygiene is critical to the individual with chronic pain and often is hard to obtain. Various

medications may provide short-term benefit. While sleeping pills, so-called minor tranquilizers, and anti-anxiety agents are commonly prescribed in chronic pain, pain specialists rarely, if ever, recommend them for long-term use. They can be habit-forming, and they may impair function and memory more than opioid pain relievers. There is also concern that they may increase pain and depression over the long-term. (Feinberg, 2008) See Insomnia treatment. Ambien CR offers no significant clinical advantage over regular release Zolpidem. Ambien CR is approved for chronic use, but chronic use of hypnotics in general is discouraged, as outlined in Insomnia treatment. Ambien CR causes a greater frequency of dizziness, drowsiness, and headache compared to immediate release Zolpidem. (Ambien & Ambien CR package insert) Cognitive behavioral therapy (CBT) should be an important part of an insomnia treatment plan. A study of patients with persistent insomnia found that the addition of Zolpidem immediate release to CBT was modestly beneficial during acute (first 6 weeks) therapy, but better long-term outcomes were achieved when Zolpidem IR was discontinued and maintenance CBT continued. (Morin, 2009) Due to adverse effects, FDA now requires lower doses for Zolpidem. The dose of Zolpidem for women should be lowered from 10 mg to 5 mg for IR products (Ambien, Edluar, Zolpimist, and generic) and from 12.5 mg to 6.25 mg for ER products (Ambien CR). The ER product is still more risky than IR. In laboratory studies, 15% of women and 3% of men who took a 10-milligram dose of Ambien had potentially dangerous concentrations of the drug in their blood eight hours later. Among those who took Ambien CR, the problem was more common: 33% of women and 25% of men had blood concentrations that would raise the risk of a motor vehicle accident eight hours later. Even at the lower dose of Ambien CR now recommended by the FDA, 15% of women and 5% of men still had high levels of the drug in their system in the morning. (FDA, 2013) According to SAMHSA, Zolpidem is linked to a sharp increase in ED visits, so it should be used safely for only a short period of time. According to the documents available for review, the patient does not carry diagnoses of insomnia. Furthermore the patient has been using this medication for long-term treatment. Therefore, at this time, the requirements for treatment have not been met and medical necessity for Ambien is not established.

1 Prescription for percocet 10/325mg #150: Upheld

Claims Administrator guideline: Decision based on MTUS Chronic Pain Treatment Guidelines.

MAXIMUS guideline: Decision based on MTUS Chronic Pain Treatment Guidelines Opioids, On-Going Management Page(s): 74-97.

Decision rationale: According to the MTUS Chronic Pain Medical Treatment Guidelines section on Opioids, On-Going Management, p 74-97, (a) Prescriptions from a single practitioner taken as directed, and all prescriptions from a single pharmacy. (b) The lowest possible dose should be prescribed to improve pain and function. (c) Office: Ongoing review and documentation of pain relief, functional status, appropriate medication use, and side effects. Pain assessment should include: current pain; the least reported pain over the period since last assessment; average pain; intensity of pain after taking the opioid; how long it takes for pain relief; and how long pain relief lasts. Satisfactory response to treatment may be indicated by the patient's decreased pain, increased level of function, or improved quality of life. Information from family members or other caregivers should be considered in determining the patient's

response to treatment. The 4 A's for Ongoing Monitoring: Four domains have been proposed as most relevant for ongoing monitoring of chronic pain patients on opioids: pain relief, side effects, physical and psychosocial functioning, and the occurrence of any potentially aberrant (or nonadherent) drug-related behaviors. These domains have been summarized as the "4 A's" (analgesia, activities of daily living, adverse side effects, and aberrant drug taking behaviors). The monitoring of these outcomes over time should affect therapeutic decisions and provide a framework for documentation of the clinical use of these controlled drugs. (Passik, 2000)

(d) Home: To aid in pain and functioning assessment, the patient should be requested to keep a pain diary that includes entries such as pain triggers, and incidence of end-of-dose pain. It should be emphasized that using this diary will help in tailoring the opioid dose. This should not be a requirement for pain management.

(e) Use of drug screening or inpatient treatment with issues of abuse, addiction, or poor pain control.

(f) Documentation of misuse of medications (doctor-shopping, uncontrolled drug escalation, drug diversion).

(g) Continuing review of overall situation with regard to nonopioid means of pain control.

(h) Consideration of a consultation with a multidisciplinary pain clinic if doses of opioids are required beyond what is usually required for the condition or pain does not improve on opioids in 3 months. Consider a psych consult if there is evidence of depression, anxiety or irritability. Additionally, the MTUS states that continued use of opioids requires (a) the patient has returned to work, (b) the patient has improved functioning and pain. There is no current documentation of baseline pain, pain score with use of opioids, functional improvement on current regimen, side effects or review of potentially aberrant drug taking behaviors as outlined in the MTUS and as required for ongoing treatment. Therefore, at this time, the requirements for treatment have not been met and medical necessity for Percocet is not established.