

Case Number:	CM14-0187020		
Date Assigned:	11/18/2014	Date of Injury:	09/13/2008
Decision Date:	01/06/2015	UR Denial Date:	10/07/2014
Priority:	Standard	Application Received:	11/10/2014

HOW THE IMR FINAL DETERMINATION WAS MADE

MAXIMUS Federal Services sent the complete case file to an expert reviewer. He/she has no affiliation with the employer, employee, providers or the claims administrator. The expert reviewer is Board Certified in General Preventive Medicine and is licensed to practice in Indiana. He/she has been in active clinical practice for more than five years and is currently working at least 24 hours a week in active practice. The expert reviewer was selected based on his/her clinical experience, education, background, and expertise in the same or similar specialties that evaluate and/or treat the medical condition and disputed items/services. He/she is familiar with governing laws and regulations, including the strength of evidence hierarchy that applies to Independent Medical Review determinations.

CLINICAL CASE SUMMARY

The expert reviewer developed the following clinical case summary based on a review of the case file, including all medical records:

This employee is a 58 year old female with date of injury of 9/13/2008. A review of the medical records indicates that the patient is undergoing treatment for intervertebral disc disease and lumbago. Subjective complaints include continued 8/10 lower back pain with radiation and numbness down the left leg. Objective findings include limited range of motion of the lumbar spine with tenderness to palpation of the paravertebrals and a positive straight leg raise bilaterally. Treatment has included Tramadol, Naproxen, and Omeprazole. The utilization review dated 10/7/2014 partially-certified Tramadol 50mg #60, Naproxen 550mg #60, and Omeprazole 20mg #60.

IMR ISSUES, DECISIONS AND RATIONALES

The Final Determination was based on decisions for the disputed items/services set forth below:

Tramadol 50mg #60: Upheld

Claims Administrator guideline: Decision based on MTUS Chronic Pain Treatment Guidelines Opioids Page(s): 91.

MAXIMUS guideline: Decision based on MTUS Chronic Pain Treatment Guidelines Opioids, Tramadol, Ultram Page(s): 74-96, 113, 123. Decision based on Non-MTUS Citation Official Disability Guidelines (ODG) Pain (Chronic) - Medications for acute pain (analgesics), Tramadol (Ultram®)

Decision rationale: Ultram is the brand name version of tramadol, which is classified as central acting synthetic opioids. California MTUS states regarding tramadol that "A therapeutic trial of opioids should not be employed until the patient has failed a trial of non-opioid analgesics. Before initiating therapy, the patient should set goals, and the continued use of opioids should be contingent on meeting these goals." Official Disability Guidelines (ODG) further states, "Tramadol is not recommended as a first-line oral analgesic because of its inferior efficacy to a combination of Hydrocodone/ acetaminophen." The treating physician did not provide sufficient documentation that the patient has failed a trial of non-opioid analgesics at the time of prescription or in subsequent medical notes. Additionally, no documentation was provided which discussed the setting of goals for the use of tramadol prior to the initiation of this medication. The original utilization review recommended weaning and modified the request, which is appropriate. As such, the request for tramadol #60 is not medically necessary.

Naproxen Sodium 550mg #60: Upheld

Claims Administrator guideline: Decision based on MTUS Chronic Pain Treatment Guidelines NSAIDS.

MAXIMUS guideline: Decision based on MTUS Chronic Pain Treatment Guidelines NSAIDs (non-steroidal anti-inflammatory drugs) Page(s): 67-73. Decision based on Non-MTUS Citation Official Disability Guidelines (ODG) Pain (Chronic), Naproxen, NSAIDs (non-steroidal anti-inflammatory drugs)

Decision rationale: California MTUS recommends NSAIDs for osteoarthritis "at the lowest dose for the shortest period in patients with moderate to severe pain. Acetaminophen may be considered for initial therapy for patients with mild to moderate pain, and in particular, for those with gastrointestinal, cardiovascular or renovascular risk factors. NSAIDs appear to be superior to acetaminophen, particularly for patients with moderate to severe pain. There is no evidence to recommend one drug in this class over another based on efficacy." MTUS further specifies that NSAIDs should be used cautiously in patients with hypertension. Official Disability Guidelines (ODG) states, "Recommended as an option. Naproxen is a non-steroidal anti-inflammatory drug (NSAID) for the relief of the signs and symptoms of osteoarthritis." The employee does not have osteoarthritis, and there is no documentation showing a failure of acetaminophen. Therefore, the request for Naproxyn is not medically necessary.

Omeprazole 20mg #60: Upheld

Claims Administrator guideline: Decision based on MTUS Chronic Pain Treatment Guidelines NSAIDS.

MAXIMUS guideline: Decision based on MTUS Chronic Pain Treatment Guidelines NSAIDs, GI symptoms & cardiovascular risk Page(s): 68-69. Decision based on Non-MTUS Citation Official Disability Guidelines (ODG) Pain (Chronic), NSAIDs, GI symptoms & cardiovascular risk

Decision rationale: California MTUS and Official Disability Guidelines ODG states, "Determine if the patient is at risk for gastrointestinal events: (1) age > 65 years; (2) history of peptic ulcer, GI bleeding or perforation; (3) concurrent use of ASA, corticosteroids, and/or an anticoagulant; or (4) high dose/multiple NSAID (e.g., NSAID + low-dose ASA)." And "Patients at intermediate risk for gastrointestinal events and no cardiovascular disease:(1) A non-selective NSAID with either a PPI (Proton Pump Inhibitor, for example, 20 mg omeprazole daily) or misoprostol (200 four times daily) or(2) a Cox-2 selective agent. Long-term PPI use (> 1 year) has been shown to increase the risk of hip fracture (adjusted odds ratio 1.44)." The medical documents provided do not establish the patient as having documented GI bleeding/perforation/peptic ulcer or other GI risk factors as outlined in MTUS. As such, the request for Omeprazole 20mg is not medically necessary.