

Case Number:	CM14-0195940		
Date Assigned:	12/03/2014	Date of Injury:	09/03/2008
Decision Date:	01/20/2015	UR Denial Date:	10/27/2014
Priority:	Standard	Application Received:	11/23/2014

HOW THE IMR FINAL DETERMINATION WAS MADE

MAXIMUS Federal Services sent the complete case file to an expert reviewer. He/she has no affiliation with the employer, employee, providers or the claims administrator. The expert reviewer is Board Certified in Emergency Medicine and is licensed to practice in New York. He/she has been in active clinical practice for more than five years and is currently working at least 24 hours a week in active practice. The expert reviewer was selected based on his/her clinical experience, education, background, and expertise in the same or similar specialties that evaluate and/or treat the medical condition and disputed items/services. He/she is familiar with governing laws and regulations, including the strength of evidence hierarchy that applies to Independent Medical Review determinations.

CLINICAL CASE SUMMARY

The expert reviewer developed the following clinical case summary based on a review of the case file, including all medical records:

The injured worker suffered a work related injury on 09/03/08. Per the physician notes from 09/26/14 diagnoses include discogenic lumbar condition with 3 prior MRIs which showed disc herniation at L5-S1, ankle inflammation status post arthroscopy in 2009 with no improvement, and depression. The injured worker presented with a cane for walking. He complains of left ankle pain, described as constant, motion loss, stiffness, swelling, buckling popping and limping. He is status post 2 ESI with no documented improvement. He was recommended to have a back brace, hot and cold wrap, and a TENS unit, a pool membership, medications, Hyalgan injection to the ankle, and rocker bottom shoe. The current recommendation is for Norco, Flexeril, and Tramadol. These treatments were denied by the Claims Administrator and were subsequently appealed for Independent Medical Review.

IMR ISSUES, DECISIONS AND RATIONALES

The Final Determination was based on decisions for the disputed items/services set forth below:

Norco #60: Upheld

Claims Administrator guideline: Decision based on MTUS Chronic Pain Treatment Guidelines Opioids.

MAXIMUS guideline: Decision based on MTUS Chronic Pain Treatment Guidelines Pain Interventions and Guidelines Page(s): 11, 74-96.

Decision rationale: Norco is the compounded medication containing hydrocodone and acetaminophen. Chronic Pain Medical Treatment Guidelines state that opioids are not recommended as a first line therapy. Opioid should be part of a treatment plan specific for the patient and should follow criteria for use. Criteria for use include establishment of a treatment plan, determination if pain is nociceptive or neuropathic, failure of pain relief with non-opioid analgesics, setting of specific functional goals, and opioid contract with agreement for random drug testing. If analgesia is not obtained, opioids should be discontinued. The patient should be screened for likelihood that he or she could be weaned from the opioids if there is no improvement in pain or function. It is recommended for short term use if first-line options, such as acetaminophen or NSAIDs have failed. Opioids may be a safer choice for patients with cardiac and renal disease than antidepressants or anticonvulsants. Acetaminophen is recommended for treatment of chronic pain & acute exacerbations of chronic pain. Acetaminophen overdose is a well-known cause of acute liver failure. Hepatotoxicity from therapeutic doses is unusual. Renal insufficiency occurs in 1 to 2% of patients with overdose. The recommended dose for mild to moderate pain is 650 to 1000 mg orally every 4 hours with a maximum of 4 g/day. In this case the patient has been using Norco since at least January 2013 and has not obtained analgesia. In addition there is no documentation that the patient has signed an opioid contract or is participating in urine drug testing. Criteria for long-term opioid use have not been met. The request is not medically necessary.

Flexeril 10MG #60: Upheld

Claims Administrator guideline: Decision based on MTUS Chronic Pain Treatment Guidelines Anti-Epilepsy drugs.

MAXIMUS guideline: Decision based on MTUS Chronic Pain Treatment Guidelines Pain Interventions and Guidelines Page(s): 63.

Decision rationale: Flexeril is the muscle relaxant (Cyclobenzaprine). Cyclobenzaprine is recommended as an option, for a short course of therapy. It has been found to be more effective than placebo with greater adverse side effects. Its greatest effect is in the first 4 days. Treatment should be brief. Non-sedating muscle relaxants are recommended with caution as a second-line option for short-term treatment (less than two weeks) of acute exacerbations in patients with chronic LBP. Muscle relaxants may be effective in reducing pain and muscle tension, and increasing mobility. However, in most LBP cases, they show no benefit beyond NSAIDs in pain and overall improvement. Also there is no additional benefit shown in combination with NSAIDs. Efficacy appears to diminish over time, and prolonged use of some medications in this class may lead to dependence. Sedation is the most commonly reported adverse effect of muscle relaxant medications. These drugs should be used with caution in patients driving motor vehicles or operating heavy machinery. In this case the patient had been using Flexeril since at least January 2013. The duration of treatment surpasses the recommended short-term duration of two weeks. The request is not medically necessary.

Tramadol ER 150mg #30: Upheld

Claims Administrator guideline: Decision based on MTUS Chronic Pain Treatment Guidelines Opioids Page(s): 93-94.

MAXIMUS guideline: Decision based on MTUS Chronic Pain Treatment Guidelines Pain Interventions and Guidelines Page(s): 74-96.

Decision rationale: Tramadol is a synthetic opioid affecting the central nervous system. It has several side effects, which include increasing the risk of seizure in patients taking SSRI's, TCA's and other opioids. Chronic Pain Medical Treatment Guidelines state that opioids are not recommended as a first line therapy. Opioid should be part of a treatment plan specific for the patient and should follow criteria for use. Criteria for use include establishment of a treatment plan, determination if pain is nociceptive or neuropathic, failure of pain relief with non-opioid analgesics, setting of specific functional goals, and opioid contract with agreement for random drug testing. If analgesia is not obtained, opioids should be discontinued. The patient should be screened for likelihood that he or she could be weaned from the opioids if there is no improvement in pain or function. It is recommended for short-term use if first-line options, such as acetaminophen or NSAIDS have failed. In this case the medication was not prescribed for short term use and the criteria for opioid use were not met. In this case the patient had been taking opioids since at least January 2013 and had not obtained analgesia. In addition there is no documentation that the patient has signed an opioid contract or is participating in urine drug testing. Criteria for long-term opioid use have not been met. The request is not medically necessary.