

Case Number:	CM14-0196754		
Date Assigned:	12/04/2014	Date of Injury:	11/01/2008
Decision Date:	01/23/2015	UR Denial Date:	10/28/2014
Priority:	Standard	Application Received:	11/24/2014

HOW THE IMR FINAL DETERMINATION WAS MADE

MAXIMUS Federal Services sent the complete case file to an expert reviewer. He/she has no affiliation with the employer, employee, providers or the claims administrator. The expert reviewer is Board Certified in Internal Medicine and is licensed to practice in District of Columbia. He/she has been in active clinical practice for more than five years and is currently working at least 24 hours a week in active practice. The expert reviewer was selected based on his/her clinical experience, education, background, and expertise in the same or similar specialties that evaluate and/or treat the medical condition and disputed items/services. He/she is familiar with governing laws and regulations, including the strength of evidence hierarchy that applies to Independent Medical Review determinations.

CLINICAL CASE SUMMARY

The expert reviewer developed the following clinical case summary based on a review of the case file, including all medical records:

This is a 54 year old patient who sustained injury on Nov 1 2008. He developed moderate to severe lower back pain, bilateral hand pain and bilateral shoulder pain and neck pain. He also had issues with insomnia. It was recommended that the patient have aqua therapy of 18 sessions.

IMR ISSUES, DECISIONS AND RATIONALES

The Final Determination was based on decisions for the disputed items/services set forth below:

Aquatic Therapy 3 x 6: Overturned

Claims Administrator guideline: Decision based on MTUS Chronic Pain Treatment Guidelines Aquatic Therapy.

MAXIMUS guideline: Decision based on MTUS Chronic Pain Treatment Guidelines Page(s): 22,46-47.

Decision rationale: Per MTUS, it is recommended as an optional form of exercise therapy, where available, as an alternative to land-based physical therapy. Aquatic therapy (including swimming) can minimize the effects of gravity, so it is specifically recommended where reduced weight bearing is desirable, for example extreme obesity. For recommendations on the number of supervised visits, see Physical medicine. Water exercise improved some components of health-related quality of life, balance, and stair climbing in females with fibromyalgia, but regular exercise and higher intensities may be required to preserve most of these gains. (Tomas-

Carus, 2007). Exercise: Exercise is recommended. There is strong evidence that exercise programs, including aerobic conditioning and strengthening, is superior to treatment programs that do not include exercise. There is no sufficient evidence to support the recommendation of any particular exercise regimen over any other exercise regimen. A therapeutic exercise program should be initiated at the start of any treatment or rehabilitation program, unless exercise is contraindicated. Such programs should emphasize education, independence, and the importance of an on-going exercise regime. (State, 2002) (Airaksinen, 2006) A recent study of the long term impact of aerobic exercise on musculoskeletal pain, in a prospective cohort of 866 healthy seniors followed for 14 years, found that exercise was associated with a substantial and significant reduction in pain even after adjusting for gender, baseline BMI and attrition, and despite the fact that fractures, a significant predictor of pain, were slightly more common among exercisers. (Bruce, 2005) A recent trial concluded that active physical treatment, cognitive-behavioral treatment, and the two combined each resulted in equally significant improvement, much better compared to no treatment. (The cognitive treatment focused on encouraging increased physical activity.) (Smeets, 2006) Progressive walking, simple strength training, and stretching improved functional status, key symptoms, and self-efficacy in patients with fibromyalgia. (Rooks, 2007) Physical conditioning in chronic pain patients can have immediate and long-term benefits, according to a low-quality study presented at the American Academy of Pain Medicine 24th Annual Meeting. (Burlinson, 2008) Physical therapy in warm-water has been effective and highly recommended in persons with fibromyalgia. In this RCT, an aquatic exercise program including one-hour, supervised, water-based exercise sessions, three times per week for 8 months, was found to be cost-effective in terms of both health care costs and societal costs. (Gusi, 2008) An educational technique known as the Alexander technique, along with exercise, is effective for long-term relief of chronic low back pain, according to the results of a randomized trial reported in the BMJ. (Little, 2008). Per guidelines cited, the patient could receive this therapy if thought to be clinically helpful. The recommended number of visits is also within the parameters suggested by MTUS. Therefore, the request is medically necessary.