

Case Number:	CM14-0197421		
Date Assigned:	12/05/2014	Date of Injury:	08/07/2000
Decision Date:	01/28/2015	UR Denial Date:	10/27/2014
Priority:	Standard	Application Received:	11/24/2014

HOW THE IMR FINAL DETERMINATION WAS MADE

MAXIMUS Federal Services sent the complete case file to an expert reviewer. He/she has no affiliation with the employer, employee, providers or the claims administrator. The expert reviewer is Board Certified in Anesthesiology, has a subspecialty in Pain medicine and Acupuncture and is licensed to practice in California. He/she has been in active clinical practice for more than five years and is currently working at least 24 hours a week in active practice. The expert reviewer was selected based on his/her clinical experience, education, background, and expertise in the same or similar specialties that evaluate and/or treat the medical condition and disputed items/services. He/she is familiar with governing laws and regulations, including the strength of evidence hierarchy that applies to Independent Medical Review determinations.

CLINICAL CASE SUMMARY

The expert reviewer developed the following clinical case summary based on a review of the case file, including all medical records:

The California MTUS is silent on medial branch blocks. Per the ODG guidelines, facet joint medial branch blocks are not recommended except as a diagnostic tool, citing minimal evidence for treatment. The ODG indicates that criteria for facet joint diagnostic blocks (injections) are as follows: 1. One set of diagnostic medial branch blocks is required with a response of 70%. The pain response should last at least 2 hours for Lidocaine. 2. Limited to patients with low-back pain that is non-radicular and at no more than two levels bilaterally. 3. There is documentation of failure of conservative treatment (including home exercise, PT and NSAIDs) prior to the procedure for at least 4-6 weeks. 4. No more than 2 facet joint levels are injected in one session (see above for medial branch block levels). 5. Recommended volume of no more than 0.5 cc of injectate is given to each joint. 6. No pain medication from home should be taken for at least 4 hours prior to the diagnostic block and for 4 to 6 hours afterward. 7. Opioids should not be given as a "sedative" during the procedure. 8. The use of IV sedation (including other agents such as midazolam) may be grounds to negate the results of a diagnostic block, and should only be given in cases of extreme anxiety. 9. The patient should document pain relief with an instrument such as a VAS scale, emphasizing the importance of recording the maximum pain relief and maximum duration of pain. The patient should also keep medication use and activity logs to support subjective reports of better pain control. 10. Diagnostic facet blocks should not be performed in patients in whom a surgical procedure is anticipated. (Resnick, 2005) 11. Diagnostic facet blocks should not be performed in patients who have had a previous fusion procedure at the planned injection level. [Exclusion Criteria that would require UR physician review: Previous fusion at the targeted level. (Franklin, 2008)] Since the request is for greater than the recommended 2 facet joint levels at a time, the request is not medically necessary.

IMR ISSUES, DECISIONS AND RATIONALES

The Final Determination was based on decisions for the disputed items/services set forth below:

Left lumbar epidural steroid injection to L5-S1: Upheld

Claims Administrator guideline: Decision based on MTUS Chronic Pain Treatment Guidelines Epidural Steroid Injections (ESIs). Decision based on Non-MTUS Citation Official Disability Guidelines (ODG), Low Back, Lumbar & Thoracic (Acute & Chronic)

MAXIMUS guideline: Decision based on MTUS Chronic Pain Treatment Guidelines Epidural Steroid Injections Page(s): 46.

Decision rationale: Per the California MTUS Chronic Pain Medical Treatment Guidelines epidural steroid injections are used to reduce pain and inflammation, restoring range of motion and thereby facilitating progress in more active treatment programs and avoiding surgery, but this treatment alone offers no significant long-term benefit. The criteria for the use of epidural steroid injections are as follows: 1) Radiculopathy must be documented by physical examination and corroborated by imaging studies and/or electrodiagnostic testing. 2) Initially unresponsive to conservative treatment (exercises, physical methods, NSAIDs and muscle relaxants). 3) Injections should be performed using fluoroscopy (live x-ray) for guidance. 4) If used for diagnostic purposes, a maximum of two injections should be performed. A second block is not recommended if there is inadequate response to the first block. Diagnostic blocks should be at an interval of at least one to two weeks between injections. 5) No more than two nerve root levels should be injected using transforaminal blocks. 6) No more than one interlaminar level should be injected at one session. 7) In the therapeutic phase, repeat blocks should be based on continued objective documented pain and functional improvement, including at least 50% pain relief with associated reduction of medication use for six to eight weeks, with a general recommendation of no more than 4 blocks per region per year. (Manchikanti, 2003) (CMS, 2004) (Boswell, 2007) 8) Current research does not support a "series-of-three" injections in either the diagnostic or therapeutic phase. We recommend no more than 2 ESI injections. The documentation submitted for review indicates that the injured worker's previous epidural injection provided an improvement of 40%, and there was no documentation of an associated reduction of medication use. As the criteria for repeat block is not met, the request is not medically necessary.