

Case Number:	CM14-0198088		
Date Assigned:	12/08/2014	Date of Injury:	09/25/1997
Decision Date:	01/30/2015	UR Denial Date:	10/31/2014
Priority:	Standard	Application Received:	11/25/2014

HOW THE IMR FINAL DETERMINATION WAS MADE

MAXIMUS Federal Services sent the complete case file to an expert reviewer. He/she has no affiliation with the employer, employee, providers or the claims administrator. The expert reviewer is Board Certified in Family Practice, and is licensed to practice in California. He/she has been in active clinical practice for more than five years and is currently working at least 24 hours a week in active practice. The expert reviewer was selected based on his/her clinical experience, education, background, and expertise in the same or similar specialties that evaluate and/or treat the medical condition and disputed items/services. He/she is familiar with governing laws and regulations, including the strength of evidence hierarchy that applies to Independent Medical Review determinations.

CLINICAL CASE SUMMARY

The expert reviewer developed the following clinical case summary based on a review of the case file, including all medical records:

Patient had a date of injury on 9/25/1997. Diagnosis includes: occipital neuralgia, cervicgia, cervical degenerative disc disease, chronic pain syndrome, and shoulder pain.

IMR ISSUES, DECISIONS AND RATIONALES

The Final Determination was based on decisions for the disputed items/services set forth below:

Terocin Lotion 2.5-0.025-10-25% with 1 refill: Upheld

Claims Administrator guideline: Decision based on MTUS Chronic Pain Treatment Guidelines.

MAXIMUS guideline: Decision based on MTUS Chronic Pain Treatment Guidelines Topical analgesics Page(s): 111 and 112.

Decision rationale: According to guidelines topical analgesic are largely experimental in use with few randomized controlled trials to determine efficacy or safety. Primarily recommended for neuropathic pain when trials of antidepressants and anticonvulsants have failed. There is little to no research to support the use of many of these agents. Any compounded product that contains at least one drug (or drug class) that is not recommended is not recommended. Terocin contains Menthol which is not supported and thus not medically necessary.

Celebrex 200mg #60 with 1 refill: Upheld

Claims Administrator guideline: Decision based on MTUS Chronic Pain Treatment Guidelines.

MAXIMUS guideline: Decision based on MTUS Chronic Pain Treatment Guidelines NSAIDS Page(s): 67.

Decision rationale: According to guidelines NSAIDs are used for Osteoarthritis (including knee and hip): Recommended at the lowest dose for the shortest period in patients with moderate to severe pain. Acetaminophen may be considered for initial therapy for patients with mild to moderate pain, and in particular, for those with gastrointestinal, cardiovascular or renovascular risk factors. NSAIDs appear to be superior to acetaminophen, particularly for patients with moderate to severe pain there should be functional improvement. According to the medical records there is no improvement with prolonged usage of NSAIDs and thus is not medically necessary.

Paxil 40mg #30: Upheld

Claims Administrator guideline: The Claims Administrator did not base their decision on the MTUS. Decision based on Non-MTUS Citation Official Disability Guidelines

MAXIMUS guideline: The Expert Reviewer did not base their decision on the MTUS. Decision based on Non-MTUS Citation Official Disability Guidelines (ODG), antidepressants.

Decision rationale: According to guidelines it states antidepressants can be used as an option for chronic pain. The patient has not tried other available treatment options and thus Paxil is not medically necessary.

Norco 10/325mg, #120: Upheld

Claims Administrator guideline: Decision based on MTUS Chronic Pain Treatment Guidelines.

MAXIMUS guideline: Decision based on MTUS Chronic Pain Treatment Guidelines Opioids Page(s): 76-82.

Decision rationale: According to guidelines it states opioids should only be continued if there is functional improvement. It also states chronic use of opioids can lead to dependence and addiction. According to the patient's medical records it does not state the patient has functional improvement with Norco usage and thus is not medically necessary.