

Case Number:	CM14-0200754		
Date Assigned:	12/11/2014	Date of Injury:	01/02/2013
Decision Date:	01/28/2015	UR Denial Date:	11/20/2014
Priority:	Standard	Application Received:	12/01/2014

HOW THE IMR FINAL DETERMINATION WAS MADE

MAXIMUS Federal Services sent the complete case file to an expert reviewer. He/she has no affiliation with the employer, employee, providers or the claims administrator. The expert reviewer is Board Certified in Neurology, has a subspecialty in Neuromuscular Medicine and is licensed to practice in New Jersey. He/she has been in active clinical practice for more than five years and is currently working at least 24 hours a week in active practice. The expert reviewer was selected based on his/her clinical experience, education, background, and expertise in the same or similar specialties that evaluate and/or treat the medical condition and disputed items/services. He/she is familiar with governing laws and regulations, including the strength of evidence hierarchy that applies to Independent Medical Review determinations.

CLINICAL CASE SUMMARY

The expert reviewer developed the following clinical case summary based on a review of the case file, including all medical records:

The patient is a 45-year-old woman who sustained a work-related injury on January 2, 2013. Subsequently, she developed hand pain. According to the orthopedic re-evaluation report dated November 11, 2014, the patient reported moderate to severe right 5th finger pain, tingling and restricted range of motion and flexion. The patient's right hand has a 5th finger that was not included when she made a fist. She lacked about 4 cm pulp to palm. It was very tight on the dorsal capsule and the extensor mechanism. The patient was diagnosed with posttraumatic arthrosis of the right 5th digit DIP joint, fracture dislocation of the right 5th digit DIP joint, anxiety and depression, chronic pain, insomnia, and weakness of flexor digitorum profundus of the 5th finger on the right hand. The provider requested authorization for Prilosec, Xanax, Naproxen, Tramadol, Topical cream of Gabapentin, Ketoprofen and Tramadol, urine toxicology, and follow-up with hand specialist.

IMR ISSUES, DECISIONS AND RATIONALES

The Final Determination was based on decisions for the disputed items/services set forth below:

Prilosec 20mg, #90: Upheld

Claims Administrator guideline: Decision based on MTUS Chronic Pain Treatment Guidelines Page(s): 68.

MAXIMUS guideline: Decision based on MTUS Chronic Pain Treatment Guidelines NSAIDs, GI symptoms & cardiovascular risk Page(s): 68.

Decision rationale: According to MTUS guidelines, Omeprazole is indicated when NSAID are used in patients with intermediate or high risk for gastrointestinal events . The risk for gastrointestinal events are: (1) age > 65 years; (2) history of peptic ulcer, GI bleeding or perforation; (3) concurrent use of ASA, corticosteroids, and/or an anticoagulant; or (4) high dose/multiple NSAID (e.g., NSAID + low-dose ASA). Recent studies tend to show that H. Pylori does not act synergistically with NSAIDS to develop gastroduodenal lesions. There is no documentation that the patient has GI issue that requires the use of Prilosec. There is no documentation in the patient's chart supporting that she is at intermediate or high risk for developing gastrointestinal events. Therefore, Prilosec 20mg #90 prescription is not medically necessary.

Xanax 1mg #60: Upheld

Claims Administrator guideline: Decision based on MTUS Chronic Pain Treatment Guidelines Page(s): 24.

MAXIMUS guideline: Decision based on MTUS Chronic Pain Treatment Guidelines Benzodiazepines Page(s): 24.

Decision rationale: According to MTUS guidelines, benzodiazepines are not recommended for long term use for pain management because of unproven long term efficacy and because of the risk of dependence. Most guidelines limit their use to 4 weeks. There is a report of anxiety and depression and the use and failure of antidepressant was not documented. Therefore the use of Xanax is not medically necessary.

Naproxen 550mg #60: Upheld

Claims Administrator guideline: Decision based on MTUS Chronic Pain Treatment Guidelines Page(s): 67.

MAXIMUS guideline: Decision based on MTUS Chronic Pain Treatment Guidelines NON SELECTIVE NSAIDS Page(s): 72.

Decision rationale: Naproxen (Naprosyn): delayed release (EC-Naprosyn), as Sodium salt (Anaprox, Anaprox DS, Aleve [otc]) Generic available; extended-release (Naprelan): 375 mg. Different dose strengths and formulations of the drug are not necessarily bioequivalent. Dosing Information: Osteoarthritis or ankylosing spondylitis: Dividing the daily dose into 3 doses versus 2 doses for immediate-release and delayed-release formulations generally does not affect response. Morning and evening doses do not have to be equal in size. The dose may be increased to 1500 mg/day of naproxyn for limited periods when a higher level of analgesic/anti-inflammatory activity is required (for up to 6 months). Naprosyn or naproxyn: 250-500 mg PO twice daily. Anaprox: 275-550 mg PO twice daily. (Total dose may be increased to 1650 mg a day for limited periods). EC-Naprosyn: 375 mg or 500 mg twice daily. The tablet should not be broken, crushed or chewed to maintain integrity of the enteric coating. Naprelan: Two 375 mg tablets (750 mg) PO once daily or two 500 mg tablets (1000 mg) once daily. If required (and a

lower dose was tolerated) Naprelan can be increased to 1500 mg once daily for limited periods (when higher analgesia is required). Pain: Naprosyn or naproxyn: 250-500 mg PO twice daily. The maximum dose on day one should not exceed 1250 mg and 1000 mg on subsequent days. Anaprox: 275-550 mg PO twice daily. The maximum dose on day one should not exceed 1375 mg and 1100 mg on subsequent days. Extended-release Naprelan: Not recommended due to delay in absorption. (Naprelan Package Insert) There is no documentation of the rationale behind the long-term use of Naproxen. NSAID should be used for the shortest duration and the lowest dose. There is no documentation from the patient file that the provider titrated Naproxen to the lowest effective dose and used it for the shortest period possible. Naproxen was used without clear documentation of its efficacy. Furthermore, there is no documentation that the provider followed the patient for NSAID adverse reactions that are not limited to GI side effect, but also may affect the renal function. Therefore, the request for Naproxen 550 mg #60 is not medically necessary.

Tramadol 150mg, two (2) times per day: Upheld

Claims Administrator guideline: Decision based on MTUS Chronic Pain Treatment Guidelines Page(s): 80.

MAXIMUS guideline: Decision based on MTUS Chronic Pain Treatment Guidelines Tramadol Page(s): 113.

Decision rationale: According to MTUS guidelines, Ultram (Tramadol) is a synthetic opioid indicated for the pain management but not recommended as a first line oral analgesic. In addition and according to MTUS guidelines, ongoing use of opioids should follow specific rules: (a) Prescriptions from a single practitioner taken as directed, and all prescriptions from a single pharmacy. (b) The lowest possible dose should be prescribed to improve pain and function. (c) Office: Ongoing review and documentation of pain relief, functional status, appropriate medication use, and side effects. Four domains have been proposed as most relevant for ongoing monitoring of chronic pain patients on opioids: pain relief, side effects, physical and psychosocial functioning, and the occurrence of any potentially aberrant (or non-adherent) drug-related behaviors. These domains have been summarized as the "4 A's" (analgesia, activities of daily living, adverse side effects, and aberrant drug taking behaviors). The monitoring of these outcomes over time should affect therapeutic decisions and provide a framework. Although, Tramadol may be needed to help with the patient pain, there is no clear evidence of objective and recent functional and pain improvement from its previous use. The patient has not been working for over 6 months. There is no objective documentation of pain severity level to justify the use of tramadol in this patient. There is no clear documentation of the efficacy/safety of previous use of tramadol. There is no recent evidence of objective monitoring of compliance of the patient with her medications. Therefore, the prescription of Tramadol 150 mg is not medically necessary.

Topical cream of Gabapentin, Ketoprofen and Tramadol: Upheld

Claims Administrator guideline: Decision based on MTUS Chronic Pain Treatment Guidelines Page(s): 111.

MAXIMUS guideline: Decision based on MTUS Chronic Pain Treatment Guidelines Topical Analgesics Page(s): 111.

Decision rationale: According to MTUS, in Chronic Pain Medical Treatment guidelines section Topical Analgesics (page 111), topical analgesics are largely experimental in use with few randomized controlled trials to determine efficacy or safety. Many agents are combined to other pain medications for pain control. That is limited research to support the use of many of these agents. Furthermore, according to MTUS guidelines, any compounded product that contains at least one drug or drug class that is not recommended is not recommended. There is no proven efficacy of topical application of the component of Ketoprofen. Furthermore, oral form of these medications was not attempted, and there is no documentation of failure or adverse reaction from first line pain medications. Based on the above, the use of Topical cream of Gabapentin, Ketoprofen and Tramadol is not medically necessary.

Urine Toxicology: Upheld

Claims Administrator guideline: The Claims Administrator did not base their decision on the MTUS. Decision based on Non-MTUS Citation Official Disability Guidelines (ODG), Pain

MAXIMUS guideline: Decision based on MTUS Chronic Pain Treatment Guidelines Opioids, steps to avoid misuse/addiction Page(s): 77-78; 94.

Decision rationale: According to MTUS guidelines, urine toxicology screens are indicated to avoid misuse/addiction. (j) Consider the use of a urine drug screen to assess for the use or the presence of illegal drugs. There is no evidence that the patient is taking any medication that requires a drug screen or has a history of use of illicit drugs. Therefore, the request for Urine drug screen is not medically necessary.

Follow-up with hand specialist: Upheld

Claims Administrator guideline: The Claims Administrator did not base their decision on the MTUS. Decision based on Non-MTUS Citation Official Disability Guidelines (ODG), Hand

MAXIMUS guideline: Decision based on MTUS Chronic Pain Treatment Guidelines Chronic pain programs, early intervention Page(s): 32-33.

Decision rationale: According to MTUS guidelines, the presence of red flags may indicate the need for specialty consultation. In addition, the requesting physician should provide a documentation supporting the medical necessity for a pain management evaluation with a specialist. The documentation should include the reasons, the specific goals and end point for using the expertise of a specialist. In this case, there is no clear documentation for the rationale for the request for an office visit for hand specialist. The requesting physician did not provide a documentation supporting the medical necessity for this visit. The provider documentation should include the reasons, the specific goals and end point for using the expertise of a specialist. Therefore, the request for follow-up hand specialist visit is not medically necessary.