

Case Number:	CM14-0200775		
Date Assigned:	12/11/2014	Date of Injury:	09/05/2008
Decision Date:	01/30/2015	UR Denial Date:	10/30/2014
Priority:	Standard	Application Received:	12/01/2014

HOW THE IMR FINAL DETERMINATION WAS MADE

MAXIMUS Federal Services sent the complete case file to an expert reviewer. He/she has no affiliation with the employer, employee, providers or the claims administrator. The expert reviewer is Board Certified in Neurology, has a subspecialty in Neuromuscular Medicine and is licensed to practice in New Jersey. He/she has been in active clinical practice for more than five years and is currently working at least 24 hours a week in active practice. The expert reviewer was selected based on his/her clinical experience, education, background, and expertise in the same or similar specialties that evaluate and/or treat the medical condition and disputed items/services. He/she is familiar with governing laws and regulations, including the strength of evidence hierarchy that applies to Independent Medical Review determinations.

CLINICAL CASE SUMMARY

The expert reviewer developed the following clinical case summary based on a review of the case file, including all medical records:

The patient is a 47 year old man who sustained a work-related injury on September 5 2008. Subsequently, the patient developed a chronic back pain for which he underwent lumbar laminectomy on 2011. The patient was treated with pain medications, physical therapy and chiropractic sessions as well as epidural injections and functional restoration program. His EMG performed on 2013 showed L5 radiculopathy. His MRI of the lumbar spine performed on 2013 showed lumbar disc protrusion. According to a progress report dated on October 14 2014, the patient was complaining of ongoing back pain. The patient physical examination demonstrated lumbar tenderness with reduced range of motion and positive straight raising test. The patient was diagnosed with post laminectomy syndrome, low back pain and degenerative disc disease. The provider requested authorization for Tramadol.

IMR ISSUES, DECISIONS AND RATIONALES

The Final Determination was based on decisions for the disputed items/services set forth below:

Tramadol HCL 50 mg: Upheld

Claims Administrator guideline: Decision based on MTUS Chronic Pain Treatment Guidelines Opioids.

MAXIMUS guideline: Decision based on MTUS Chronic Pain Treatment Guidelines Tramadol; Criteria for use of opioids Page(s): 113, 76-79.

Decision rationale: According to MTUS guidelines, Ultram is a synthetic opioid indicated for the pain management but not recommended as a first line oral analgesic. Although, Ultram may be needed to help with the patient pain, it may not help with the weaning process from opioids. Ultram could be used if exacerbation of pain after or during the weaning process. In addition and according to MTUS guidelines, ongoing use of opioids should follow specific rules:(a) Prescriptions from a single practitioner taken as directed, and all prescriptions from a single pharmacy.(b) The lowest possible dose should be prescribed to improve pain and function.(c) Office: Ongoing review and documentation of pain relief, functional status, appropriate medication use, and side effects. Pain assessment should include: current pain; the least reported pain over the period since last assessment; average pain; intensity of pain after taking the opioid; how long it takes for pain relief; and how long pain relief lasts. Satisfactory response to treatment may be indicated by the patient's decreased pain, increased level of function, or improved quality of life. Information from family members or other caregivers should be considered in determining the patient's response to treatment. The 4 A's for Ongoing Monitoring: Four domains have been proposed as most relevant for ongoing monitoring of chronic pain patients on opioids: pain relief, side effects, physical and psychosocial functioning, and the occurrence of any potentially aberrant (or non-adherent) drug-related behaviors. These domains have been summarized as the "4 A's" (analgesia, activities of daily living, adverse side effects, and aberrant drug taking behaviors). The monitoring of these outcomes over time should affect therapeutic decisions and provide a framework. There is no clear evidence of objective and recent functional and pain improvement with previous use of opioids (Tramadol). There no clear documentation of the efficacy/safety of previous use of Tramadol. There is no recent evidence of objective monitoring of compliance of the patient with his medication. There is no clear justification for the need to continue the use of Tramadol. Therefore, the prescription of 1 Prescription for Tramadol HCL 50 mg is not medically necessary at this time.

Lyrica 100 mg # 90: Upheld

Claims Administrator guideline: Decision based on MTUS Chronic Pain Treatment Guidelines.

MAXIMUS guideline: Decision based on MTUS Chronic Pain Treatment Guidelines Lyrica Page(s): 20.

Decision rationale: According to MTUS guidelines, Lyrica is an anti-epilepsy drug (AEDs - also referred to as anti-convulsant), which has been shown to be effective for treatment of diabetic; painful neuropathy and post-therapeutic neuralgia; and has been considered as a first-line treatment for neuropathic pain. There is no clear documentation of neuropathic pain in this patient that required and responded to previous use of Lyrica. In addition, there is no clear proven efficacy of Lyrica for back pain. Therefore, Lyrica 100 mg # 90 is not medically necessary.

Patches of Butrans 15 mcg, # 4: Upheld

Claims Administrator guideline: The Claims Administrator did not base their decision on the MTUS. Decision based on Non-MTUS Citation Official Disability Guidelines (ODG)

MAXIMUS guideline: Decision based on MTUS Chronic Pain Treatment Guidelines Criteria for use of opioids Page(s): 78-79.

Decision rationale: (a) Prescriptions from a single practitioner taken as directed, and all prescriptions from a single pharmacy.(b) The lowest possible dose should be prescribed to improve pain and function.(c) Office: Ongoing review and documentation of pain relief, functional status,appropriate medication use, and side effects. Pain assessment should include: currentpain; the least reported pain over the period since last assessment; average pain; intensity of pain after taking the opioid; how long it takes for pain relief; and how long pain relief lasts. Satisfactory response to treatment may be indicated by the patient's decreased pain, increased level of function, or improved quality of life. Information from family members or other caregivers should be considered in determining the patient's response to treatment. The 4 A's for Ongoing Monitoring: Four domains have been proposed as most relevant for ongoing monitoring of chronic pain patients on opioids: pain relief, side effects, physical and psychosocial functioning, and the occurrence of any potentially aberrant (or non-adherent) drug-related behaviors. These domains have been summarized as the "4 A's" (analgesia, activities of daily living, adverse side effects, and aberrant drug taking behaviors). The monitoring of these outcomes over time should affect therapeutic decisions and provide a framework.According to MTUS guidelines, Butrans is recommended to treat opiate addiction. There is no clear documentation of patient improvement in level of function, quality of life, adequate follow up or absence of side effects and aberrant behavior with previous use of opioids. The patient continued to have significant pain with Butrans. There is no justification to use multiple opioids. There is no recent documentation of recent opioid addiction. Therefore, the request for Patches of Butrans 15 mcg, # 4 is not medically necessary.