

Case Number:	CM14-0200944		
Date Assigned:	12/11/2014	Date of Injury:	02/17/2014
Decision Date:	05/15/2015	UR Denial Date:	11/18/2014
Priority:	Standard	Application Received:	12/01/2014

HOW THE IMR FINAL DETERMINATION WAS MADE

MAXIMUS Federal Services sent the complete case file to an expert reviewer. He/she has no affiliation with the employer, employee, providers or the claims administrator. He/she has been in active clinical practice for more than five years and is currently working at least 24 hours a week in active practice. The expert reviewer was selected based on his/her clinical experience, education, background, and expertise in the same or similar specialties that evaluate and/or treat the medical condition and disputed items/Service. He/she is familiar with governing laws and regulations, including the strength of evidence hierarchy that applies to Independent Medical Review determinations.

The Expert Reviewer has the following credentials:

State(s) of Licensure: North Carolina

Certification(s)/Specialty: Family Practice

CLINICAL CASE SUMMARY

The expert reviewer developed the following clinical case summary based on a review of the case file, including all medical records:

The injured worker is a 41 year old female with an industrial injury dated February 17, 2014. The injured worker diagnoses include cervical sprain/strain, thoracic sprain/strain, lateral epicondylitis, medial epicondylitis, elbow sprain/strain, rotator cuff syndrome and shoulder sprain/strain. He has been treated with diagnostic studies, prescribed medications and periodic follow up visits. According to the progress note dated 11/12/2014, the injured worker reported elbow pain, rated a 3/10 and a 1/10 with medication. The injured worker also reported that thoracic spine, cervical spine and shoulder/arm are feeling better. Objective findings revealed mild tenderness, spasm and decrease range of motion in the thoracic spine, lumbosacral spine, and cervical spine. The treating physician prescribed services for physical therapy and outpatient chiropractic treatment to cervical, thoracic, hand, left shoulder, left elbow consisting of manipulation, myofascial release, therapeutic exercise, e-stimulation and traction 2 times a week for 3 weeks.

IMR ISSUES, DECISIONS AND RATIONALES

The Final Determination was based on decisions for the disputed items/services set forth below:

Outpatient chiropractic treatment to cervical, thoracic, hand, left shoulder, left elbow consisting of manipulation, myofascial release, therapeutic exercise, e-stimulation and traction 2 times a week for 3 weeks: Upheld

Claims Administrator guideline: Decision based on MTUS ACOEM Chapter 2 General Approach to Initial Assessment and Documentation.

MAXIMUS guideline: Decision based on MTUS Chronic Pain Treatment Guidelines manual manipulation Page(s): 58-59.

Decision rationale: The California chronic pain medical guidelines section on manual manipulation states: Recommended for chronic pain if caused by musculoskeletal conditions. Manual Therapy is widely used in the treatment of musculoskeletal pain. The intended goal or effect of Manual Medicine is the achievement of positive symptomatic or objective measurable gains in functional improvement that facilitate progression in the patient's therapeutic exercise program and return to productive activities. Manipulation is manual therapy that moves a joint beyond the physiologic range-of-motion but not beyond the anatomic range-of-motion. Low back: Recommended as an option. Therapeutic care: Trial of 6 visits over 2 weeks, with evidence of objective functional improvement, total of up to 18 visits over 6-8 weeks. Elective/maintenance care: Not medically necessary. Recurrences/flare-ups: Need to reevaluate treatment success, if RTW achieved then 1-2 visits every 4-6 months. Ankle & Foot: Not recommended. Carpal tunnel syndrome: Not recommended. Forearm, Wrist, & Hand: Not recommended. Knee: Not recommended. Treatment Parameters from state guidelines: a. Time to produce effect: 4 to 6 treatments. Manual manipulation is recommended form of treatment for chronic pain. However, the requested amount of therapy sessions is in excess of the recommendations per the California MTUS. The California MTUS states there should be not more than 6 visits over 2 weeks and documented evidence of functional improvement before continuation of therapy. The request is for 3 weeks. This does not meet criteria guidelines and thus is not medically necessary.

Physical therapy to cervical, thoracic, hand, left shoulder, left elbow consisting of manipulation, myofascial release, therapeutic exercise, e-stimulation and traction 2 times a week for 3 weeks: Overturned

Claims Administrator guideline: Decision based on MTUS ACOEM Chapter 2 General Approach to Initial Assessment and Documentation.

MAXIMUS guideline: Decision based on MTUS Chronic Pain Treatment Guidelines physical medicine Page(s): 98-99.

Decision rationale: The California chronic pain medical treatment guidelines section on physical medicine states: Recommended as indicated below. Passive therapy (those treatment modalities that do not require energy expenditure on the part of the patient) can provide short-term relief during the early phases of pain treatment and are directed at controlling symptoms such as pain, inflammation and swelling and to improve the rate of healing soft tissue injuries. They can be used sparingly with active therapies to help control swelling, pain and inflammation during the rehabilitation process. Active therapy is based on the philosophy that therapeutic exercise and/or activity are beneficial for restoring flexibility, strength, endurance, function, range of motion, and can alleviate discomfort. Active therapy requires an internal effort by the

individual to complete a specific exercise or task. This form of therapy may require supervision from a therapist or medical provider such as verbal, visual and/or tactile instruction(s). Patients are instructed and expected to continue active therapies at home as an extension of the treatment process in order to maintain improvement levels. Home exercise can include exercise with or without mechanical assistance or resistance and functional activities with assistive devices. (Colorado, 2002) (Airaksinen, 2006) Patient-specific hand therapy is very important in reducing swelling, decreasing pain, and improving range of motion in CRPS. (Li, 2005) The use of active treatment modalities (e.g., exercise, education, activity modification) instead of passive treatments is associated with substantially better clinical outcomes. In a large case series of patients with low back pain treated by physical therapists, those adhering to guidelines for active rather than passive treatments incurred fewer treatment visits, cost less, and had less pain and less disability. The overall success rates were 64.7% among those adhering to the active treatment recommendations versus 36.5% for passive treatment. (Fritz, 2007) Physical Medicine Guidelines: Allow for fading of treatment frequency (from up to 3 visits per week to 1 or less), plus active self-directed home Physical Medicine. Myalgia and myositis, unspecified (ICD9 729.1): 9-10 visits over 8 weeks Neuralgia, neuritis, and radiculitis, unspecified (ICD9 729.2) 8-10 visits over 4 weeks Reflex sympathetic dystrophy (CRPS) (ICD9 337.2): 24 visits over 16 weeks. The goal of physical therapy is graduation to home therapy after a certain amount of recommended sessions. The request is not in excess of these recommendations per the California MTUS. Physical therapy is a recommended treatment .Therefore, the request is medically necessary.