

| | | | |
|-----------------------|--------------|------------------------------|------------|
| Case Number: | CM14-0202975 | | |
| Date Assigned: | 12/15/2014 | Date of Injury: | 12/17/2001 |
| Decision Date: | 02/06/2015 | UR Denial Date: | 11/07/2014 |
| Priority: | Standard | Application Received: | 12/04/2014 |

HOW THE IMR FINAL DETERMINATION WAS MADE

MAXIMUS Federal Services sent the complete case file to an expert reviewer. He/she has no affiliation with the employer, employee, providers or the claims administrator. The expert reviewer is Board Certified in Anesthesiology, has a subspecialty in Acupuncture and Pain Medicine and is licensed to practice in California. He/she has been in active clinical practice for more than five years and is currently working at least 24 hours a week in active practice. The expert reviewer was selected based on his/her clinical experience, education, background, and expertise in the same or similar specialties that evaluate and/or treat the medical condition and disputed items/services. He/she is familiar with governing laws and regulations, including the strength of evidence hierarchy that applies to Independent Medical Review determinations.

CLINICAL CASE SUMMARY

The expert reviewer developed the following clinical case summary based on a review of the case file, including all medical records:

61y/o female injured worker with date of injury 12/17/01 with related cervical and lumbar spine pain. Per progress report dated 10/20/14, the injured worker complained of residual cervical spine and lumbar spine pain. Pain in her lumbar spine was centered over bilateral SI joints. She also had left knee pain with swelling. Per physical exam, there was decreased range of motion secondary to pain and stiffness, Spurling's test was positive bilaterally. Examination of the lumbar spine revealed tenderness over the bilateral lumbar paraspinal musculature. Sensation was diminished to light touch and pinprick. DTRs were 1+ throughout. Treatment to date has included physical therapy and medication management. The date of UR decision was 11/7/14.

IMR ISSUES, DECISIONS AND RATIONALES

The Final Determination was based on decisions for the disputed items/services set forth below:

Fexmid 7.5mg #120: Upheld

Claims Administrator guideline: Decision based on MTUS Chronic Pain Treatment Guidelines Page(s): 64-66.

MAXIMUS guideline: Decision based on MTUS Chronic Pain Treatment Guidelines Muscle Relaxants Page(s): 64.

Decision rationale: With regard to muscle relaxants, the MTUS CPMTG states: "Recommend non-sedating muscle relaxants with caution as a second-line option for short-term treatment of

acute exacerbations in patients with chronic LBP. (Chou, 2007) (Mens, 2005) (Van Tulder, 1998) (van Tulder, 2003) (van Tulder, 2006) (Schnitzer, 2004) (See, 2008) Muscle relaxants may be effective in reducing pain and muscle tension, and increasing mobility. However, in most LBP cases, they show no benefit beyond NSAIDs in pain and overall improvement." Regarding Fexmid: "Recommended for a short course of therapy. Limited, mixed-evidence does not allow for a recommendation for chronic use. Cyclobenzaprine is a skeletal muscle relaxant and a central nervous system depressant with similar effects to tricyclic antidepressants (e.g. amitriptyline). Cyclobenzaprine is more effective than placebo in the management of back pain, although the effect is modest and comes at the price of adverse effects."The documentation submitted for review indicates that the injured worker was dispensed this medication on 9/12/14, as it is recommended for short-term use, the request is not medically necessary.

Prilosec 20mg #90: Upheld

Claims Administrator guideline: Decision based on MTUS Chronic Pain Treatment Guidelines NSAIDs, GI symptoms & cardiovascular risk.

MAXIMUS guideline: Decision based on MTUS Chronic Pain Treatment Guidelines NSAIDs, GI Symptoms & Cardiovascular Risk Page(s): 68.

Decision rationale: In the treatment of dyspepsia secondary to NSAID therapy, the MTUS recommends stopping the NSAID, switching to a different NSAID, or considering the use of an H2-receptor antagonist or a PPI. The MTUS Chronic Pain Medical Treatment Guidelines recommend the use of proton pump inhibitors in conjunction with NSAIDs in situations in which the patient is at risk for gastrointestinal events including: (1) age > 65 years; (2) history of peptic ulcer, GI bleeding or perforation; (3) concurrent use of ASA, corticosteroids, and/or an anticoagulant; or (4) high dose/multiple NSAID (e.g., NSAID + low-dose ASA). CPMTG guidelines further specify: "Recommendations: Patients with no risk factor and no cardiovascular disease: Non-selective NSAIDs OK (e.g. ibuprofen, naproxen, etc.) Patients at intermediate risk for gastrointestinal events and no cardiovascular disease: (1) A non-selective NSAID with either a PPI (Proton Pump Inhibitor, for example, 20 mg omeprazole daily) or misoprostol (200 g four times daily) or (2) a Cox-2 selective agent. Long-term PPI use (> 1 year) has been shown to increase the risk of hip fracture (adjusted odds ratio 1.44). Patients at high risk for gastrointestinal events with no cardiovascular disease: A Cox-2 selective agent plus a PPI if absolutely necessary. Patients at high risk of gastrointestinal events with cardiovascular disease: If GI risk is high the suggestion is for a low-dose Cox-2 plus low dose Aspirin (for cardioprotection) and a PPI. If cardiovascular risk is greater than GI risk the suggestion is naproxyn plus low-dose aspirin plus a PPI. (Laine, 2006) (Scholmerich, 2006) (Nielsen, 2006) (Chan, 2004) (Gold, 2007) (Laine, 2007)"While it is noted that the injured worker is being treated with fenoprofen, As there is no documentation of peptic ulcer, GI bleeding or perforation, or cardiovascular disease in the records available for my review, the injured worker's risk for gastrointestinal events is low, as such, medical necessity cannot be affirmed. It should be noted that the UR physician has certified a modification of the request for #30.

Flurbiprofen/Menthol/Camphor/Capsaicin compound cream 30gm: Upheld

Claims Administrator guideline: Decision based on MTUS Chronic Pain Treatment Guidelines Topical Analgesics Page(s): 111-113.

MAXIMUS guideline: Decision based on MTUS Chronic Pain Treatment Guidelines Topical Analgesics Page(s): 111-113.

Decision rationale: Capsaicin may have an indication for chronic lower back pain in this context. Per MTUS p 112 "Indications: There are positive randomized studies with capsaicin cream in patients with osteoarthritis, fibromyalgia, and chronic non-specific back pain, but it should be considered experimental in very high doses. Although topical capsaicin has moderate to poor efficacy, it may be particularly useful (alone or in conjunction with other modalities) in patients whose pain has not been controlled successfully with conventional therapy." Per MTUS with regard to Flurbiprofen (p112), "(Biswal, 2006) These medications may be useful for chronic musculoskeletal pain, but there are no long-term studies of their effectiveness or safety. Indications: Osteoarthritis and tendinitis, in particular, that of the knee and elbow or other joints that are amenable to topical treatment: Recommended for short-term use (4-12 weeks). There is little evidence to utilize topical NSAIDs for treatment of osteoarthritis of the spine, hip or shoulder." The documentation contains no evidence of osteoarthritis or tendinitis. Flurbiprofen is not indicated. Regarding the use of multiple medications, MTUS p60 states "Only one medication should be given at a time, and interventions that are active and passive should remain unchanged at the time of the medication change. A trial should be given for each individual medication. Analgesic medications should show effects within 1 to 3 days, and the analgesic effect of antidepressants should occur within 1 week. A record of pain and function with the medication should be recorded. (Mens, 2005) The recent AHRQ review of comparative effectiveness and safety of analgesics for osteoarthritis concluded that each of the analgesics was associated with a unique set of benefits and risks, and no currently available analgesic was identified as offering a clear overall advantage compared with the others." Therefore, it would be optimal to trial each medication individually. The CA MTUS, ODG, National Guidelines Clearinghouse, and ACOEM provide no evidence-based recommendations regarding the topical application of menthol or camphor. It is the opinion of this IMR reviewer that a lack of endorsement, a lack of mention, inherently implies a lack of recommendation, or a status equivalent to "not recommended". Since several components are not medically indicated, then the overall product is not indicated per MTUS as outlined below. Note the statement on page 111: Any compounded product that contains at least one drug (or drug class) that is not recommended is not recommended. The request is not medically necessary.

Flurbiprofen/Menthol/Camphor/Capsaicin compound cream 120gm: Upheld

Claims Administrator guideline: Decision based on MTUS Chronic Pain Treatment Guidelines Page(s): 111-113.

MAXIMUS guideline: Decision based on MTUS Chronic Pain Treatment Guidelines Topical Analgesics Page(s): 60; 111-113.

Decision rationale: Capsaicin may have an indication for chronic lower back pain in this context. Per MTUS p 112 "Indications: There are positive randomized studies with capsaicin

cream in patients with osteoarthritis, fibromyalgia, and chronic non-specific back pain, but it should be considered experimental in very high doses. Although topical capsaicin has moderate to poor efficacy, it may be particularly useful (alone or in conjunction with other modalities) in patients whose pain has not been controlled successfully with conventional therapy." Per MTUS with regard to Flurbiprofen (p112), "(Biswal, 2006) These medications may be useful for chronic musculoskeletal pain, but there are no long-term studies of their effectiveness or safety. Indications: Osteoarthritis and tendinitis, in particular, that of the knee and elbow or other joints that are amenable to topical treatment: Recommended for short-term use (4-12 weeks). There is little evidence to utilize topical NSAIDs for treatment of osteoarthritis of the spine, hip or shoulder." The documentation contains no evidence of osteoarthritis or tendinitis. Flurbiprofen is not indicated. Regarding the use of multiple medications, MTUS p60 states "Only one medication should be given at a time, and interventions that are active and passive should remain unchanged at the time of the medication change. A trial should be given for each individual medication. Analgesic medications should show effects within 1 to 3 days, and the analgesic effect of antidepressants should occur within 1 week. A record of pain and function with the medication should be recorded. (Mens, 2005) The recent AHRQ review of comparative effectiveness and safety of analgesics for osteoarthritis concluded that each of the analgesics was associated with a unique set of benefits and risks, and no currently available analgesic was identified as offering a clear overall advantage compared with the others." Therefore, it would be optimal to trial each medication individually. The CA MTUS, ODG, National Guidelines Clearinghouse, and ACOEM provide no evidence-based recommendations regarding the topical application of menthol or camphor. It is the opinion of this IMR reviewer that a lack of endorsement, a lack of mention, inherently implies a lack of recommendation, or a status equivalent to "not recommended". Since several components are not medically indicated, then the overall product is not indicated per MTUS as outlined below. Note the statement on page 111: Any compounded product that contains at least one drug (or drug class) that is not recommended is not recommended. The request is not medically necessary.