

Case Number:	CM14-0204713		
Date Assigned:	12/17/2014	Date of Injury:	09/03/2009
Decision Date:	03/05/2015	UR Denial Date:	11/07/2014
Priority:	Standard	Application Received:	12/08/2014

HOW THE IMR FINAL DETERMINATION WAS MADE

MAXIMUS Federal Services sent the complete case file to an expert reviewer. He/she has no affiliation with the employer, employee, providers or the claims administrator. He/she has been in active clinical practice for more than five years and is currently working at least 24 hours a week in active practice. The expert reviewer was selected based on his/her clinical experience, education, background, and expertise in the same or similar specialties that evaluate and/or treat the medical condition and disputed items/Service. He/she is familiar with governing laws and regulations, including the strength of evidence hierarchy that applies to Independent Medical Review determinations.

The Expert Reviewer has the following credentials:
 State(s) of Licensure: California, District of Columbia, Maryland
 Certification(s)/Specialty: Anesthesiology, Pain Management

CLINICAL CASE SUMMARY

The expert reviewer developed the following clinical case summary based on a review of the case file, including all medical records:

59y/o female injured worker with date of injury 9/3/09 with related neck, low back, and posterior lateral thigh pain. Per progress report dated 7/24/14, pain was rated 8/10 in intensity. Pain was described as neck spasms with constant dull achy pain in the low back and sharp shooting pain into the bilateral posterolateral lower extremities. Per physical exam, there was tenderness over the left side of the low back over the supragluteal region. There was tenderness to palpation of the bilateral trochanteric bursae. Tightness was noted about the bilateral iliotibial bands. Straight leg raise was positive bilaterally. The documentation submitted for review did not state whether physical therapy was utilized. Treatment to date has included medication management. The date of UR decision was 11/7/14.

IMR ISSUES, DECISIONS AND RATIONALES

The Final Determination was based on decisions for the disputed items/services set forth below:

Left Sided SI Joint Injection: Upheld

Claims Administrator guideline: The Claims Administrator did not cite any medical evidence for its decision.

MAXIMUS guideline: The Expert Reviewer did not base their decision on the MTUS. Decision based on Non-MTUS Citation Official Disability Guidelines (ODG) Hip & Pelvis, Sacroiliac Joint Injections.

Decision rationale: The MTUS is silent on the use of sacroiliac joint injections. Per ODG TWC with regard to sacroiliac joint injections: "Recommended as an option if failed at least 4-6 weeks of aggressive conservative therapy as indicated below." Criteria for the use of sacroiliac blocks: 1. The history and physical should suggest the diagnosis (with documentation of at least 3 positive exam findings as listed above). 2. Diagnostic evaluation must first address any other possible pain generators. 3. The patient has had and failed at least 4-6 weeks of aggressive conservative therapy including PT, home exercise and medication management. 4. Blocks are performed under fluoroscopy. (Hansen, 2003) 5. A positive diagnostic response is recorded as 80% for the duration of the local anesthetic. If the first block is not positive, a second diagnostic block is not performed. 6. If steroids are injected during the initial injection, the duration of pain relief should be at least 6 weeks with at least > 70% pain relief recorded for this period. 7. In the treatment or therapeutic phase (after the stabilization is completed), the suggested frequency for repeat blocks is 2 months or longer between each injection, provided that at least >70% pain relief is obtained for 6 weeks. 8. The block is not to be performed on the same day as a lumbar epidural steroid injection (ESI), transforaminal ESI, facet joint injection or medial branch block. 9. In the treatment or therapeutic phase, the interventional procedures should be repeated only as necessary judging by the medical necessity criteria, and these should be limited to a maximum of 4 times for local anesthetic and steroid blocks over a period of 1 year. The documentation submitted for review did not contain 3 positive exam findings (Cranial Shear Test; Extension Test; Flamingo Test; Fortin Finger Test; Gaenslen's Test; Gillet's Test (One Legged-Stork Test); Patrick's Test (FABER); Pelvic Compression Test; Pelvic Distraction Test; Pelvic Rock Test; Resisted Abduction Test (REAB); Sacroiliac Shear Test; Standing Flexion Test; Seated Flexion Test; Thigh Thrust Test (POSH).) suggesting the diagnosis of SI joint dysfunction. Furthermore, it was not indicated whether the injured worker has undergone conservative therapy in the form of physical therapy. As the criteria was not met, the request is not medically necessary.