

Case Number:	CM14-0207991		
Date Assigned:	12/19/2014	Date of Injury:	02/15/2012
Decision Date:	02/17/2015	UR Denial Date:	11/07/2014
Priority:	Standard	Application Received:	12/12/2014

HOW THE IMR FINAL DETERMINATION WAS MADE

MAXIMUS Federal Services sent the complete case file to an expert reviewer. He/she has no affiliation with the employer, employee, providers or the claims administrator. The expert reviewer is Board Certified in Emergency Medicine, and is licensed to practice in New York. He/she has been in active clinical practice for more than five years and is currently working at least 24 hours a week in active practice. The expert reviewer was selected based on his/her clinical experience, education, background, and expertise in the same or similar specialties that evaluate and/or treat the medical condition and disputed items/services. He/she is familiar with governing laws and regulations, including the strength of evidence hierarchy that applies to Independent Medical Review determinations.

CLINICAL CASE SUMMARY

The expert reviewer developed the following clinical case summary based on a review of the case file, including all medical records:

The patient is a 64-year-old male who was injured on February 15, 2012. The patient continued to experience pain in his lumbar spine with minor radiation in right leg. Physical examination was notable for muscle spasm, antalgic gait, and negative straight leg raise. Diagnoses included S/P decompression and posterolateral fusion and cervical disc herniations with bilateral radiculopathy. Treatment included medications, trigger point injections, physical therapy, aquatherapy, acupuncture, and surgery. Requests for authorization for Zanaflex 4 mg # 84, Norco 10/325 #126, and Colace 100 mg #80 were submitted for consideration.

IMR ISSUES, DECISIONS AND RATIONALES

The Final Determination was based on decisions for the disputed items/services set forth below:

Zanaflex 4mg Qty 84: Upheld

Claims Administrator guideline: Decision based on MTUS Chronic Pain Treatment Guidelines.

MAXIMUS guideline: Decision based on MTUS Chronic Pain Treatment Guidelines Pain Interventions and Guidelines Page(s): 63, 64.

Decision rationale: Zanaflex is the muscle relaxant, tizanidine. Tizanidine is a muscle relaxant that acts centrally as an alpha2-adrenergic agonist that is FDA approved for management of spasticity. Side effects include somnolence, dizziness, dry mouth, hypotension, weakness, and

hepatotoxicity. Non-sedating muscle relaxants are recommended with caution as a second-line option for short-term treatment (less than two weeks) of acute exacerbations in patients with chronic LBP. Muscle relaxants may be effective in reducing pain and muscle tension, and increasing mobility. However, in most LBP cases, they show no benefit beyond NSAIDs in pain and overall improvement. Also there is no additional benefit shown in combination with NSAIDs. Efficacy appears to diminish over time, and prolonged use of some medications in this class may lead to dependence. Sedation is the most commonly reported adverse effect of muscle relaxant medications. These drugs should be used with caution in patients driving motor vehicles or operating heavy machinery. In this case the patient has been using Zanaflex since at least April 2013. The duration of treatment surpasses the recommended short-term duration of two weeks. The request should not be authorized.

Norco 10/325mg Qty 126: Upheld

Claims Administrator guideline: Decision based on MTUS Chronic Pain Treatment Guidelines.

MAXIMUS guideline: Decision based on MTUS Chronic Pain Treatment Guidelines Pain Interventions and Guidelines Page(s): 11.

Decision rationale: Norco is the compounded medication containing hydrocodone and acetaminophen. Chronic Pain Medical Treatment Guidelines state that opioids are not recommended as a first line therapy. Opioid should be part of a treatment plan specific for the patient and should follow criteria for use. Criteria for use include establishment of a treatment plan, determination if pain is nociceptive or neuropathic, failure of pain relief with non-opioid analgesics, setting of specific functional goals, and opioid contract with agreement for random drug testing. If analgesia is not obtained, opioids should be discontinued. The patient should be screened for likelihood that he or she could be weaned from the opioids if there is no improvement in pain or function. It is recommended for short term use if first-line options, such as acetaminophen or NSAIDs have failed. Opioids may be a safer choice for patients with cardiac and renal disease than antidepressants or anticonvulsants. Acetaminophen is recommended for treatment of chronic pain & acute exacerbations of chronic pain. Acetaminophen overdose is a well-known cause of acute liver failure. Hepatotoxicity from therapeutic doses is unusual. Renal insufficiency occurs in 1 to 2% of patients with overdose. The recommended dose for mild to moderate pain is 650 to 1000 mg orally every 4 hours with a maximum of 4 g/day. In this case the patient has been using the Norco since at least April 2013 and has not obtained analgesia. In addition there is no documentation that the patient has signed an opioid contract. Criteria for long-term opioid use have not been met. The request should not be authorized.

Colace 100mg Qty 80: Upheld

Claims Administrator guideline: The Claims Administrator did not cite any medical evidence for its decision.

MAXIMUS guideline: The Expert Reviewer did not base their decision on the MTUS. Decision based on Non-MTUS Citation Official Disability Guidelines (ODG) Pain Opioid-induced constipation treatment Colace is docusate; Drugs for Irritable Bowel Syndrome Treatment Guidelines from The Medical Letter - July 1, 2011.

Decision rationale: Colace is docusate, a stool softener that works by increasing the amount of water that is absorbed by the stool in the gut. Opioid-induced constipation is a common adverse effect of long-term opioid use because the binding of opioids to peripheral opioid receptors in the gastrointestinal (GI) tract results in absorption of electrolytes, such as chloride, with a subsequent reduction in small intestinal fluid. Activation of enteric opioid receptors also results in abnormal GI motility. Constipation occurs commonly in patients receiving opioids and can be severe enough to cause discontinuation of therapy. If prescribing opioids has been determined to be appropriate, then ODG recommend that prophylactic treatment of constipation should be initiated. First-line: When prescribing an opioid, and especially if it will be needed for more than a few days, there should be an open discussion with the patient that this medication may be constipating, and the first steps should be identified to correct this. Simple treatments include increasing physical activity, maintaining appropriate hydration by drinking enough water, and advising the patient to follow a proper diet, rich in fiber. These can reduce the chance and severity of opioid-induced constipation and constipation in general. In addition, some laxatives may help to stimulate gastric motility. Other over-the-counter medications can help loosen otherwise hard stools, add bulk, and increase water content of the stool. In this case there is no documentation that the patient is suffering from constipation or that trial of alternative therapies such as increasing physical activity or maintaining hydration has failed. The request should not be authorized.