

Case Number:	CM14-0211019		
Date Assigned:	12/23/2014	Date of Injury:	03/20/2008
Decision Date:	02/13/2015	UR Denial Date:	12/09/2014
Priority:	Standard	Application Received:	12/16/2014

HOW THE IMR FINAL DETERMINATION WAS MADE

MAXIMUS Federal Services sent the complete case file to an expert reviewer. He/she has no affiliation with the employer, employee, providers or the claims administrator. The expert reviewer is Board Certified in Family Practice, and is licensed to practice in North Carolina. He/she has been in active clinical practice for more than five years and is currently working at least 24 hours a week in active practice. The expert reviewer was selected based on his/her clinical experience, education, background, and expertise in the same or similar specialties that evaluate and/or treat the medical condition and disputed items/services. He/she is familiar with governing laws and regulations, including the strength of evidence hierarchy that applies to Independent Medical Review determinations.

CLINICAL CASE SUMMARY

The expert reviewer developed the following clinical case summary based on a review of the case file, including all medical records:

The patient is a 56-year-old with a reported date of injury of 03/20/2008. The patient has the diagnoses of lumbar radiculitis, chronic lumbar sprain/strain, cervical myofascial pain syndrome and cervical radiculitis. Per the progress notes from the requesting physician dated 12/01/2014, the patient had complaints of persistent neck and back pain. The physical exam noted decreased range of motion in the lumbar and cervical spine. Treatment plan recommendations included increase in gabapentin, physical therapy and MRI of the lumbar spine.

IMR ISSUES, DECISIONS AND RATIONALES

The Final Determination was based on decisions for the disputed items/services set forth below:

6 Physical Therapy Sessions to the Low Back: Upheld

Claims Administrator guideline: Decision based on MTUS Chronic Pain Treatment Guidelines.

MAXIMUS guideline: Decision based on MTUS Chronic Pain Treatment Guidelines Physical medicine Page(s): 98-99.

Decision rationale: The California chronic pain medical treatment guidelines section on physical medicine states: Recommended as indicated below. Passive therapy (those treatment modalities that do not require energy expenditure on the part of the patient) can provide short term relief during the early phases of pain treatment and are directed at controlling symptoms

such as pain, inflammation and swelling and to improve the rate of healing soft tissue injuries. They can be used sparingly with active therapies to help control swelling, pain and inflammation during the rehabilitation process. Active therapy is based on the philosophy that therapeutic exercise and/or activity are beneficial for restoring flexibility, strength, endurance, function, range of motion, and can alleviate discomfort. Active therapy requires an internal effort by the individual to complete a specific exercise or task. This form of therapy may require supervision from a therapist or medical provider such as verbal, visual and/or tactile instruction(s). Patients are instructed and expected to continue active therapies at home as an extension of the treatment process in order to maintain improvement levels. Home exercise can include exercise with or without mechanical assistance or resistance and functional activities with assistive devices. (Colorado, 2002) (Airaksinen, 2006) Patient-specific hand therapy is very important in reducing swelling, decreasing pain, and improving range of motion in CRPS. (Li, 2005) The use of active treatment modalities (e.g., exercise, education, activity modification) instead of passive treatments is associated with substantially better clinical outcomes. In a large case series of patients with low back pain treated by physical therapists, those adhering to guidelines for active rather than passive treatments incurred fewer treatment visits, cost less, and had less pain and less disability. The overall success rates were 64.7% among those adhering to the active treatment recommendations versus 36.5% for passive treatment. (Fritz, 2007) Physical Medicine Guidelines - Allow for fading of treatment frequency (from up to 3 visits per week to 1 or less), plus active self-directed home Physical Medicine. Myalgia and myositis, unspecified (ICD9 729.1): 9-10 visits over 8 weeks Neuralgia, neuritis, and radiculitis, unspecified (ICD9 729.2) 8-10 visits over 4 weeks Reflex sympathetic dystrophy (CRPS) (ICD9 337.2): 24 visits over 16 weeks The California MTUS does recommend physical therapy as a treatment option for chronic pain. However this patient has chronic low back pain with an injury date of 03/20/2008. The progress notes indicate the back pain is worsening with constant pressure and a burning sensation radiating to the right lower extremity. However there is no indication of acute exacerbation of chronic low back pain which might warrant physical therapy. Therefore the request is not medically necessary.

Gabapentin 600mg (dosage/quantity unspecified): Overturned

Claims Administrator guideline: The Claims Administrator did not cite any medical evidence for its decision.

MAXIMUS guideline: Decision based on MTUS Chronic Pain Treatment Guidelines
Gabapentin Page(s): 18.

Decision rationale: The California chronic pain medical treatment guidelines section on Gabapentin states: Gabapentin (Neurontin, Gabarone, generic available) has been shown to be effective for treatment of diabetic painful neuropathy and postherpetic neuralgia and has been considered as a first-line treatment for neuropathic pain. (Backonja, 2002) (ICSI, 2007) (Knotkova, 2007) (Eisenberg, 2007) (Attal, 2006) This RCT concluded that gabapentin monotherapy appears to be efficacious for the treatment of pain and sleep interference associated with diabetic peripheral neuropathy and exhibits positive effects on mood and quality of life. (Backonja, 1998) It has been given FDA approval for treatment of post-herpetic neuralgia. The number needed to treat (NNT) for overall neuropathic pain is 4. It has a more favorable side-

effect profile than Carbamazepine, with a number needed to harm of 2.5. (Wiffen2-Cochrane, 2005) (Zaremba, 2006) Gabapentin in combination with morphine has been studied for treatment of diabetic neuropathy and postherpetic neuralgia. When used in combination the maximum tolerated dosage of both drugs was lower than when each was used as a single agent and better analgesia occurred at lower doses of each. (Gilron-NEJM, 2005) Recommendations involving combination therapy require further study. This medication is indicated for neuropathic pain as a first line agent. The patient has the diagnosis of cervical and lumbar radiculitis. The patient has subjective complaints consistent with neuropathic pain. Therefore the request is medically necessary.

Magnetic Resonance Imaging (MRI) of the Lumbar Spine: Upheld

Claims Administrator guideline: Decision based on MTUS ACOEM Chapter 12 Low Back Complaints Page(s): 303-304. Decision based on Non-MTUS Citation Official Disability Guidelines (ODG), Low Back-Lumbar & Thoracic

MAXIMUS guideline: Decision based on MTUS ACOEM Chapter 12 Low Back Complaints Page(s): 303-304.

Decision rationale: The ACOEM chapter on low back complaints and special diagnostic studies states: Unequivocal objective findings that identify specific nerve compromise on the neurologic examination are sufficient evidence to warrant imaging in patients who do not respond to treatment and who would consider surgery an option. When the neurologic examination is less clear, however, further physiologic evidence of nerve dysfunction should be obtained before ordering an imaging study. Indiscriminant imaging will result in false-positive findings, such as disk bulges, that are not the source of painful symptoms and do not warrant surgery. If physiologic evidence indicates tissue insult or nerve impairment, the practitioner can discuss with a consultant the selection of an imaging test to define a potential cause (magnetic resonance imaging [MRI] for neural or other soft tissue, computed tomography [CT] for bony structures). Relying solely on imaging studies to evaluate the source of low back and related symptoms carries a significant risk of diagnostic confusion (false positive test results) because of the possibility of identifying a finding that was present before symptoms began and therefore has no temporal association with the symptoms. Techniques vary in their abilities to define abnormalities (Table 12-7). Imaging studies should be reserved for cases in which surgery is considered or red-flag diagnoses are being evaluated. Because the overall false-positive rate is 30% for imaging studies in patients over age 30 who do not have symptoms, the risk of diagnostic confusion is great. There is no recorded physical exam that shows nerve tissue insult or impingement. There is no recorded presence of emerging red flags on the physical exam. The patient has subjective complaints of burning in the right lower extremity but no objective findings. For these reasons, criteria for imaging as defined above per the ACOEM have not been met. Therefore the request is not medically necessary.