

Case Number:	CM14-0039187		
Date Assigned:	06/27/2014	Date of Injury:	06/12/2009
Decision Date:	04/16/2015	UR Denial Date:	03/20/2014
Priority:	Standard	Application Received:	04/03/2014

HOW THE IMR FINAL DETERMINATION WAS MADE

MAXIMUS Federal Services sent the complete case file to an expert reviewer. He/she has no affiliation with the employer, employee, providers or the claims administrator. He/she has been in active clinical practice for more than five years and is currently working at least 24 hours a week in active practice. The expert reviewer was selected based on his/her clinical experience, education, background, and expertise in the same or similar specialties that evaluate and/or treat the medical condition and disputed items/Service. He/she is familiar with governing laws and regulations, including the strength of evidence hierarchy that applies to Independent Medical Review determinations.

The Expert Reviewer has the following credentials:
 State(s) of Licensure: New Jersey, Michigan, California
 Certification(s)/Specialty: Neurology, Neuromuscular Medicine

CLINICAL CASE SUMMARY

The expert reviewer developed the following clinical case summary based on a review of the case file, including all medical records:

The injured worker is a 54 year old male who sustained an industrial injury to his left shoulder, neck and right wrist on June 12, 2009. The injured worker is status post left shoulder surgery in January 2013 and July 2013 and left wrist arthroscopy in February 2011. The injured worker was diagnosed with cervical sprain/strain, left shoulder degenerative joint disease and complex regional pain syndrome. According to the primary treating physician's progress report on December 9, 2013, the injured worker continues to experience stiffness of the left shoulder with pain against resistance. Examination noted range of motion at 145 degrees in both flexion and abduction. Current medications consist of Percocet, Norco, Naproxen, Omeprazole and Xanax. Treatment modalities consist of continuation of conservative therapy, a trial of Vimovo and prescribed medications for break through pain.

IMR ISSUES, DECISIONS AND RATIONALES

The Final Determination was based on decisions for the disputed items/services set forth below:

Percocet 10/325mg #60 3 refills: Upheld

Claims Administrator guideline: Decision based on MTUS Chronic Pain Treatment Guidelines Opioids.

MAXIMUS guideline: Decision based on MTUS Chronic Pain Treatment Guidelines Criteria for use of opioids Page(s): 179.

Decision rationale: According to MTUS guidelines, ongoing use of opioids should follow specific rules: “(a) Prescriptions from a single practitioner taken as directed, and all prescriptions from a single pharmacy. (b) The lowest possible dose should be prescribed to improve pain and function. (c) Office: Ongoing review and documentation of pain relief, functional status, appropriate medication use, and side effects. Pain assessment should include current pain; the least reported pain over the period since last assessment; average pain; intensity of pain after taking the opioid; how long it takes for pain relief; and how long pain relief lasts. Satisfactory response to treatment may be indicated by the patient's decreased pain, increased level of function, or improved quality of life. Information from family members or other caregivers should be considered in determining the patient's response to treatment. The 4 A's for Ongoing Monitoring: Four domains have been proposed as most relevant for ongoing monitoring of chronic pain patients on opioids: pain relief, side effects, physical and psychosocial functioning, and the occurrence of any potentially aberrant (or non adherent) drug-related behaviors. These domains have been summarized as the "4 A's" (analgesia, activities of daily living, adverse side effects, and aberrant drug taking behaviors). The monitoring of these outcomes over time should affect therapeutic decisions and provide a framework.” The patient have been using opioids for long period of time without recent documentation of full control of pain and without any documentation of functional or quality of life improvement. There is no clear documentation of patient improvement in level of function, quality of life, adequate follow up for absence of side effects and aberrant behavior with a previous use of narcotics. There is no justification for the use of several narcotics. Therefore, the prescription of Percocet 10/325mg, #60, with 3 refills is not medically necessary.

Norco 10/325mg #60 3 refills: Upheld

Claims Administrator guideline: Decision based on MTUS Chronic Pain Treatment Guidelines Opioids.

MAXIMUS guideline: Decision based on MTUS Chronic Pain Treatment Guidelines Criteria for use of opioids Page(s): 76-79.

Decision rationale: According to MTUS guidelines, Norco (Hydrocodone/Acetaminophen) is a synthetic opioid indicated for the pain management but not recommended as a first line oral analgesic. In addition and according to MTUS guidelines, ongoing use of opioids should follow specific rules: “(a) Prescriptions from a single practitioner taken as directed, and all prescriptions from a single pharmacy. (b) The lowest possible dose should be prescribed to improve pain and function. (c) Office: Ongoing review and documentation of pain relief, functional status, appropriate medication use, and side effects. Four domains have been proposed as most relevant for ongoing monitoring of chronic pain patients on opioids: pain relief, side effects, physical and psychosocial functioning, and the occurrence of any potentially aberrant (or non adherent) drug-related behaviors. These domains have been summarized as the "4 A's" (analgesia, activities of daily living, adverse side effects, and aberrant drug taking behaviors). The monitoring of these outcomes over time should affect therapeutic decisions and provide a framework.” According to

the patient file, there is no objective documentation of pain and functional improvement to justify continuous use of Norco. Norco was used for longtime without documentation of functional improvement or evidence of return to work or improvement of activity of daily living. Therefore, the prescription of Norco 10/325mg #60, with 3 refills is not medically necessary.

Trial Vimovo 375/20mg #60: Upheld

Claims Administrator guideline: The Claims Administrator did not base their decision on the MTUS. Decision based on Non-MTUS Citation Official Disability Guidelines.

MAXIMUS guideline: Decision based on MTUS Chronic Pain Treatment Guidelines NSAIDs, GI symptoms & cardiovascular risk Page(s): 68.

Decision rationale: Vimovo is formed by esomeprazole and naproxen. According to MTUS guidelines, Omeprazole is indicated when NSAID are used in patients with intermediate or high risk for gastrointestinal events. The risk for gastrointestinal events are: (1) age > 65 years; (2) history of peptic ulcer, GI bleeding or perforation; (3) concurrent use of ASA, corticosteroids, and/or an anticoagulant; or (4) high dose/multiple NSAID (e.g., NSAID + low-dose ASA). Recent studies tend to show that H. Pylori does not act synergistically with NSAIDS to develop gastroduodenal lesions. There is no documentation that the patient has GI issue that requires the use of nexium. There is no documentation in the patient's chart supporting that he is at intermediate or high risk for developing gastrointestinal events. In addition, there is no controlled studies supporting the superiority of the use of Vimovo to Naproxen and Omeprazol used separately. Therefore, trial Vimovo 375mg/20mg #60 prescription is not medically necessary.