

Case Number:	CM14-0064860		
Date Assigned:	07/11/2014	Date of Injury:	02/27/2014
Decision Date:	04/16/2015	UR Denial Date:	04/16/2014
Priority:	Standard	Application Received:	05/07/2014

HOW THE IMR FINAL DETERMINATION WAS MADE

MAXIMUS Federal Services sent the complete case file to an expert reviewer. He/she has no affiliation with the employer, employee, providers or the claims administrator. He/she has been in active clinical practice for more than five years and is currently working at least 24 hours a week in active practice. The expert reviewer was selected based on his/her clinical experience, education, background, and expertise in the same or similar specialties that evaluate and/or treat the medical condition and disputed items/Service. He/she is familiar with governing laws and regulations, including the strength of evidence hierarchy that applies to Independent Medical Review determinations.

The Expert Reviewer has the following credentials:
 State(s) of Licensure: New Jersey, Michigan, California
 Certification(s)/Specialty: Neurology, Neuromuscular Medicine

CLINICAL CASE SUMMARY

The expert reviewer developed the following clinical case summary based on a review of the case file, including all medical records:

The injured worker is a 51-year-old female, who sustained an industrial injury on February 27, 2014. She reported cumulative trauma injury to the neck, back, upper extremities and lower extremities. The injured worker was diagnosed as having lumbar radiculopathy, lumbar sprain/strain, right shoulder sprain/strain, left shoulder sprain/strain, right knee sprain/strain and right foot sprain/strain. Treatment to date has included diagnostic studies, physical therapy, acupuncture and medications. On January 25, 2015, the injured worker complained of intermittent neck pain, constant back pain, intermittent right shoulder pain, constant left wrist pain, constant right knee pain and constant right foot pain. Physical examination of the right shoulder and lumbar spine revealed tenderness and loss of motion. Tenderness was also noted in the left wrist. The treatment plan included more diagnostic studies, physical therapy, acupuncture, possible intra-articular injections to the right shoulder, possible surgery, left wrist splinting, possible intra-articular injections to the left wrist, possible intra-articular injection/cortisone injection to the right foot, right knee brace and medications.

IMR ISSUES, DECISIONS AND RATIONALES

The Final Determination was based on decisions for the disputed items/services set forth below:

MRI of the right shoulder: Upheld

Claims Administrator guideline: Decision based on MTUS ACOEM Chapter 9 Shoulder Complaints.

MAXIMUS guideline: Decision based on MTUS ACOEM Chapter 9 Shoulder Complaints Page(s): 209.

Decision rationale: According to MTUS guidelines, MRI of the shoulder is indicated in case of tumor, infection, ligament instability and rotator cuff injury. There is no clinical evidence or documentation of one of the above diagnosis. Therefore, MRI of the right shoulder is not medically necessary.

MRI of the left wrist: Upheld

Claims Administrator guideline: Decision based on MTUS ACOEM Chapter 11 Forearm, Wrist, and Hand Complaints. Decision based on Non-MTUS Citation Official Disability Guidelines (ODG)- Forearm, Wrist & Hand MRI's.

MAXIMUS guideline: The Expert Reviewer did not base their decision on the MTUS. Decision based on Non-MTUS Citation <http://www.odg-twc.com/index.html>.

Decision rationale: According to ODG guidelines, MRI of the wrist "Recommended as indicated below. While criteria for which patients may benefit from the addition of MRI have not been established, in selected cases where there is a high clinical suspicion of a fracture despite normal radiographs, MRI may prove useful. (ACR, 2001) (Schmitt, 2003) (Valeri, 1999) (Duer, 2007) Magnetic resonance imaging has been advocated for patients with chronic wrist pain because it enables clinicians to perform a global examination of the osseous and soft tissue structures. It may be diagnostic in patients with triangular fibrocartilage (TFC) and intraosseous ligament tears, occult fractures, avascular neurosis, and miscellaneous other abnormalities. Many articles dispute the value of imaging in the diagnosis of ligamentous tears, because arthroscopy may be more accurate and treatment can be performed along with the diagnosis. (Dalinka, 2000) (Tehranzadeh, 2006) For inflammatory arthritis, high-resolution in-office MRI with an average follow-up of 8 months detects changes in bony disease better than radiography, which is insensitive for detecting changes in bone erosions for this patient population in this time frame. (Chen, 2006) See also Radiography. Indications for imaging Magnetic resonance imaging (MRI): Acute hand or wrist trauma, suspect acute distal radius fracture, radiographs normal, next procedure if immediate confirmation or exclusion of fracture is required; Acute hand or wrist trauma, suspect acute scaphoid fracture, radiographs normal, next procedure if immediate confirmation or exclusion of fracture is required; Acute hand or wrist trauma, suspect gamekeeper injury (thumb MCP ulnar collateral ligament injury); Chronic wrist pain, plain films normal, suspect soft tissue tumor; Chronic wrist pain, plain film normal or equivocal, suspect Kienbock's disease; Repeat MRI is not routinely recommended, and should be reserved for a significant change in symptoms and/or findings suggestive of significant pathology. (Mays, 2008)." There is no documentation that the patient is suspected of wrist fracture. There is no indication of Wrist MRI as per ODG criteria. Therefore, the request for left wrist MRI is not medically necessary.

MRI of the lumbar spine: Upheld

Claims Administrator guideline: Decision based on MTUS ACOEM Chapter 12 Low Back Complaints. Decision based on Non-MTUS Citation Official Disability Guidelines (ODG)- Low Back MRIs.

MAXIMUS guideline: Decision based on MTUS ACOEM Chapter 12 Low Back Complaints Page(s): 303.

Decision rationale: Regarding the indications for imaging in case of back pain, MTUS guidelines stated: "Lumbar spine x rays should not be recommended in patients with low back pain in the absence of red flags for serious spinal pathology, even if the pain has persisted for at least six weeks. However, it may be appropriate when the physician believes it would aid in patient management. Unequivocal objective findings that identify specific nerve compromise on the neurologic examination are sufficient evidence to warrant imaging in patients who do not respond to treatment and who would consider surgery an option. When the neurologic examination is less clear, however, further physiologic evidence of nerve dysfunction should be obtained before ordering an imaging study. Indiscriminant imaging will result in false-positive findings, such as disk bulges, that are not the source of painful symptoms and do not warrant surgery. If physiologic evidence indicates tissue insult or nerve impairment, the practitioner can discuss with a consultant the selection of an imaging test to define a potential cause (magnetic resonance imaging [MRI] for neural or other soft tissue, computer tomography [CT] for bony structures)". Furthermore, and according to MTUS guidelines, MRI is the test of choice for patients with prior back surgery, fracture or tumors that may require surgery. The patient does not have any clear evidence of new lumbar nerve root compromise. There is no clear evidence of significant change in the patient signs or symptoms suggestive of new pathology. Therefore, the request for lumbar MRI is not medically necessary.