

<b>Case Number:</b>	CM14-0085859		
<b>Date Assigned:</b>	07/23/2014	<b>Date of Injury:</b>	08/27/2007
<b>Decision Date:</b>	04/07/2015	<b>UR Denial Date:</b>	05/20/2014
<b>Priority:</b>	Standard	<b>Application Received:</b>	06/09/2014

### HOW THE IMR FINAL DETERMINATION WAS MADE

MAXIMUS Federal Services sent the complete case file to an expert reviewer. He/she has no affiliation with the employer, employee, providers or the claims administrator. He/she has been in active clinical practice for more than five years and is currently working at least 24 hours a week in active practice. The expert reviewer was selected based on his/her clinical experience, education, background, and expertise in the same or similar specialties that evaluate and/or treat the medical condition and disputed items/Service. He/she is familiar with governing laws and regulations, including the strength of evidence hierarchy that applies to Independent Medical Review determinations.

The Expert Reviewer has the following credentials:  
 State(s) of Licensure: New Jersey, Michigan, California  
 Certification(s)/Specialty: Neurology, Neuromuscular Medicine

### CLINICAL CASE SUMMARY

The expert reviewer developed the following clinical case summary based on a review of the case file, including all medical records:

The injured worker is a 53-year-old male, who sustained an industrial injury on 8/27/07. On 6/9/14, the injured worker submitted an application for IMR for review of Tramadol 50mg #60. The treating provider has reported the injured worker complained of low back pain radiating down right leg. The diagnoses have included cervical, thoracic and lumbar multilevel disc protrusions. Treatment to date has included status post L5-S1 laminectomy (prior to injury 1987), physical therapy, aquatic therapy, chiropractic care, medications. On 5/20/14 Utilization Review non-certified Tramadol 50mg #60 The MTUS Guidelines were cited.

### IMR ISSUES, DECISIONS AND RATIONALES

The Final Determination was based on decisions for the disputed items/services set forth below:

**Tramadol 50mg #60:** Upheld

**Claims Administrator guideline:** Decision based on MTUS Chronic Pain Treatment Guidelines Page(s): 76.

**MAXIMUS guideline:** Decision based on MTUS Chronic Pain Treatment Guidelines Tramadol Page(s): 113.

**Decision rationale:** According to MTUS guidelines, Ultram is a synthetic opioid indicated for the pain management but not recommended as a first line oral analgesic. Although, Ultram may be needed to help with the patient pain, it may not help with the weaning process from opioids. Ultram could be used if exacerbation of pain after or during the weaning process. In addition and according to MTUS guidelines, ongoing use of opioids should follow specific rules: (a) Prescriptions from a single practitioner taken as directed, and all prescriptions from a single pharmacy. (b) The lowest possible dose should be prescribed to improve pain and function. (c) Office: Ongoing review and documentation of pain relief, functional status, appropriate medication use, and side effects. Pain assessment should include: current pain; the least reported pain over the period since last assessment; average pain; intensity of pain after taking the opioid; how long it takes for pain relief; and how long pain relief lasts. Satisfactory response to treatment may be indicated by the patient's decreased pain, increased level of function, or improved quality of life. Information from family members or other caregivers should be considered in determining the patient's response to treatment. The 4 A's for Ongoing Monitoring: Four domains have been proposed as most relevant for ongoing monitoring of chronic pain patients on opioids: pain relief, side effects, physical and psychosocial functioning, and the occurrence of any potentially aberrant (or non adherent) drug-related behaviors. These domains have been summarized as the "4 A's" (analgesia, activities of daily living, adverse side effects, and aberrant drug taking behaviors). The monitoring of these outcomes over time should affect therapeutic decisions and provide a framework. There is no clear recent and objective documentation of pain and functional improvement in this patient with previous use of Tramadol. There is no clear documentation of compliance and UDS for previous use of tramadol. Therefore, the prescription of Tramadol 50mg #60 is not medically necessary.