

Case Number:	CM15-0000326		
Date Assigned:	01/09/2015	Date of Injury:	10/15/2004
Decision Date:	09/21/2015	UR Denial Date:	12/15/2014
Priority:	Standard	Application Received:	01/02/2015

HOW THE IMR FINAL DETERMINATION WAS MADE

MAXIMUS Federal Services sent the complete case file to an expert reviewer. He/she has no affiliation with the employer, employee, providers or the claims administrator. He/she has been in active clinical practice for more than five years and is currently working at least 24 hours a week in active practice. The expert reviewer was selected based on his/her clinical experience, education, background, and expertise in the same or similar specialties that evaluate and/or treat the medical condition and disputed items/Service. He/she is familiar with governing laws and regulations, including the strength of evidence hierarchy that applies to Independent Medical Review determinations.

The Expert Reviewer has the following credentials:

State(s) of Licensure: Iowa, Illinois, Hawaii

Certification(s)/Specialty: Preventive Medicine, Occupational Medicine, Public Health & General Preventive Medicine

CLINICAL CASE SUMMARY

The expert reviewer developed the following clinical case summary based on a review of the case file, including all medical records:

This 42-year-old male sustained an industrial injury on 10/15/04. He subsequently reported neck and shoulder pain. Diagnoses include cervical pain, cervical radiculopathy, cervical disc displacement and degeneration of the cervical spine. Treatments to date include prescription pain medications. The injured worker continues to have limited use of the left shoulder and hand. Upon examination, the provider requested modified use of left hand with no repetitive power grasping. A request for Elavil, Ambien, Norco, Naproxen and Protonix medications was made by the treating physician.

IMR ISSUES, DECISIONS AND RATIONALES

The Final Determination was based on decisions for the disputed items/services set forth below:

Elavil 75mg #60: Upheld

Claims Administrator guideline: Decision based on MTUS Chronic Pain Treatment Guidelines.

MAXIMUS guideline: Decision based on MTUS Chronic Pain Treatment Guidelines Chronic pain Page(s): 13. Decision based on Non-MTUS Citation Official Disability Guidelines (ODG) Chronic Pain, TCAs.

Decision rationale: MTUS states that "Amitriptyline is a tricyclic antidepressant. Tricyclics are generally considered a first-line agent unless they are ineffective, poorly tolerated, or contraindicated." ODG states "Assessment of treatment efficacy should include not only pain outcomes, but also an evaluation of function, changes in use of other analgesic medication, sleep quality and duration, and psychological assessment. Side effects, including excessive sedation (especially that which would affect work performance) should be assessed. (Additional side effects are listed below for each specific drug.) It is recommended that these outcome measurements should be initiated at one week of treatment with a recommended trial of at least 4 weeks. The optimal duration of treatment is not known because most double-blind trials have been of short duration (6-12 weeks). It has been suggested that if pain is in remission for 3-6 months, a gradual tapering of anti-depressants may be undertaken." ODG states "Dosing Information: Amitriptyline: Neuropathic pain: The starting dose may be as low as 10-25 mg at night, with increases of 10-25 mg once or twice a week up to 100 mg/day. (ICSI, 2007) The lowest effective dose should be used (Dworkin, 2007)." The treating physician has not provided evidence of improved pain control, improved function or sleep quality from Elavil to warrant ongoing therapy. As such, the request for Elavil 75mg #60 was not medically necessary.

Ambien 1m #30: Upheld

Claims Administrator guideline: The Claims Administrator did not base their decision on the MTUS. Decision based on Non-MTUS Citation Official Disability Guidelines (ODG), Pain.

MAXIMUS guideline: The Expert Reviewer did not base their decision on the MTUS. Decision based on Non-MTUS Citation Official Disability Guidelines (ODG) Pain, Zolpidem, insomnia treatment.

Decision rationale: The CA MTUS silent regarding this topic. ODG states that zolpidem is a prescription short acting non-benzodiazepine hypnotic, which is approved for short-term treatment of insomnia. There has been no discussion of the patient's sleep hygiene or the need for variance from the guidelines, such as "(a) Wake at the same time everyday; (b) Maintain a consistent bedtime; (c) Exercise regularly (not within 2 to 4 hours of bedtime); (d) Perform relaxing activities before bedtime; (e) Keep your bedroom quiet and cool; (f) Do not watch the clock; (g) Avoid caffeine and nicotine for at least six hours before bed; (h) Only drink in moderation; & (i) Avoid napping. Medical documents also do not include results of these first line treatments, if they were used in treatment of the patient's insomnia. ODG additionally states "The specific component of insomnia should be addressed: (a) Sleep onset; (b) Sleep maintenance; (c) Sleep quality; & (d) Next-day functioning." Medical documents provided do not detail these components. As such, the request for Ambien 1m #30 is not medically necessary at this time.

Norco 10/325mg #60: Upheld

Claims Administrator guideline: Decision based on MTUS Chronic Pain Treatment Guidelines opioids.

MAXIMUS guideline: Decision based on MTUS Chronic Pain Treatment Guidelines Opioids Page(s): 74-96. Decision based on Non-MTUS Citation Official Disability Guidelines (ODG) Neck and Upper Back (Acute and Chronic), Low Back - Lumbar & Thoracic (Acute & Chronic), Shoulder, Pain, Opioids.

Decision rationale: ODG does not recommend the use of opioids for neck and shoulder pain "except for short use for severe cases, not to exceed 2 weeks." The patient has exceeded the 2 week recommended treatment length for opioid usage. MTUS does not discourage use of opioids past 2 weeks, but does state that "ongoing review and documentation of pain relief, functional status, appropriate medication use, and side effects. Pain assessment should include: current pain; the least reported pain over the period since last assessment; average pain; intensity of pain after taking the opioid; how long it takes for pain relief; and how long pain relief lasts. Satisfactory response to treatment may be indicated by the patient's decreased pain, increased level of function, or improved quality of life." The treating physician does not fully document the least reported pain over the period since last assessment, intensity of pain after taking opioid, pain relief, increased level of function, or improved quality of life. As such, the request for Norco 10/325mg #60 is not medically necessary.

Naproxen 375mg #90: Upheld

Claims Administrator guideline: Decision based on MTUS Chronic Pain Treatment Guidelines anti-inflammatory drug.

MAXIMUS guideline: Decision based on MTUS Chronic Pain Treatment Guidelines NSAIDs (non-steroidal anti-inflammatory drugs) Page(s): 67-73. Decision based on Non-MTUS Citation Official Disability Guidelines (ODG) Pain (Chronic), Naproxen, NSAIDs (non-steroidal anti-inflammatory drugs).

Decision rationale: MTUS specifies four recommendations regarding NSAID use: 1) Osteoarthritis (including knee and hip): Recommended at the lowest dose for the shortest period in patients with moderate to severe pain. 2) Back Pain - Acute exacerbations of chronic pain: Recommended as a second-line treatment after acetaminophen. In general, there is conflicting evidence that NSAIDs are more effective than acetaminophen for acute LBP. 3) Back Pain - Chronic low back pain: Recommended as an option for short-term symptomatic relief. A Cochrane review of the literature on drug relief for low back pain (LBP) suggested that NSAIDs were no more effective than other drugs such as acetaminophen, narcotic analgesics, and muscle relaxants. The review also found that NSAIDs had more adverse effects than placebo and acetaminophen but fewer effects than muscle relaxants and narcotic analgesics. 4) Neuropathic pain: There is inconsistent evidence for the use of these medications to treat long-term neuropathic pain, but they may be useful to treat breakthrough and mixed pain conditions such as osteoarthritis (and other nociceptive pain) in with neuropathic pain. The medical documents do not indicate that the patient is being treated for osteoarthritis. Additionally, the treating physician does not document failure of primary (Tylenol) treatment. Progress notes do not indicate how long the patient has been on naproxen, but the MTUS guidelines recommend against long-term use. Additionally, the medical documentation provided does not indicate objective functional improvement with the use of Naproxen. As such, the request for Naproxen 375mg #90 is not medically necessary.

Protonix 20mg #60: Upheld

Claims Administrator guideline: Decision based on MTUS Chronic Pain Treatment Guidelines proton pump inhibitors.

MAXIMUS guideline: Decision based on MTUS Chronic Pain Treatment Guidelines NSAIDs, GI symptoms & cardiovascular risk Page(s): 68-69. Decision based on Non-MTUS Citation Official Disability Guidelines (ODG) Pain (Chronic), NSAIDs, GI symptoms & cardiovascular risk.

Decision rationale: Protonix is the brand name version of Pantoprazole, which is a proton pump inhibitor. MTUS states, "Determine if the patient is at risk for gastrointestinal events: (1) age > 65 years; (2) history of peptic ulcer, GI bleeding or perforation; (3) concurrent use of ASA, corticosteroids, and/or an anticoagulant; or (4) high dose/multiple NSAID (e. g. , NSAID + low-dose ASA)." And "Patients at intermediate risk for gastrointestinal events and no cardiovascular disease: (1) A non-selective NSAID with either a PPI (Proton Pump Inhibitor, for example, 20 mg omeprazole daily) or misoprostol (200g four times daily) or (2) a Cox-2 selective agent. Long-term PPI use (> 1 year) has been shown to increase the risk of hip fracture (adjusted odds ratio 1. 44)." ODG states, "If a PPI is used, omeprazole OTC tablets or lansoprazole 24HR OTC are recommended for an equivalent clinical efficacy and significant cost savings. Products in this drug class have demonstrated equivalent clinical efficacy and safety at comparable doses, including esomeprazole (Nexium), lansoprazole (Prevacid), omeprazole (Prilosec), pantoprazole (Protonix), dexlansoprazole (Dexilant), and rabeprazole (Aciphex). (Shi, 2008) A trial of omeprazole or lansoprazole is recommended before Nexium therapy. The other PPIs, Protonix, Dexilant, and Aciphex, should also be second-line. According to the latest AHRQ Comparative Effectiveness Research, all of the commercially available PPIs appeared to be similarly effective. (AHRQ, 2011)." The patient does not meet the age recommendations for increased GI risk. The medical documents provided do not indicate history of peptic ulcer, GI bleeding or perforation. Medical records do not indicate that the patient is on ASA, corticosteroids, and/or an anticoagulant; or high dose/multiple NSAID. Additionally per guidelines, Pantoprazole is considered second line therapy and the treating physician has not provided detailed documentation of a failed trial of omeprazole and/or lansoprazole. As such, the request for Protonix 20mg #60 is not medically necessary.