

Case Number:	CM15-0001064		
Date Assigned:	01/12/2015	Date of Injury:	09/16/2010
Decision Date:	04/08/2015	UR Denial Date:	12/15/2014
Priority:	Standard	Application Received:	01/05/2015

HOW THE IMR FINAL DETERMINATION WAS MADE

MAXIMUS Federal Services sent the complete case file to an expert reviewer. He/she has no affiliation with the employer, employee, providers or the claims administrator. He/she has been in active clinical practice for more than five years and is currently working at least 24 hours a week in active practice. The expert reviewer was selected based on his/her clinical experience, education, background, and expertise in the same or similar specialties that evaluate and/or treat the medical condition and disputed items/Service. He/she is familiar with governing laws and regulations, including the strength of evidence hierarchy that applies to Independent Medical Review determinations.

The Expert Reviewer has the following credentials:
 State(s) of Licensure: North Carolina
 Certification(s)/Specialty: Family Practice

CLINICAL CASE SUMMARY

The expert reviewer developed the following clinical case summary based on a review of the case file, including all medical records:

The injured worker is a 57 year old female with an industrial injury dated 09/16/2010 a motor vehicle accident resulting in injury to neck back and ribs. Diagnoses include lumbar discopathy. Diagnostic testing has included MRI of the cervical spine (11/12/2012), x-ray of the lumbar spine (09/05/2013), and electro diagnostic studies. Previous treatments have included conservative measures, medications, physical therapy, surgeries, and injections. A progress note dated 09/05/2013, reports occasional pain in the low back radiating to the buttocks and to the legs and associated with numbness and tingling. The objective examination revealed pain and tenderness right across the iliac crest and into the lumbosacral spine, guarded and restricted range of motion, and dysesthesia in the L4-S1 dermatomes. The treating physician is requesting retrospective ondansetron and tramadol which were denied by the utilization review. On 12/15/2014, Utilization Review non-certified retrospective prescriptions for ondansetron tablets 8mg #30 times 2 (date of service 09/05/2013), and tramadol hydrochloride ER 150mg #90 (date of service 09/05/2013, noting the MTUS guidelines were cited. On 01/05/2015, the injured worker submitted an application for IMR for review of ondansetron tablets 8mg #30 times 2 (date of service 09/05/2013), and tramadol hydrochloride ER 150mg #90 (date of service 09/05/2013).

IMR ISSUES, DECISIONS AND RATIONALES

The Final Determination was based on decisions for the disputed items/services set forth below:

Ondansetron Tablets 8mg #30x 2, (DOS 09/05/2013): Upheld

Claims Administrator guideline: The Claims Administrator did not base their decision on the MTUS. Decision based on Non-MTUS Citation Official Disability Guidelines (ODG) Pain, Antiemetics.

MAXIMUS guideline: The Expert Reviewer did not base their decision on the MTUS. Decision based on Non-MTUS Citation ODG, zofran.

Decision rationale: The California MTUS and the ACOEM do not specifically address the requested medication. Per the Official Disability Guidelines section on Ondanset, the medication is indicated for the treatment of nausea and vomiting associated with chemotherapy, radiation therapy or post-operatively. The medication is not indicated for the treatment of nausea and vomiting associated with chronic opioid use. The patient does not have a malignancy diagnosis. There is also no indication that the patient has failed more traditional first line medication such as promethazine or Compazine. For these reasons the request is not certified.

Tramadol Hydrochloride ER 150mg #90 (DOS 09/05/2013): Upheld

Claims Administrator guideline: Decision based on MTUS Chronic Pain Treatment Guidelines Opioids Page(s): 93-94.

MAXIMUS guideline: Decision based on MTUS Chronic Pain Treatment Guidelines opioids Page(s): 76-84.

Decision rationale: The California chronic pain medical treatment guidelines section on opioids states for ongoing management: On-Going Management. Actions Should Include: (a) Prescriptions from a single practitioner taken as directed, and all prescriptions from a single pharmacy. (b) The lowest possible dose should be prescribed to improve pain and function. (c) Office: Ongoing review and documentation of pain relief, functional status, appropriate medication use, and side effects. Pain assessment should include: current pain; the least reported pain over the period since last assessment; average pain; intensity of pain after taking the opioid; how long it takes for pain relief; and how long pain relief lasts. Satisfactory response to treatment may be indicated by the patient's decreased pain, increased level of function, or improved quality of life. Information from family members or other caregivers should be considered in determining the patient's response to treatment. The 4 A's for Ongoing Monitoring: Four domains have been proposed as most relevant for ongoing monitoring of chronic pain patients on opioids: pain relief, side effects, physical and psychosocial functioning, and the occurrence of any potentially aberrant (or nonadherent) drug-related behaviors. These domains have been summarized as the "4 A's" (analgesia, activities of daily living, adverse side effects, and aberrant drug taking behaviors). The monitoring of these outcomes over time should affect therapeutic decisions and provide a framework for documentation of the clinical use of these controlled drugs. (Passik, 2000) (d) Home: To aid in pain and functioning assessment, the patient should be requested to keep a pain dairy that includes entries such as pain triggers, and incidence of end-of-dose pain. It should be emphasized that using this diary will help in tailoring

the opioid dose. This should not be a requirement for pain management. (e) Use of drug screening or inpatient treatment with issues of abuse, addiction, or poor pain control. (f) Documentation of misuse of medications (doctor-shopping, uncontrolled drug escalation, drug diversion). (g) Continuing review of overall situation with regard to nonopioid means of pain control. (h) Consideration of a consultation with a multidisciplinary pain clinic if doses of opioids are required beyond what is usually required for the condition or pain does not improve on opioids in 3 months. Consider a psych consult if there is evidence of depression, anxiety or irritability. Consider an addiction medicine consult if there is evidence of substance misuse.

When to Continue Opioids (a) If the patient has returned to work. (b) If the patient has improved functioning and pain. (Washington, 2002) (Colorado, 2002) (Ontario, 2000) (VA/DoD, 2003) (Maddox-AAPM/APS, 1997) (Wisconsin, 2004) (Warfield, 2004) The long-term use of this medication class is not recommended per the California MTUS unless there documented evidence of benefit with measurable outcome measures and improvement in function. There is no documented improvement in VAS scores. There are also no objective measurements of improvement in function. Therefore, criteria for the ongoing use of opioids have not been met and the request is not certified.