MAXIMUS Federal Services

Case Number:	CM15-0006798		
Date Assigned:	01/26/2015	Date of Injury:	04/30/2009
Decision Date:	03/17/2015	UR Denial Date:	12/19/2014
Priority:	Standard	Application Received:	01/13/2015

HOW THE IMR FINAL DETERMINATION WAS MADE

MAXIMUS Federal Services sent the complete case file to an expert reviewer. He/she has no affiliation with the employer, employee, providers or the claims administrator. He/she has been in active clinical practice for more than five years and is currently working at least 24 hours a week in active practice. The expert reviewer was selected based on his/her clinical experience, education, background, and expertise in the same or similar specialties that evaluate and/or treat the medical condition and disputed items/Service. He/she is familiar with governing laws and regulations, including the strength of evidence hierarchy that applies to Independent Medical Review determinations.

The Expert Reviewer has the following credentials:
State(s) of Licensure: Ohio, West Virginia
Certification(s)/Specialty: Preventive Medicine, Occupational Medicine, Medical Toxicology

CLINICAL CASE SUMMARY

The expert reviewer developed the following clinical case summary based on a review of the case file, including all medical records:

The injured worker is a 66 year old male, who sustained an industrial injury on April 30, 2009. He has reported mid and lower back pain and pain of the left knee. The diagnoses have included chondromalacia of the patella, lumbar spine disc bulges and thoracic spine disc bulges. Treatment to date has included meds, physical therapy, Cold therapy, lumbar spine facet blocks, injections, home exercises and imaging studies. Currently, the injured worker complains of continued lower back pain radiating to the left leg and knee pain. The treating physician is requesting prescriptions for Norco, Anaprox and Fexmid.On December 19, 2014 Utilization Review non-certified the request for prescriptions for Norco, Anaprox and Fexmid noting the lack of documentation to support the medical necessity of the medications. The MTUS chronic pain medical treatment guidelines were cited in the decision.

IMR ISSUES, DECISIONS AND RATIONALES

The Final Determination was based on decisions for the disputed items/services set forth below:

Norco 5 MG #15: Upheld

Claims Administrator guideline: Decision based on MTUS Chronic Pain Treatment Guidelines Opioids.

MAXIMUS guideline: Decision based on MTUS Chronic Pain Treatment Guidelines opioids Page(s): 74-96. Decision based on Non-MTUS Citation Low Back - Lumbar & Thoracic (Acute & Chronic), Pain, Opioids

Decision rationale: ODG does not recommend the use of opioids for neck, low back, and shoulder pain "except for short use for severe cases, not to exceed 2 weeks." The patient has exceeded the 2 week recommended treatment length for opioid usage. MTUS does not discourage use of opioids past 2 weeks, but does state that "ongoing review and documentation of pain relief, functional status, appropriate medication use, and side effects. Pain assessment should include: current pain; the least reported pain over the period since last assessment; average pain; intensity of pain after taking the opioid; how long it takes for pain relief; and how long pain relief lasts. Satisfactory response to treatment may be indicated by the patient's decreased pain, increased level of function, or improved quality of life." The treating physician does not fully document the least reported pain over the period since last assessment, intensity of pain after taking opioid, pain relief, increased level of function, or improved quality of life. As such, the request for Norco 5mg x15 is deemed not medically necessary.

Anaprox DS 550 MG #60: Upheld

Claims Administrator guideline: Decision based on MTUS Chronic Pain Treatment Guidelines.

MAXIMUS guideline: Decision based on MTUS Chronic Pain Treatment Guidelines NSAIDs (non-steroidal anti-inflammatory drugs) Page(s): 67-73. Decision based on Non-MTUS Citation) Pain (Chronic), Naproxen, NSAIDs (non-steroidal anti-inflammatory drugs)

Decision rationale: CA-MTUS specifies four recommendations regarding NSAID use: 1) Osteoarthritis (including knee and hip): Recommended at the lowest dose for the shortest period in patients with moderate to severe pain. 2) Back Pain - Acute exacerbations of chronic pain: Recommended as a second-line treatment after acetaminophen. In general, there is conflicting evidence that NSAIDs are more effective that acetaminophen for acute LBP. 3) Back Pain - Chronic low back pain: Recommended as an option for short-term symptomatic relief. A Cochrane review of the literature on drug relief for low back pain (LBP) suggested that NSAIDs were no more effective than other drugs such as acetaminophen, narcotic analgesics, and muscle relaxants. The review also found that NSAIDs had more adverse effects than placebo and acetaminophen but fewer effects than muscle relaxants and narcotic analgesics. 4) Neuropathic pain: There is inconsistent evidence for the use of these medications to treat longterm neuropathic pain, but they may be useful to treat breakthrough and mixed pain conditions such as osteoarthritis (and other nociceptive pain) in with neuropathic pain. The medical documents do not indicate that the patient is being treated for osteoarthritis. Additionally, the treating physician does not document failure of primary (Tylenol) treatment. Progress notes do not indicate specifically how long the patient has been on naproxen but it appears to have been for an extended period with no documentation of functional improvement due to its use. MTUS guidelines recommend against long-term use and without documentation as to why this guideline should be extended in this case the request for anaprox 550mg must be deemed not medically necessary.

Fexmid 7.5 MG #60: Upheld

Claims Administrator guideline: Decision based on MTUS Chronic Pain Treatment Guidelines.

MAXIMUS guideline: Decision based on MTUS Chronic Pain Treatment Guidelines Cyclobenzaprine, Medications for chronic pain, Antispasmodics Page(s): 41-42, 60-61, 64-66. Decision based on Non-MTUS Citation Pain, Cyclobenzaprine (Flexerilï¿½) UpToDate, Flexeril

Decision rationale: MTUS Chronic Pain Medical Treatment states for Cyclobenzaprine, "Recommended as an option, using a short course of therapy... The effect is greatest in the first 4 days of treatment, suggesting that shorter courses may be better. (Browning, 2001) Treatment should be brief." The medical documents indicate that patient is far in excess of the initial treatment window and period.Additionally, MTUS outlines that "Relief of pain with the use of medications is generally temporary, and measures of the lasting benefit from this modality should include evaluating the effect of pain relief in relationship to improvements in function and increased activity. Before prescribing any medication for pain the following should occur: (1) determine the aim of use of the medication; (2) determine the potential benefits and adverse effects; (3) determine the patient's preference. Only one medication should be given at a time, and interventions that are active and passive should remain unchanged at the time of the medication change. A trial should be given for each individual medication. Analgesic medications should show effects within 1 to 3 days, and the analgesic effect of antidepressants should occur within 1 week. A record of pain and function with the medication should be recorded. (Mens, 2005)? Uptodate "flexeril" also recommends "Do not use longer than 2-3 weeks". Medical documents do not fully detail the components outlined in the guidelines above and do not establish the need for long term/chronic usage of cyclobenzaprine. As such, the request for Flexeril 7.5mg #60 is deemed not medically necessary.