

Case Number:	CM15-0009438		
Date Assigned:	02/12/2015	Date of Injury:	05/04/1992
Decision Date:	04/03/2015	UR Denial Date:	12/29/2014
Priority:	Standard	Application Received:	01/16/2015

HOW THE IMR FINAL DETERMINATION WAS MADE

MAXIMUS Federal Services sent the complete case file to an expert reviewer. He/she has no affiliation with the employer, employee, providers or the claims administrator. He/she has been in active clinical practice for more than five years and is currently working at least 24 hours a week in active practice. The expert reviewer was selected based on his/her clinical experience, education, background, and expertise in the same or similar specialties that evaluate and/or treat the medical condition and disputed items/Service. He/she is familiar with governing laws and regulations, including the strength of evidence hierarchy that applies to Independent Medical Review determinations.

The Expert Reviewer has the following credentials:
State(s) of Licensure: California, District of Columbia, Maryland
Certification(s)/Specialty: Anesthesiology, Pain Management

CLINICAL CASE SUMMARY

The expert reviewer developed the following clinical case summary based on a review of the case file, including all medical records:

The following clinical case summary was developed based on a review of the case file, including all medical records: The injured worker is a 56 year old male, who sustained an industrial injury reported on 5/4/1992. He has reported radiating low back pain. The diagnoses were noted to have included lumbar discopathy; and right knee meniscal tear. Treatments to date have included consultations; diagnostic imaging and urine studies; and medication management. The work status classification for this injured worker was not noted. On 12/29/2014, Utilization Review (UR) non-certified, for medical necessity, the request, made on 12/29/2014, for: Flurbiprofen 25%/Menthol 10%/Camphor 3%/Capsaicin 0.0375% topical cream, #30gm for radiating leg pain; Flurbiprofen 25%/Menthol 10%/Camphor 3%/Capsaicin 0.0375% topical cream, #120gm for radiating leg pain; Prilosec 20mg #90. The Medical Treatment Utilization Schedule, topical analgesics, proton-pump inhibitors, was cited.

IMR ISSUES, DECISIONS AND RATIONALES

The Final Determination was based on decisions for the disputed items/services set forth below:

Flurbiprofen25%/Menthol10%/Camphor3%/Capsaicin0.0375% topical cream, #30gm:
Upheld

Claims Administrator guideline: Decision based on MTUS Chronic Pain Treatment Guidelines.

MAXIMUS guideline: Decision based on MTUS Chronic Pain Treatment Guidelines Topical Analgesic Page(s): 60, 111-113.

Decision rationale: Capsaicin may have an indication for chronic lower back pain in this context. Per MTUS p 112 Indications: There are positive randomized studies with capsaicin cream in patients with osteoarthritis, fibromyalgia, and chronic non-specific back pain, but it should be considered experimental in very high doses. Although topical capsaicin has moderate to poor efficacy, it may be particularly useful (alone or in conjunction with other modalities) in patients whose pain has not been controlled successfully with conventional therapy. Per MTUS with regard to Flurbiprofen (p112), "(Biswal, 2006) these medications may be useful for chronic musculoskeletal pain, but there are no long-term studies of their effectiveness or safety. Indications: Osteoarthritis and tendinitis, in particular, that of the knee and elbow or other joints that are amenable to topical treatment: Recommended for short-term use (4-12 weeks). There is little evidence to utilize topical NSAIDs for treatment of osteoarthritis of the spine, hip or shoulder." The documentation contains no evidence of osteoarthritis or tendinitis. Flurbiprofen is not indicated. Regarding the use of multiple medications, MTUS p60 states "Only one medication should be given at a time, and interventions that are active and passive should remain unchanged at the time of the medication change. A trial should be given for each individual medication. Analgesic medications should show effects within 1 to 3 days, and the analgesic effect of antidepressants should occur within 1 week. A record of pain and function with the medication should be recorded. (Mens, 2005) The recent AHRQ review of comparative effectiveness and safety of analgesics for osteoarthritis concluded that each of the analgesics was associated with a unique set of benefits and risks, and no currently available analgesic was identified as offering a clear overall advantage compared with the others." Therefore, it would be optimal to trial each medication individually. The CA MTUS, ODG, National Guidelines Clearinghouse, and ACOEM provide no evidence-based recommendations regarding the topical application of menthol or camphor. It is the opinion of this IMR reviewer that a lack of endorsement, a lack of mention, inherently implies a lack of recommendation, or a status equivalent to "not recommended". Since several components are not medically indicated, then the overall product is not indicated per MTUS as outlined below. Note the statement on page 111: Any compounded product that contains at least one drug (or drug class) that is not recommended is not recommended. The request is not medically necessary.

Flurbiprofen25%/Menthol10%/Camphor3%/Capsaicin0.0375% topical cream #120gm:
Upheld

Claims Administrator guideline: Decision based on MTUS Chronic Pain Treatment Guidelines.

MAXIMUS guideline: Decision based on MTUS Chronic Pain Treatment Guidelines Topical Analgesics Page(s): 60, 111-113.

Decision rationale: Capsaicin may have an indication for chronic lower back pain in this context. Per MTUS p 112 Indications: There are positive randomized studies with capsaicin cream in patients with osteoarthritis, fibromyalgia, and chronic non-specific back pain, but it should be considered experimental in very high doses. Although topical capsaicin has moderate to poor efficacy, it may be particularly useful (alone or in conjunction with other modalities) in patients whose pain has not been controlled successfully with conventional therapy. Per MTUS with regard to Flurbiprofen (p112), "(Biswal, 2006) these medications may be useful for chronic musculoskeletal pain, but there are no long-term studies of their effectiveness or safety. Indications: Osteoarthritis and tendinitis, in particular, that of the knee and elbow or other joints that are amenable to topical treatment: Recommended for short-term use (4-12 weeks). There is little evidence to utilize topical NSAIDs for treatment of osteoarthritis of the spine, hip or shoulder." The documentation contains no evidence of osteoarthritis or tendinitis. Flurbiprofen is not indicated. Regarding the use of multiple medications, MTUS p60 states "Only one medication should be given at a time, and interventions that are active and passive should remain unchanged at the time of the medication change. A trial should be given for each individual medication. Analgesic medications should show effects within 1 to 3 days, and the analgesic effect of antidepressants should occur within 1 week. A record of pain and function with the medication should be recorded. (Mens, 2005) The recent AHRQ review of comparative effectiveness and safety of analgesics for osteoarthritis concluded that each of the analgesics was associated with a unique set of benefits and risks, and no currently available analgesic was identified as offering a clear overall advantage compared with the others." Therefore, it would be optimal to trial each medication individually. The CA MTUS, ODG, National Guidelines Clearinghouse, and ACOEM provide no evidence-based recommendations regarding the topical application of menthol or camphor. It is the opinion of this IMR reviewer that a lack of endorsement, a lack of mention, inherently implies a lack of recommendation, or a status equivalent to "not recommended". Since several components are not medically indicated, then the overall product is not indicated per MTUS as outlined below. Note the statement on page 111: Any compounded product that contains at least one drug (or drug class) that is not recommended is not recommended. The request is not medically necessary.

Prilosec 20mg #90: Upheld

Claims Administrator guideline: Decision based on MTUS Chronic Pain Treatment Guidelines.

MAXIMUS guideline: Decision based on MTUS Chronic Pain Treatment Guidelines NSAIDs, GI symptoms & cardiovascular risk Page(s): 68.

Decision rationale: In the treatment of dyspepsia secondary to NSAID therapy, the MTUS recommends stopping the NSAID, switching to a different NSAID, or considering the use of an H2-receptor antagonist or a PPI. The MTUS Chronic Pain Medical Treatment Guidelines recommend the use of proton pump inhibitors in conjunction with NSAIDs in situations in which the patient is at risk for gastrointestinal events including: (1) age > 65 years; (2) history of peptic ulcer, GI bleeding or perforation; (3) concurrent use of ASA, corticosteroids, and/or an anticoagulant; or (4) high dose/multiple NSAID (e.g., NSAID + low-dose ASA). CPMTG guidelines further specify: "Recommendations: Patients with no risk factor and no cardiovascular

disease: Non-selective NSAIDs OK (e.g, ibuprofen, naproxen, etc.) Patients at intermediate risk for gastrointestinal events and no cardiovascular disease: (1) A non-selective NSAID with either a PPI (Proton Pump Inhibitor, for example, 20 mg omeprazole daily) or misoprostol (200 g four times daily) or (2) a Cox-2 selective agent. Long-term PPI use (> 1 year) has been shown to increase the risk of hip fracture (adjusted odds ratio 1.44). Patients at high risk for gastrointestinal events with no cardiovascular disease: A Cox-2 selective agent plus a PPI if absolutely necessary. Patients at high risk of gastrointestinal events with cardiovascular disease: If GI risk is high the suggestion is for a low-dose Cox-2 plus low dose Aspirin (for cardioprotection) and a PPI. If cardiovascular risk is greater than GI risk the suggestion is naproxyn plus low-dose aspirin plus a PPI. (Laine, 2006) (Scholmerich, 2006) (Nielsen, 2006) (Chan, 2004) (Gold, 2007) (Laine, 2007)" as there is no documentation of peptic ulcer, GI bleeding or perforation, or cardiovascular disease in the records available for my review, the injured worker's risk for gastrointestinal events is low, as such, medical necessity cannot be affirmed.