

Case Number:	CM15-0101462		
Date Assigned:	06/03/2015	Date of Injury:	09/12/2003
Decision Date:	07/02/2015	UR Denial Date:	05/06/2015
Priority:	Standard	Application Received:	05/27/2015

HOW THE IMR FINAL DETERMINATION WAS MADE

MAXIMUS Federal Services sent the complete case file to an expert reviewer. He/she has no affiliation with the employer, employee, providers or the claims administrator. He/she has been in active clinical practice for more than five years and is currently working at least 24 hours a week in active practice. The expert reviewer was selected based on his/her clinical experience, education, background, and expertise in the same or similar specialties that evaluate and/or treat the medical condition and disputed items/Service. He/she is familiar with governing laws and regulations, including the strength of evidence hierarchy that applies to Independent Medical Review determinations.

The Expert Reviewer has the following credentials:
 State(s) of Licensure: North Carolina
 Certification(s)/Specialty: Family Practice

CLINICAL CASE SUMMARY

The expert reviewer developed the following clinical case summary based on a review of the case file, including all medical records:

The injured worker is a 58 year old female, who sustained an industrial injury on 9/12/03. Initial complaints were not reviewed. The injured worker was diagnosed as having right wrist carpal tunnel syndrome; right shoulder surgery x2. Treatment to date has included status post right shoulder surgery x2 (no dates); status post right carpal tunnel release (no dates). Currently, the PR-2 notes dated 2/19/15 indicated the injured worker complains of right shoulder pain rated at 7/10; constant right wrist pain rated as 9/10 with numbness and tingling. Objective findings show range of motion: flexion 120 degrees, extension 25 degrees, abduction 110 degrees, adduction 40 degrees, internal rotation 60 degrees and external rotation 60 degrees. The right wrist range of motion: flexion 40 degrees; extension 35 degrees, radial deviation 10 degrees, ulnar deviation 20 degrees. The provider has requested Ambien 10mg #30; Norco 10/325mg #140 and Soma 350mg #90.

IMR ISSUES, DECISIONS AND RATIONALES

The Final Determination was based on decisions for the disputed items/services set forth below:

Ambien 10mg #30: Upheld

Claims Administrator guideline: The Claims Administrator did not base their decision on the MTUS. Decision based on Non-MTUS Citation Official Disability Guidelines (ODG) Pain Chapter, Zolpidem (Ambien).

MAXIMUS guideline: The Expert Reviewer did not base their decision on the MTUS. Decision based on Non-MTUS Citation ODG, insomnia.

Decision rationale: The California MTUS and the ACOEM do not specifically address this medication. Per the official disability guidelines recommend pharmacological agents for insomnia only is used after careful evaluation of potential causes of sleep disturbance. Primary insomnia is usually addressed pharmacologically. Secondary insomnia may be treated with pharmacological and/or psychological measures. Pharmacological treatment consists of four main categories: Benzodiazepines, Non-benzodiazepines, Melatonin and melatonin receptor agonists and over the counter medications. Sedating antidepressants have also been used to treat insomnia however, there is less evidence to support their use for insomnia, but they may be an option inpatient's with coexisting depression. The patient does not have the diagnosis of primary insomnia or depression There is no provided clinical documentation of failure of sleep hygiene measures/counseling. Therefore, the request is not medically necessary.

Norco 10/325mg #140: Upheld

Claims Administrator guideline: Decision based on MTUS Chronic Pain Treatment Guidelines Opioids, criteria for use.

MAXIMUS guideline: Decision based on MTUS Chronic Pain Treatment Guidelines opioids Page(s): 76-84.

Decision rationale: The California chronic pain medical treatment guidelines section on opioids states for ongoing management: On-Going Management. Actions Should Include: (a) Prescriptions from a single practitioner taken as directed, and all prescriptions from a single pharmacy. (b) The lowest possible dose should be prescribed to improve pain and function. (c) Office: Ongoing review and documentation of pain relief, functional status, appropriate medication use, and side effects. Pain assessment should include: current pain; the least reported pain over the period since last assessment; average pain; intensity of pain after taking the opioid; how long it takes for pain relief; and how long pain relief lasts. Satisfactory response to treatment may be indicated by the patient's decreased pain, increased level of function, or improved quality of life. Information from family members or other caregivers should be considered in determining the patient's response to treatment. The 4 A's for Ongoing Monitoring: Four domains have been proposed as most relevant for ongoing monitoring of chronic pain patients on opioids: pain relief, side-effects, physical and psychosocial functioning, and the occurrence of any potentially aberrant (or non-adherent) drug-related behaviors. These domains have been summarized as the "4 A's" (analgesia, activities of daily living, adverse side effects, and aberrant drug taking behaviors). The monitoring of these outcomes over time should affect therapeutic decisions and provide a framework for documentation of the clinical use of these controlled drugs. (Passik, 2000) (d) Home: To aid in pain and functioning assessment, the patient should be requested to keep a pain diary that includes entries such as pain triggers, and incidence of end-of-dose pain. It should be emphasized that using this diary will help in tailoring the opioid dose. This should not be a requirement for pain management. (e) Use of drug screening or inpatient treatment with issues of abuse, addiction, or poor pain control. (f) Documentation of misuse of medications (doctor- shopping, uncontrolled drug escalation, drug diversion). (g) Continuing review of overall situation with regard to non-opioid means of pain control. (h) Consideration of

a consultation with a multidisciplinary pain clinic if doses of opioids are required beyond what is usually required for the condition or pain does not improve on opioids in 3 months. Consider a psych consult if there is evidence of depression, anxiety or irritability. Consider an addiction medicine consult if there is evidence of substance misuse. When to Continue Opioids; (a) If the patient has returned to work. (b) If the patient has improved functioning and pain. (Washington, 2002) (Colorado, 2002) (Ontario, 2000) (VA/DoD, 2003) (Maddox-AAPM/APS, 1997) (Wisconsin, 2004) (Warfield, 2004). The long-term use of this medication class is not recommended per the California MTUS unless there documented evidence of benefit with measurable outcome measures and improvement in function. There is no documented significant improvement in VAS scores for significant periods of time. There are no objective measurements of improvement in function. Therefore all criteria for the ongoing use of opioids have not been met and the request is not medically necessary.

Soma 350mg #90: Upheld

Claims Administrator guideline: Decision based on MTUS Chronic Pain Treatment Guidelines Muscle Relaxants (for pain).

MAXIMUS guideline: Decision based on MTUS Chronic Pain Treatment Guidelines muscle relaxants Page(s): 63-65.

Decision rationale: The California chronic pain medical treatment guidelines section on muscle relaxants states: Recommend non-sedating muscle relaxants with caution as a second-line option for short-term treatment of acute exacerbations in patients with chronic LBP. (Chou, 2007) (Mens, 2005) (Van Tulder, 1998) (van Tulder, 2003) (van Tulder, 2006) (Schnitzer, 2004) (See, 2008) Muscle relaxants may be effective in reducing pain and muscle tension, and increasing mobility. However, in most LBP cases, they show no benefit beyond NSAIDs in pain and overall improvement. Also, there is no additional benefit shown in combination with NSAIDs. Efficacy appears to diminish over time, and prolonged use of some medications in this class may lead to dependence. (Homik, 2004) (Chou, 2004) This medication is not intended for long-term use per the California MTUS. The medication has not been prescribed for the flare-up of chronic low back pain. This is not an approved use for the medication. For these reasons, criteria for the use of this medication have not been met. Therefore, the request is not medically necessary.