

Case Number:	CM15-0102090		
Date Assigned:	06/04/2015	Date of Injury:	12/16/2000
Decision Date:	07/03/2015	UR Denial Date:	05/26/2015
Priority:	Standard	Application Received:	05/27/2015

HOW THE IMR FINAL DETERMINATION WAS MADE

MAXIMUS Federal Services sent the complete case file to an expert reviewer. He/she has no affiliation with the employer, employee, providers or the claims administrator. He/she has been in active clinical practice for more than five years and is currently working at least 24 hours a week in active practice. The expert reviewer was selected based on his/her clinical experience, education, background, and expertise in the same or similar specialties that evaluate and/or treat the medical condition and disputed items/Service. He/she is familiar with governing laws and regulations, including the strength of evidence hierarchy that applies to Independent Medical Review determinations.

The Expert Reviewer has the following credentials:

State(s) of Licensure: North Carolina

Certification(s)/Specialty: Family Practice

CLINICAL CASE SUMMARY

The expert reviewer developed the following clinical case summary based on a review of the case file, including all medical records:

The injured worker is a 64 year old male with an industrial injury dated 12/16/2000. The mechanism of injury is documented as a fall landing on a concrete floor striking his head, shoulder and lower back. His diagnoses included post laminectomy syndrome, lumbar; cervicgia, and lumbago. Prior treatment included medications and diagnostics. He presents on 03/20/2015 with complaints of lower back sore in the morning before he took his pain medication. His medications were Norco, Lidoderm patch, Norflex and Ultram. The progress note dated 03/20/2015 documents bulging disk cervical 3, 4, and 5 and bulging disk lumbar 5 and 6. There was tenderness at lumbar 4-5 with no palpable paraspinal muscle spasm. The injured worker walked with a slow guarded gait. Bilateral toe walk was demonstrated but he was unable to heel walk on left side due to heel pain. MRI of lumbar spine had been requested. There were no reports of MRI in the submitted records. The request is for urine analysis, lumbar epidural steroid facet injection lumbar 4-sacral 1 times 2 and post-op physical therapy three times four.

IMR ISSUES, DECISIONS AND RATIONALES

The Final Determination was based on decisions for the disputed items/services set forth below:

Lumbar epidural steroid facet injection L4-S1 x 2: Upheld

Claims Administrator guideline: The Claims Administrator did not base their decision on the MTUS. Decision based on Non-MTUS Citation Official Disability Guidelines, Criteria for the use of epidural steroid injections.

MAXIMUS guideline: Decision based on MTUS Chronic Pain Treatment Guidelines epidural steroid injections Page(s): 46.

Decision rationale: The California chronic pain medical treatment guidelines section on epidural steroid injections (ESI) states: Criteria for the use of Epidural steroid injections: Note: The purpose of ESI is to reduce pain and inflammation, restoring range of motion and thereby facilitating progress in more active treatment programs, and avoiding surgery, but this treatment alone offers no significant long-term functional benefit. 1) Radiculopathy must be documented by physical examination and corroborated by imaging studies and/or electrodiagnostic testing. 2) Initially unresponsive to conservative treatment (exercises, physical methods, NSAIDs and muscle relaxants). 3) Injections should be performed using fluoroscopy (live x-ray) for guidance. 4) If used for diagnostic purposes, a maximum of two injections should be performed. A second block is not recommended if there is inadequate response to the first block. Diagnostic blocks should be at an interval of at least one to two weeks between injections. 5) No more than two nerve root levels should be injected using transforaminal blocks. 6) No more than one interlaminar level should be injected at one session. 7) In the therapeutic phase, repeat blocks should be based on continued objective documented pain and functional improvement, including at least 50% pain relief with associated reduction of medication use for six to eight weeks, with a general recommendation of no more than 4 blocks per region per year. (Manchikanti, 2003) (CMS, 2004) (Boswell, 2007) 8) Current research does not support a series-of-three injections in either the diagnostic or therapeutic phase. We recommend no more than 2 ESI injections. The request is not for diagnostic injection so a series of two is not indicated per the California MTUS and the request is not medically necessary.

Labs; Urine analysis: Overturned

Claims Administrator guideline: Decision based on MTUS Chronic Pain Treatment Guidelines Opioids. Decision based on Non-MTUS Citation Official Disability Guidelines, Urine drug testing.

MAXIMUS guideline: Decision based on MTUS Chronic Pain Treatment Guidelines opioids Page(s): 76-84.

Decision rationale: The California chronic pain medical treatment guidelines section on opioids states: On-Going Management. Actions Should Include: (a) Prescriptions from a single practitioner taken as directed, and all prescriptions from a single pharmacy. (b) The lowest possible dose should be prescribed to improve pain and function. (c) Office: Ongoing review and documentation of pain relief, functional status, appropriate medication use, and side effects. Pain assessment should include: current pain; the least reported pain over the period since last assessment; average pain; intensity of pain after taking the opioid; how long it takes for pain relief; and how long pain relief lasts. Satisfactory response to treatment may be indicated by the patient's decreased pain, increased level of function, or improved quality of life. Information

from family members or other caregivers should be considered in determining the patient's response to treatment. The 4 A's for Ongoing Monitoring: Four domains have been proposed as most relevant for ongoing monitoring of chronic pain patients on opioids: pain relief, side effects, physical and psychosocial functioning, and the occurrence of any potentially aberrant (or non-adherent) drug-related behaviors. These domains have been summarized as the "4 A's" (analgesia, activities of daily living, adverse side effects, and aberrant drug taking behaviors). The monitoring of these outcomes over time should affect therapeutic decisions and provide a framework for documentation of the clinical use of these controlled drugs. (Passik, 2000) (d) Home: To aid in pain and functioning assessment, the patient should be requested to keep a pain diary that includes entries such as pain triggers, and incidence of end-of-dose pain. It should be emphasized that using this diary will help in tailoring the opioid dose. This should not be a requirement for pain management. (e) Use of drug screening or inpatient treatment with issues of abuse, addiction, or poor pain control. (f) Documentation of misuse of medications (doctor- shopping, uncontrolled drug escalation, drug diversion). (g) Continuing review of overall situation with regard to non-opioid means of pain control. (h) Consideration of a consultation with a multidisciplinary pain clinic if doses of opioids are required beyond what is usually required for the condition or pain does not improve on opioids in 3 months. Consider a psych consult if there is evidence of depression, anxiety or irritability. Consider an addiction medicine consult if there is evidence of substance misuse. The California MTUS does recommend urine drug screens as part of the criteria for ongoing use of opioids .The patient was on opioids at the time of request and therefore the request is medically necessary.

Post-op physical therapy 3 x 4: Upheld

Claims Administrator guideline: Decision based on MTUS Chronic Pain Treatment Guidelines Physical Medicine Page(s): 98-99.

MAXIMUS guideline: Decision based on MTUS Chronic Pain Treatment Guidelines physical medicine Page(s): 98-99.

Decision rationale: The California chronic pain medical treatment guidelines section on physical medicine states: Recommended as indicated below. Passive therapy (those treatment modalities that do not require energy expenditure on the part of the patient) can provide short term relief during the early phases of pain treatment and are directed at controlling symptoms such as pain, inflammation and swelling and to improve the rate of healing soft tissue injuries. They can be used sparingly with active therapies to help control swelling, pain and inflammation during the rehabilitation process. Active therapy is based on the philosophy that therapeutic exercise and/or activity are beneficial for restoring flexibility, strength, endurance, function, range of motion, and can alleviate discomfort. Active therapy requires an internal effort by the individual to complete a specific exercise or task. This form of therapy may require supervision from a therapist or medical provider such as verbal, visual and/or tactile instruction(s). Patients are instructed and expected to continue active therapies at home as an extension of the treatment process in order to maintain improvement levels. Home exercise can include exercise with or without mechanical assistance or resistance and functional activities with assistive devices. (Colorado, 2002) (Airaksinen, 2006) Patient-specific hand therapy is very important in reducing swelling, decreasing pain, and improving range of motion in CRPS. (Li, 2005) The use of active treatment modalities (e.g., exercise, education, activity modification) instead of passive treatments is associated with substantially better clinical outcomes. In a large case series of patients with low back pain treated by physical therapists,

those adhering to guidelines for active rather than passive treatments incurred fewer treatment visits, cost less, and had less pain and less disability. The overall success rates were 64.7% among those adhering to the active treatment recommendations versus 36.5% for passive treatment. (Fritz, 2007) Physical Medicine Guidelines: Allow for fading of treatment frequency (from up to 3 visits per week to 1 or less), plus active self-directed home Physical Medicine. Myalgia and myositis, unspecified (ICD9 729.1): 9-10 visits over 8 weeks-Neuralgia, neuritis, and radiculitis, unspecified (ICD9 729.2) 8-10 visits over 4 weeks, Reflex sympathetic dystrophy (CRPS) (ICD9 337.2): 24 visits over 16 weeks. The goal of physical therapy is graduation to home therapy after a certain amount of recommended sessions. The request is in excess of these recommendations per the California MTUS. There is no explanation why the patient would not be moved to home therapy after completing the recommended amount of supervised sessions.. Therefore the request is not medically necessary.