

Case Number:	CM15-0103643		
Date Assigned:	06/08/2015	Date of Injury:	09/22/2013
Decision Date:	07/07/2015	UR Denial Date:	05/22/2015
Priority:	Standard	Application Received:	05/29/2015

HOW THE IMR FINAL DETERMINATION WAS MADE

MAXIMUS Federal Services sent the complete case file to an expert reviewer. He/she has no affiliation with the employer, employee, providers or the claims administrator. He/she has been in active clinical practice for more than five years and is currently working at least 24 hours a week in active practice. The expert reviewer was selected based on his/her clinical experience, education, background, and expertise in the same or similar specialties that evaluate and/or treat the medical condition and disputed items/Service. He/she is familiar with governing laws and regulations, including the strength of evidence hierarchy that applies to Independent Medical Review determinations.

The Expert Reviewer has the following credentials:
 State(s) of Licensure: New Jersey, Alabama, California
 Certification(s)/Specialty: Neurology, Neuromuscular Medicine

CLINICAL CASE SUMMARY

The expert reviewer developed the following clinical case summary based on a review of the case file, including all medical records:

The injured worker is a 62-year-old male, who sustained an industrial injury on 09/22/2013 when the handle of the air brake he was lifting broke up, and caused the left arm to fully extend. He has reported subsequent neck and upper extremity pain and was diagnosed with chronic pain and left cervical radiculitis. Treatment to date has included oral pain medication, physical therapy, cervical epidural injections and surgery. In a progress note dated 05/15/2015, the injured worker complained of increasing right hand pain, chronic neck pain with radiation to the left upper extremity and anxiety and depression about an upcoming anterior cervical decompression surgery that was scheduled. The physician noted that the injured worker had completed 5 out of 6 psychotherapy and biofeedback sessions with gradually decreasing pain, improved catastrophizing, mild depression and mild anxiety. The plan was to continue biofeedback therapy. A request for authorization of 6 sessions of biofeedback was submitted.

IMR ISSUES, DECISIONS AND RATIONALES

The Final Determination was based on decisions for the disputed items/services set forth below:

Six sessions of biofeedback: Upheld

Claims Administrator guideline: Decision based on MTUS Chronic Pain Treatment Guidelines Biofeedback. Decision based on Non-MTUS Citation Official Disability Guidelines, Biofeedback Therapy.

MAXIMUS guideline: The Expert Reviewer did not base their decision on the MTUS. Decision based on Non-MTUS Citation Biofeedback. <http://www.odg-twc.com/index.html>.

Decision rationale: According to ODG guidelines, biofeedback "Not recommended as a stand-alone treatment, but recommended as an option in a cognitive behavioral therapy (CBT) program to facilitate exercise therapy and return to activity. There is good evidence that biofeedback helps in back muscle strengthening, but evidence is insufficient to demonstrate the effectiveness of biofeedback for treatment of chronic pain. Biofeedback may be approved if it facilitates entry into a CBT treatment program, where there is strong evidence of success. As with yoga, since outcomes from biofeedback are very dependent on the highly motivated self-disciplined patient, we recommend approval only when requested by such a patient, but not adoption for use by any patient. EMG biofeedback may be used as part of a behavioral treatment program, with the assumption that the ability to reduce muscle tension will be improved through feedback of data regarding degree of muscle tension to the subject. The potential benefits of biofeedback include pain reduction because the patient may gain a feeling that he is in control and pain is a manageable symptom. Biofeedback techniques are likely to use surface EMG feedback so the patient learns to control the degree of muscle contraction. The available evidence does not clearly show whether biofeedback's effects exceed nonspecific placebo effects. It is also unclear whether biofeedback adds to the effectiveness of relaxation training alone. The application of biofeedback to patients with CRPS is not well researched. However, based on CRPS symptomology, temperature or skin conductance feedback modalities may be of particular interest. (Keefe, 1981) (Nouwen, 1983) (Bush, 1985) (Croce, 1986) (Stuckey, 1986) (Asfour, 1990) (Altmaier, 1992) (Flor, 1993) (Newton-John, 1995) (Spence, 1995) (Vlaeyen, 1995) (NIH-JAMA, 1996) (van Tulder, 1997) (Buckelew, 1998) (Hasenbring, 1999) (Dursun, 2001) (van Santen, 2002) (Astin, 2002) (State, 2002) (BlueCross BlueShield, 2004) This recent report on 11 chronic whiplash patients found that, after 4 weeks of myofeedback training, there was a trend for decreased disability in 36% of the patients. The authors recommended a randomized-controlled trial to further explore the effects of myofeedback training. (Voerman, 2006) See also Cognitive behavioral therapy (Psychological treatment) and Cognitive intervention (Behavioral treatment) in the Low Back Chapter. Functional MRI has been proposed as a method to control brain activation of pain. See Functional imaging of brain responses to pain. ODG biofeedback therapy guidelines: Screen for patients with risk factors for delayed recovery, as well as motivation to comply with a treatment regimen that requires self-discipline. Initial therapy for these "at risk" patients should be physical therapy exercise instruction, using a cognitive motivational approach to PT. Possibly consider biofeedback referral in conjunction with CBT after 4 weeks: Initial trial of 3-4 psychotherapy visits over 2 weeks, With evidence of objective functional improvement, total of up to 6-10 visits over 5-6 weeks (individual sessions). Patients may continue biofeedback exercises at home." The patient anxiety and depression improved with previous biofeedback therapy and the need for more sessions in addition to the 6 performed sessions is established. However approving 6 more sessions cannot be done without documentation of efficacy of the first additional sessions. Therefore, the request for Six sessions of biofeedback is not medically necessary.