

Case Number:	CM15-0104000		
Date Assigned:	06/08/2015	Date of Injury:	02/26/2011
Decision Date:	07/08/2015	UR Denial Date:	05/12/2015
Priority:	Standard	Application Received:	05/29/2015

HOW THE IMR FINAL DETERMINATION WAS MADE

MAXIMUS Federal Services sent the complete case file to an expert reviewer. He/she has no affiliation with the employer, employee, providers or the claims administrator. He/she has been in active clinical practice for more than five years and is currently working at least 24 hours a week in active practice. The expert reviewer was selected based on his/her clinical experience, education, background, and expertise in the same or similar specialties that evaluate and/or treat the medical condition and disputed items/Service. He/she is familiar with governing laws and regulations, including the strength of evidence hierarchy that applies to Independent Medical Review determinations.

The Expert Reviewer has the following credentials:

State(s) of Licensure: North Carolina

Certification(s)/Specialty: Family Practice

CLINICAL CASE SUMMARY

The expert reviewer developed the following clinical case summary based on a review of the case file, including all medical records:

This 33-year-old female sustained an industrial injury on 2/26/11. She subsequently reported knee and low back pain. Diagnoses include right knee derangement of meniscus. Treatments to date include x-ray and MRI testing, TENS therapy, knee brace, physical therapy and prescription pain medications. The injured worker continues to experience right knee pain. Upon examination, reduced range of motion of the right lower extremity was noted. A request for Physical therapy 2 times a week for 6 weeks and TENS unit was made by the treating physician.

IMR ISSUES, DECISIONS AND RATIONALES

The Final Determination was based on decisions for the disputed items/services set forth below:

Physical therapy 2 times a week for 6 weeks: Upheld

Claims Administrator guideline: Decision based on MTUS Chronic Pain Treatment Guidelines Physical Medicine Page(s): 98-99.

MAXIMUS guideline: Decision based on MTUS Chronic Pain Treatment Guidelines physical medicine Page(s): 98-99.

Decision rationale: The California chronic pain medical treatment guidelines section on physical medicine states: Recommended as indicated below. Passive therapy (those treatment modalities that do not require energy expenditure on the part of the patient) can provide short term relief during the early phases of pain treatment and are directed at controlling symptoms such as pain, inflammation and swelling and to improve the rate of healing soft tissue injuries. They can be used sparingly with active therapies to help control swelling, pain and inflammation during the rehabilitation process. Active therapy is based on the philosophy that therapeutic exercise and/or activity are beneficial for restoring flexibility, strength, endurance, function, range of motion, and can alleviate discomfort. Active therapy requires an internal effort by the individual to complete a specific exercise or task. This form of therapy may require supervision from a therapist or medical provider such as verbal, visual and/or tactile instruction(s). Patients are instructed and expected to continue active therapies at home as an extension of the treatment process in order to maintain improvement levels. Home exercise can include exercise with or without mechanical assistance or resistance and functional activities with assistive devices. (Colorado, 2002) (Airaksinen, 2006) Patient-specific hand therapy is very important in reducing swelling, decreasing pain, and improving range of motion in CRPS. (Li, 2005) The use of active treatment modalities (e.g., exercise, education, activity modification) instead of passive treatments is associated with substantially better clinical outcomes. In a large case series of patients with low back pain treated by physical therapists, those adhering to guidelines for active rather than passive treatments incurred fewer treatment visits, cost less, and had less pain and less disability. The overall success rates were 64.7% among those adhering to the active treatment recommendations versus 36.5% for passive treatment. (Fritz, 2007) Physical Medicine Guidelines -Allow for fading of treatment frequency (from up to 3 visits per week to 1 or less), plus active self-directed home Physical Medicine. Myalgia and myositis, unspecified (ICD9 729.1): 9-10 visits over 8 weeks. Neuralgia, neuritis, and radiculitis, unspecified (ICD9 729.2): 8-10 visits over 4 weeks. Reflex sympathetic dystrophy (CRPS) (ICD9 337.2): 24 visits over 16 weeks. The requested amount of physical therapy is in excess of California chronic pain medical treatment guidelines. There is no objective explanation why the patient would need excess physical therapy and not be transitioned to active self-directed physical medicine. The request is not medically necessary.

TENS unit: Upheld

Claims Administrator guideline: Decision based on MTUS Chronic Pain Treatment Guidelines Page(s): 114-116.

MAXIMUS guideline: Decision based on MTUS Chronic Pain Treatment Guidelines TENS Page(s): 114.

Decision rationale: The California chronic pain medical treatment guidelines section on transcutaneous electrical nerve stimulation states: TENS, chronic pain (transcutaneous electrical nerve stimulation) Not recommended as a primary treatment modality, but a one-month home-based TENS trial may be considered as a noninvasive conservative option, if used as an adjunct to a program of evidence-based functional restoration, for the conditions described below. While TENS may reflect the long-standing accepted standard of care within many medical communities, the results of studies are inconclusive; the published trials do not provide

information on the stimulation parameters which are most likely to provide optimum pain relief, nor do they answer questions about long-term effectiveness. (Carroll-Cochrane, 2001) Several published evidence-based assessments of transcutaneous electrical nerve stimulation (TENS) have found that evidence is lacking concerning effectiveness. One problem with current studies is that many only evaluated single-dose treatment, which may not reflect the use of this modality in a clinical setting. Other problems include statistical methodology, small sample size, influence of placebo effect, and difficulty comparing the different outcomes that were measured. This treatment option is recommended as an adjunct to a program of evidence based functional restoration. In addition, there must be a 30-day trial period with objective measurements of improvement. These criteria have not been met and the request is not medically necessary.