

Case Number:	CM15-0104081		
Date Assigned:	06/08/2015	Date of Injury:	07/09/2012
Decision Date:	07/08/2015	UR Denial Date:	05/04/2015
Priority:	Standard	Application Received:	05/30/2015

HOW THE IMR FINAL DETERMINATION WAS MADE

MAXIMUS Federal Services sent the complete case file to an expert reviewer. He/she has no affiliation with the employer, employee, providers or the claims administrator. He/she has been in active clinical practice for more than five years and is currently working at least 24 hours a week in active practice. The expert reviewer was selected based on his/her clinical experience, education, background, and expertise in the same or similar specialties that evaluate and/or treat the medical condition and disputed items/Service. He/she is familiar with governing laws and regulations, including the strength of evidence hierarchy that applies to Independent Medical Review determinations.

The Expert Reviewer has the following credentials:
 State(s) of Licensure: North Carolina
 Certification(s)/Specialty: Family Practice

CLINICAL CASE SUMMARY

The expert reviewer developed the following clinical case summary based on a review of the case file, including all medical records:

The injured worker is a 41-year-old female, with a reported date of injury of 07/09/2012. The diagnoses include chronic left shoulder pain, status post rotator cuff repair and biceps tenodesis, left shoulder adhesive capsulitis, status post left shoulder manipulation and arthroscopic debridement, and status post anterior cervical discectomy and fusion. Treatments to date have included anterior cervical discectomy and fusion on 10/06/2014, oral medication, and physical therapy. The progress report dated 04/15/2015 indicates that the injured worker had persistent neck and left shoulder pain. The objective findings include spasms in the cervical paraspinal muscles, stiffness in the cervical spine, cervical forward flexion at 20 degrees, cervical extension at 15 degrees, cervical side rotation and bending at 45 degrees with pain, and limited mobility in the left shoulder due to pain. The progress report dated 03/20/2015 indicates that the injured worker reported her pain severity was 7.5 out of 10. She had not been able to use any medications for pain for the previous three weeks or so, because she had been denied at the pharmacy. It was noted that she had been doing fairly well using Butrans and Norco to relieve her pain symptoms. They reduced her pain level to 2-3 out of 10 and allowed her to increase her activity. She had not sign of sedation and reported no side effects with the medication use. The treating physician requested Norco 10/325mg #120, with three refills and Butrans patch 10mg #4, with three refills.

IMR ISSUES, DECISIONS AND RATIONALES

The Final Determination was based on decisions for the disputed items/services set forth below:

Norco 10/325mg #120 with 3 refills: Upheld

Claims Administrator guideline: Decision based on MTUS Chronic Pain Treatment Guidelines Opioids, Specific Drug List, Hydrocodone/Acetaminophen, Opioids, Criteria for Use, Weaning of Medications Page(s): 91, 78-80, 124.

MAXIMUS guideline: Decision based on MTUS Chronic Pain Treatment Guidelines opioids Page(s): 76-84.

Decision rationale: The California chronic pain medical treatment guidelines section on opioids states for ongoing management: On-Going Management. Actions Should Include: (a) Prescriptions from a single practitioner taken as directed, and all prescriptions from a single pharmacy. (b) The lowest possible dose should be prescribed to improve pain and function. (c) Office: Ongoing review and documentation of pain relief, functional status, appropriate medication use, and side effects. Pain assessment should include: current pain; the least reported pain over the period since last assessment; average pain; intensity of pain after taking the opioid; how long it takes for pain relief; and how long pain relief lasts. Satisfactory response to treatment may be indicated by the patient's decreased pain, increased level of function, or improved quality of life. Information from family members or other caregivers should be considered in determining the patient's response to treatment. The 4 A's for Ongoing Monitoring: Four domains have been proposed as most relevant for ongoing monitoring of chronic pain patients on opioids: pain relief, side effects, physical and psychosocial functioning, and the occurrence of any potentially aberrant (or nonadherent) drug-related behaviors. These domains have been summarized as the "4 A's" (analgesia, activities of daily living, adverse side effects, and aberrant drug taking behaviors). The monitoring of these outcomes over time should affect therapeutic decisions and provide a framework for documentation of the clinical use of these controlled drugs. (Passik, 2000) (d) Home: To aid in pain and functioning assessment, the patient should be requested to keep a pain diary that includes entries such as pain triggers, and incidence of end-of-dose pain. It should be emphasized that using this diary will help in tailoring the opioid dose. This should not be a requirement for pain management. (e) Use of drug screening or inpatient treatment with issues of abuse, addiction, or poor pain control. (f) Documentation of misuse of medications (doctor-shopping, uncontrolled drug escalation, drug diversion). (g) Continuing review of overall situation with regard to nonopioid means of pain control. (h) Consideration of a consultation with a multidisciplinary pain clinic if doses of opioids are required beyond what is usually required for the condition or pain does not improve on opioids in 3 months. Consider a psych consult if there is evidence of depression, anxiety or irritability. Consider an addiction medicine consult if there is evidence of substance misuse. When to Continue Opioids (a) If the patient has returned to work, (b) If the patient has improved functioning and pain. (Washington, 2002) (Colorado, 2002) (Ontario, 2000) (VA/DoD, 2003) (Maddox-AAPM/APS, 1997) (Wisconsin, 2004) (Warfield, 2004) The long-term use of this medication class is not recommended per the California MTUS unless there documented evidence of benefit with measurable outcome measures and improvement in function. There is documented significant decrease in objective pain measures such as VAS scores for significant periods of time. There are no objective measures of improvement of function. Therefore all criteria for the ongoing use of opioids have not been met and the request is not medically necessary.

Butrans patch 10mg #4 with 3 refills: Upheld

Claims Administrator guideline: Decision based on MTUS Chronic Pain Treatment Guidelines Topical Analgesics, Buprenorphine Page(s): 111, 26-37.

MAXIMUS guideline: Decision based on MTUS Chronic Pain Treatment Guidelines opioids Page(s): 76-84.

Decision rationale: The California chronic pain medical treatment guidelines section on opioids states for ongoing management: On-Going Management. Actions Should Include: (a) Prescriptions from a single practitioner taken as directed, and all prescriptions from a single pharmacy. (b) The lowest possible dose should be prescribed to improve pain and function. (c) Office: Ongoing review and documentation of pain relief, functional status, appropriate medication use, and side effects. Pain assessment should include: current pain; the least reported pain over the period since last assessment; average pain; intensity of pain after taking the opioid; how long it takes for pain relief; and how long pain relief lasts. Satisfactory response to treatment may be indicated by the patient's decreased pain, increased level of function, or improved quality of life. Information from family members or other caregivers should be considered in determining the patient's response to treatment. The 4 A's for Ongoing Monitoring: Four domains have been proposed as most relevant for ongoing monitoring of chronic pain patients on opioids: pain relief, side effects, physical and psychosocial functioning, and the occurrence of any potentially aberrant (or nonadherent) drug-related behaviors. These domains have been summarized as the "4 A's" (analgesia, activities of daily living, adverse side effects, and aberrant drug taking behaviors). The monitoring of these outcomes over time should affect therapeutic decisions and provide a framework for documentation of the clinical use of these controlled drugs. (Passik, 2000) (d) Home: To aid in pain and functioning assessment, the patient should be requested to keep a pain diary that includes entries such as pain triggers, and incidence of end-of-dose pain. It should be emphasized that using this diary will help in tailoring the opioid dose. This should not be a requirement for pain management. (e) Use of drug screening or inpatient treatment with issues of abuse, addiction, or poor pain control. (f) Documentation of misuse of medications (doctor-shopping, uncontrolled drug escalation, drug diversion). (g) Continuing review of overall situation with regard to nonopioid means of pain control. (h) Consideration of a consultation with a multidisciplinary pain clinic if doses of opioids are required beyond what is usually required for the condition or pain does not improve on opioids in 3 months. Consider a psych consult if there is evidence of depression, anxiety or irritability. Consider an addiction medicine consult if there is evidence of substance misuse. When to Continue Opioids (a) If the patient has returned to work, (b) If the patient has improved functioning and pain. (Washington, 2002) (Colorado, 2002) (Ontario, 2000) (VA/DoD, 2003) (Maddox-AAPM/APS, 1997) (Wisconsin, 2004) (Warfield, 2004) The long-term use of this medication class is not recommended per the California MTUS unless there documented evidence of benefit with measurable outcome measures and improvement in function. There is documented significant decrease in objective pain measures such as VAS scores for significant periods of time. There are no objective measures of improvement of function. Therefore all criteria for the ongoing use of opioids have not been met and the request is not medically necessary.

