

Case Number:	CM15-0104854		
Date Assigned:	06/09/2015	Date of Injury:	02/04/2013
Decision Date:	07/10/2015	UR Denial Date:	04/30/2015
Priority:	Standard	Application Received:	06/01/2015

HOW THE IMR FINAL DETERMINATION WAS MADE

MAXIMUS Federal Services sent the complete case file to an expert reviewer. He/she has no affiliation with the employer, employee, providers or the claims administrator. He/she has been in active clinical practice for more than five years and is currently working at least 24 hours a week in active practice. The expert reviewer was selected based on his/her clinical experience, education, background, and expertise in the same or similar specialties that evaluate and/or treat the medical condition and disputed items/Service. He/she is familiar with governing laws and regulations, including the strength of evidence hierarchy that applies to Independent Medical Review determinations.

The Expert Reviewer has the following credentials:

State(s) of Licensure: Iowa, Illinois, Hawaii

Certification(s)/Specialty: Preventive Medicine, Occupational Medicine, Public Health & General Preventive Medicine

CLINICAL CASE SUMMARY

The expert reviewer developed the following clinical case summary based on a review of the case file, including all medical records:

The injured worker is a 49-year-old female, who sustained an industrial injury on 02/04/2013. According to a progress report dated 04/16/2015, the injured worker had chronic pain. Location of pain included the shoulder, hands, fingers and bilateral neck with radiation of pain to the right hand. Associated symptoms included joint stiffness and tenderness of the left shoulder. Numbness and tingling in the left upper extremity was noted. She continued to have disrupted sleep pattern. The injured worker had been taking Flexeril and reported an 80 percent decrease in pain and spasm. Adverse side effects include sedation, which helped with her insomnia. Lidoderm patches helped 80 percent with pain relief at site. Voltaren gel was used with at least 50 percent reduction in pain levels. Diagnoses included rotator cuff syndrome, displacement of cervical intervertebral disc without myelopathy and degeneration of cervical intervertebral disc. Prescriptions included Celebrex, Cyclobenzaprine, Lidocaine 5% patch, and Omeprazole. The injured worker was temporarily totally disabled until next visit pending surgical evaluation. Currently under review is the request for Cyclobenzaprine 10mg quantity 30 and Lidocaine Patch 5% quantity 30.

IMR ISSUES, DECISIONS AND RATIONALES

The Final Determination was based on decisions for the disputed items/services set forth below:

Cyclobenzaprine 10 MG Qty 30: Upheld

Claims Administrator guideline: Decision based on MTUS Chronic Pain Treatment Guidelines NSAIDs.

MAXIMUS guideline: Decision based on MTUS Chronic Pain Treatment Guidelines Cyclobenzaprine, Medications for chronic pain, Antispasmodics Page(s): 41-42, 60-61, 64-66. Decision based on Non-MTUS Citation Official Disability Guidelines (ODG) Pain, Cyclobenzaprine (Flexeril) and Other Medical Treatment Guidelines UpToDate, Flexeril.

Decision rationale: MTUS Chronic Pain Medical Treatment states for Cyclobenzaprine, "Recommended as an option, using a short course of therapy. The effect is greatest in the first 4 days of treatment, suggesting that shorter courses may be better. (Browning, 2001) Treatment should be brief." The medical documents indicate that patient is far in excess of the initial treatment window and period. Additionally, MTUS outlines, "Relief of pain with the use of medications is generally temporary, and measures of the lasting benefit from this modality should include evaluating the effect of pain relief in relationship to improvements in function and increased activity. Before prescribing any medication for pain, the following should occur: (1) determine the aim of use of the medication; (2) determine the potential benefits and adverse effects; (3) determine the patient's preference. Only one medication should be given at a time, and interventions that are active and passive should remain unchanged at the time of the medication change. A trial should be given for each individual medication. Analgesic medications should show effects within 1 to 3 days, and the analgesic effect of antidepressants should occur within 1 week. A record of pain and function with the medication should be recorded. (Mens, 2005) UpToDate "flexeril" also recommends "Do not use longer than 2-3 weeks". Medical documents do not fully detail the components outlined in the guidelines above and do not establish the need for long term/chronic usage of cyclobenzaprine. ODG states regarding cyclobenzaprine, "Recommended as an option, using a short course of therapy. The addition of cyclobenzaprine to other agents is not recommended." Several other pain medications are being requested, along with cyclobenzaprine, which ODG recommends against. As such, the request for Cyclobenzaprine 10 MG Qty 30 is not medically necessary.

Lidocaine Patch 5 Percent Qty 30: Upheld

Claims Administrator guideline: Decision based on MTUS Chronic Pain Treatment Guidelines Topical.

MAXIMUS guideline: Decision based on MTUS Chronic Pain Treatment Guidelines Lidoderm patches Page(s): 56-57. Decision based on Non-MTUS Citation Official Disability Guidelines (ODG) Pain, Topical analgesics and Other Medical Treatment Guidelines UpToDate.com, Lidocaine (topical).

Decision rationale: Chronic Pain Medical Treatment Guidelines state "Lidoderm is the brand name for a lidocaine patch produced by [REDACTED]. Topical lidocaine may be recommended for localized peripheral pain after there has been evidence of a trial of first-line therapy (tri-cyclic or SNRI anti-depressants or an AED such as gabapentin or Lyrica). This is not

a first-line treatment and is only FDA approved for post-herpetic neuralgia. Further research is needed to recommend this treatment for chronic neuropathic pain disorders other than post-herpetic neuralgia. Formulations that do not involve a dermal-patch system are generally indicated as local anesthetics and anti-pruritics. For more information and references, see Topical analgesics." ODG further details, "Criteria for use of Lidoderm patches: (a) Recommended for a trial if there is evidence of localized pain that is consistent with a neuropathic etiology. (b) There should be evidence of a trial of first-line neuropathy medications (tri-cyclic or SNRI anti-depressants or an AED such as gabapentin or Lyrica). (c) This medication is not generally recommended for treatment of osteoarthritis or treatment of myofascial pain/trigger points. (d) An attempt to determine a neuropathic component of pain should be made if the plan is to apply this medication to areas of pain that are generally secondary to non-neuropathic mechanisms (such as the knee or isolated axial low back pain). One recognized method of testing is the use of the Neuropathic Pain Scale. (e) The area for treatment should be designated as well as number of planned patches and duration for use (number of hours per day). (f) A Trial of patch treatment is recommended for a short-term period (no more than four weeks). (g) It is generally recommended that no other medication changes be made during the trial period. (h) Outcomes should be reported at the end of the trial including improvements in pain and function, and decrease in the use of other medications. If improvements cannot be determined, the medication should be discontinued. (i) Continued outcomes should be intermittently measured and if improvement does not continue, lidocaine patches should be discontinued." Medical documents provided do not indicate that the use would be for post-herpetic neuralgia. Additionally, treatment notes did not detail other first-line therapy used and what the clinical outcomes resulted. As such, the request for Lidocaine Patch 5 Percent Qty 30 is not medically necessary.