

Case Number:	CM15-0104998		
Date Assigned:	06/09/2015	Date of Injury:	11/16/2012
Decision Date:	07/10/2015	UR Denial Date:	05/19/2015
Priority:	Standard	Application Received:	06/01/2015

HOW THE IMR FINAL DETERMINATION WAS MADE

MAXIMUS Federal Services sent the complete case file to an expert reviewer. He/she has no affiliation with the employer, employee, providers or the claims administrator. He/she has been in active clinical practice for more than five years and is currently working at least 24 hours a week in active practice. The expert reviewer was selected based on his/her clinical experience, education, background, and expertise in the same or similar specialties that evaluate and/or treat the medical condition and disputed items/Service. He/she is familiar with governing laws and regulations, including the strength of evidence hierarchy that applies to Independent Medical Review determinations.

The Expert Reviewer has the following credentials:

State(s) of Licensure: North Carolina

Certification(s)/Specialty: Family Practice

CLINICAL CASE SUMMARY

The expert reviewer developed the following clinical case summary based on a review of the case file, including all medical records:

The injured worker is a 66 year old female who sustained an industrial injury on 11/16/2012. There was no mechanism of injury documented. The injured worker was diagnosed with right carpal tunnel syndrome, right middle finger and left thumb stenosing tenosynovitis, cervical myofascial pain, bilateral chondromalacia patella, right shoulder impingement and bilateral elbow collateral ligament tear. The injured worker is status post left carpal tunnel release (no date documented). Treatment to date includes conservative measures; diagnostic testing with recent right elbow, left and right knee magnetic resonance imaging (MRI) in April 2015, physical therapy and medications. According to the treating physician's progress report on April 21, 2015, the injured worker continues to experience lower back and cervical pain, bilateral knee and elbow and right shoulder pain. The injured worker rates her pain level to the various body parts at 5-7/10. Examination demonstrated diffuse left knee tenderness with positive patellofemoral compression test and crepitation with range of motion. The right shoulder noted tenderness with limited range of motion and positive impingement signs. The lumbar spine was essentially unchanged with limited range of motion and positive straight leg raise. There was noted spasm of the lumbar paraspinal muscles. Current medications are listed as Tramadol ER, Naproxen, Cyclobenzaprine, Lidoderm Patches and Pantoprazole. Treatment plan consists of continued requests for bilateral knee arthroscopy, physical therapy for the lower back; follow-up with neurologist and the current request for Cyclobenzaprine and Lidoderm Patches.

IMR ISSUES, DECISIONS AND RATIONALES

The Final Determination was based on decisions for the disputed items/services set forth below:

Cyclobenzaprine 7.5mg #90: Upheld

Claims Administrator guideline: Decision based on MTUS Chronic Pain Treatment Guidelines Muscle Relaxants Page(s): 41-42.

MAXIMUS guideline: Decision based on MTUS Chronic Pain Treatment Guidelines muscle relaxants Page(s): 63-65.

Decision rationale: The California chronic pain medical treatment guidelines section on muscle relaxants states: Recommend non-sedating muscle relaxants with caution as a second-line option for short-term treatment of acute exacerbations in patients with chronic LBP. (Chou, 2007) (Mens, 2005) (Van Tulder, 1998) (van Tulder, 2003) (van Tulder, 2006) (Schnitzer, 2004) (See, 2008) Muscle relaxants may be effective in reducing pain and muscle tension, and increasing mobility. However, in most LBP cases, they show no benefit beyond NSAIDs in pain and overall improvement. Also there is no additional benefit shown in combination with NSAIDs. Efficacy appears to diminish over time, and prolonged use of some medications in this class may lead to dependence. (Homik, 2004) (Chou, 2004) This medication is not intended for long-term use per the California MTUS. The medication has not been prescribed for the flare-up of chronic low back pain. This is not an approved use for the medication. For these reasons, criteria for the use of this medication have not been met. Therefore the request is not medically necessary.

Lidoderm Patches 5%, #2 boxes: Upheld

Claims Administrator guideline: The Claims Administrator did not cite any medical evidence for its decision.

MAXIMUS guideline: Decision based on MTUS Chronic Pain Treatment Guidelines topical lidocaine Page(s): 111-112.

Decision rationale: The California chronic pain medical treatment guidelines section on topical lidocaine states: Lidocaine Indication: Neuropathic pain Recommended for localized peripheral pain after there has been evidence of a trial of first-line therapy (tri-cyclic or SNRI anti-depressants or an AED such as gabapentin or Lyrica). Topical lidocaine, in the formulation of a dermal patch (Lidoderm) has been designated for orphan status by the FDA for neuropathic pain. Lidoderm is also used off-label for diabetic neuropathy. No other commercially approved topical formulations of lidocaine (whether creams, lotions or gels) are indicated for neuropathic pain. Non-dermal patch formulations are generally indicated as local anesthetics and anti-pruritics. Further research is needed to recommend this treatment for chronic neuropathic pain disorders other than post-herpetic neuralgia. Formulations that do not involve a dermal-patch system are generally indicated as local anesthetics and anti-pruritics. In February 2007 the FDA notified consumers and healthcare professionals of the potential hazards of the use of topical lidocaine. Those at particular risk were individuals that applied large amounts of this substance over large areas, left the products on for long periods of time, or used the agent with occlusive dressings. Systemic exposure was highly variable among patients. Only FDA-approved products

are currently recommended. (Argoff, 2006) (Dworkin, 2007) (Khaliq-Cochrane, 2007) (Knotkova, 2007) (Lexi-Comp, 2008) Non-neuropathic pain: Not recommended. There is only one trial that tested 4% lidocaine for treatment of chronic muscle pain. The results showed there was no superiority over placebo. (Scudds, 1995) This medication is recommended for localized peripheral pain. The patient has no documented failure of all first line agents indicated for the treatment of neuropathic pain as outlined above. Therefore, criteria as set forth by the California MTUS as outlined above have not been met and the request is not medically necessary.