

<b>Case Number:</b>	CM15-0108384		
<b>Date Assigned:</b>	06/15/2015	<b>Date of Injury:</b>	09/26/2012
<b>Decision Date:</b>	07/14/2015	<b>UR Denial Date:</b>	06/04/2015
<b>Priority:</b>	Standard	<b>Application Received:</b>	06/05/2015

### HOW THE IMR FINAL DETERMINATION WAS MADE

MAXIMUS Federal Services sent the complete case file to an expert reviewer. He/she has no affiliation with the employer, employee, providers or the claims administrator. He/she has been in active clinical practice for more than five years and is currently working at least 24 hours a week in active practice. The expert reviewer was selected based on his/her clinical experience, education, background, and expertise in the same or similar specialties that evaluate and/or treat the medical condition and disputed items/Service. He/she is familiar with governing laws and regulations, including the strength of evidence hierarchy that applies to Independent Medical Review determinations.

The Expert Reviewer has the following credentials:  
 State(s) of Licensure: North Carolina  
 Certification(s)/Specialty: Family Practice

### CLINICAL CASE SUMMARY

The expert reviewer developed the following clinical case summary based on a review of the case file, including all medical records:

This 58-year-old male sustained an industrial injury to the right shoulder and low back on 9/26/12. Previous treatment included magnetic resonance imaging, physical therapy, injections, extracorporeal shock wave therapy for the left hip and medications. Electromyography of bilateral lower extremities (1/14/15) was normal. The injured worker received 17 physical therapy sessions for the lumbar spine and left hip between 11/14/15 and 3/30/15. No physical exam was included in the documentation submitted for review. Current diagnoses included left hip trochanteric tendinitis and lumbar spine sprain/strain. The treatment plan included physical therapy twice a week for three weeks to the right shoulder and lumbar spine.

### IMR ISSUES, DECISIONS AND RATIONALES

The Final Determination was based on decisions for the disputed items/services set forth below:

**Outpatient physical therapy 2 times a week for 3 weeks to the right shoulder and lumbar spine:** Upheld

**Claims Administrator guideline:** Decision based on MTUS Chronic Pain Treatment Guidelines Physical Medicine.

**MAXIMUS guideline:** Decision based on MTUS Chronic Pain Treatment Guidelines physical medicine Page(s): 98-99.

**Decision rationale:** The California chronic pain medical treatment guidelines section on physical medicine states: Recommended as indicated below. Passive therapy (those treatment modalities that do not require energy expenditure on the part of the patient) can provide short term relief during the early phases of pain treatment and are directed at controlling symptoms such as pain, inflammation and swelling and to improve the rate of healing soft tissue injuries. They can be used sparingly with active therapies to help control swelling, pain and inflammation during the rehabilitation process. Active therapy is based on the philosophy that therapeutic exercise and/or activity are beneficial for restoring flexibility, strength, endurance, function, range of motion, and can alleviate discomfort. Active therapy requires an internal effort by the individual to complete a specific exercise or task. This form of therapy may require supervision from a therapist or medical provider such as verbal, visual and/or tactile instruction(s). Patients are instructed and expected to continue active therapies at home as an extension of the treatment process in order to maintain improvement levels. Home exercise can include exercise with or without mechanical assistance or resistance and functional activities with assistive devices. (Colorado, 2002) (Airaksinen, 2006) Patient-specific hand therapy is very important in reducing swelling, decreasing pain, and improving range of motion in CRPS. (Li, 2005) The use of active treatment modalities (e.g., exercise, education, activity modification) instead of passive treatments is associated with substantially better clinical outcomes. In a large case series of patients with low back pain treated by physical therapists, those adhering to guidelines for active rather than passive treatments incurred fewer treatment visits, cost less, and had less pain and less disability. The overall success rates were 64.7% among those adhering to the active treatment recommendations versus 36.5% for passive treatment. (Fritz, 2007) Physical Medicine Guidelines-Allow for fading of treatment frequency (from up to 3 visits per week to 1 or less), plus active self-directed home Physical Medicine. Myalgia and myositis, unspecified (ICD9 729.1): 9-10 visits over 8 weeks. Neuralgia, neuritis, and radiculitis, unspecified (ICD9 729.2): 8-10 visits over 4 weeks. Reflex sympathetic dystrophy (CRPS) (ICD9 337.2): 24 visits over 16 weeks. The goal of physical therapy is graduation to home therapy after a certain amount of recommended sessions. The patient has already completed a course of physical therapy. The request is in excess of these recommendations per the California MTUS. There is no reason why the patient would not be moved to home therapy after completing the recommended amount of supervised sessions. Therefore, the request is not medically necessary.

**Magnetic resonance imaging (MRI) of the right shoulder:** Upheld

**Claims Administrator guideline:** Decision based on MTUS ACOEM Chapter 9 Shoulder Complaints.

**MAXIMUS guideline:** Decision based on MTUS ACOEM Chapter 9 Shoulder Complaints Page(s): 208.

**Decision rationale:** The ACOEM chapter on shoulder complaints and imaging studies states: Primary criteria for ordering imaging studies are: Emergence of a red flag (e.g., indications of intra-abdominal or cardiac problems presenting as shoulder problems); Physiologic evidence of

tissue insult or neurovascular dysfunction (e.g., cervical root problems presenting as shoulder pain, weakness from a massive rotator cuff tear, or the presence of edema, cyanosis or Raynaud's phenomenon); Failure to progress in a strengthening program intended to avoid surgery. Clarification of the anatomy prior to an invasive procedure (e.g., a full thickness rotator cuff tear not responding to conservative treatment). The provided documentation for review fails to meet the above criteria per the ACOEM. Therefore, the request is not medically necessary.

**Magnetic resonance imaging (MRI) of the lumbar spine:** Upheld

**Claims Administrator guideline:** Decision based on MTUS ACOEM Chapter 12 Low Back Complaints.

**MAXIMUS guideline:** Decision based on MTUS ACOEM Chapter 12 Low Back Complaints  
Page(s): 303-304.

**Decision rationale:** The ACOEM chapter on low back complaints and special diagnostic studies states: Unequivocal objective findings that identify specific nerve compromise on the neurologic examination are sufficient evidence to warrant imaging in patients who do not respond to treatment and who would consider surgery an option. When the neurologic examination is less clear, however, further physiologic evidence of nerve dysfunction should be obtained before ordering an imaging study. Indiscriminant imaging will result in false-positive findings, such as disk bulges, that are not the source of painful symptoms and do not warrant surgery. If physiologic evidence indicates tissue insult or nerve impairment, the practitioner can discuss with a consultant the selection of an imaging test to define a potential cause (magnetic resonance imaging [MRI] for neural or other soft tissue, computed tomography [CT] for bony structures). Relying solely on imaging studies to evaluate the source of low back and related symptoms carries a significant risk of diagnostic confusion (false positive test results) because of the possibility of identifying a finding that was present before symptoms began and therefore has no temporal association with the symptoms. Techniques vary in their abilities to define abnormalities (Table 12-7). Imaging studies should be reserved for cases in which surgery is considered or red-flag diagnoses are being evaluated. Because the overall false-positive rate is 30% for imaging studies in patients over age 30 who do not have symptoms, the risk of diagnostic confusion is great. There is no recorded presence of emerging red flags on the physical exam. There is evidence of nerve compromise on physical exam but there is not mention of consideration for surgery or complete failure of conservative therapy. For these reasons, criteria for imaging as defined above per the ACOEM have not been met. Therefore, the request is not medically necessary.